certain groups of facts are concordant in pointing to a hidden orderliness in the midst of the apparent chaos. Such, for example, are the facts of colour mixture and the facts which demonstrate some definite relationship between colour and luminosity. These, the best established correlations in the whole field of colour vision, are those the importance of which Dr. Edridge-Green minimizes or denies.

In these scattered remarks I have attempted to show that many of Dr. Edridge-Green's statements rest upon very doubtful foundations. Stress has been laid upon the evidence against them because he has tacitly ignored it. Dogmatism may win a Pyrrhic victory, but it is only by exhaustive study, for which few have the time or opportunity, or, it may be feared, the inclination, that a sound judgment of the intricacies of the subject can be arrived at. In controversies such as this, however, it is well to remember John Locke's wise words: "Truth certainly would do well enough if she were once left to shift for herself . . . But if truth makes not her way into the understanding by her own light, she will be but the weaker for any borrowed force violence can add to her."

THE THEORY OF VISION

BY

F. W. EDRIDGE-GREEN, C.B.E.

LONDON.

IN the BRITISH JOURNAL OF OPHTHALMOLOGY for July, Mr. J. H. Parsons has commenced his promised attack on my theories of vision and colour vision.

In the case of such a very difficult subject, which has been worked at by the greatest scientific men of the world, many hesitate to form an opinion because they do not know what can be said on the other side. An attack, therefore, of this kind which might otherwise prejudice those only partially acquainted with the subject, greatly strengthens my position when it can be shown that the arguments, many very old, are mostly based on misstatements.

In this article he has dealt with the retinal portion of my theory in which I assume that the rods are not perceptive elements, but are concerned with the formation and distribution of the visual purple. Mr. Parsons does not appear to have read my book very carefully, as nearly all the points he raises have been answered there. He states that some of the evidence in favour of the rods being perceptive elements is as follows: *Comparative Anatomy*: He states that there are certain animals, as, for instance, the tortoise which have only cones and others possess only rods. This I have

already answered in my book, page 54: "It was formerly stated that a tortoise possesses only cones, but on examining the retina of a tortoise I found the rods and cones as definitely marked and distinct from each other as in man." This argument therefore is valueless.

Dr. Lindsay Johnson, whose work has not received the recognition in this country it deserves, informed me that in no animal had he failed to find a visual substance corresponding to the visual purple. I have spent many evenings with him examining his specimens. He also told me that he had discovered a large duct leading from the macular region to the lymphatic spaces of the optic nerve. Though this is strongly in support of my theory, and is probably true, I have not previously mentioned it as I was not satisfied that I saw it in the specimens shown.

The next portion of his attack has also been dealt with in my book, page 50. "Many physiologists have tried to assign different functions to the rods and cones, but all these theories have failed, because all the functions which were said to be the exclusive property of the rods have been found only gradually diminished in the fovea. For instance, von Tschermak, Hering, Hess, Garten, and I have found the Purkinje phenomenon, the variation in optical white equations by a state of light and dark adaptation, the colourless interval for spectral lights of increasing intensity, the varying phases of the after image, in the fovea, only gradually diminished. The complete absence of any qualitative change between the foveal and extra foveal regions is a very important fact in support of the hypothesis that the visual purple is the visual substance."

Mr. Parsons has such a biased style of writing. He writes: "Although there is some difference of opinion on the point, most competent observers using the most refined methods agree that Purkinje's phenomenon is absent at the fovea." An explanation of this difference of opinion is given in my book, page 78. Results depend upon the intensity of the light employed; this demonstration was given before the Physiological Society and no one failed to see the Purkinje phenomenon with the fovea. Why is Mr. Parsons so reluctant to observe these facts for himself? During the many years that I was doing nothing but original research on vision and colour vision at the Physiological Institute, at University College, why did he not visit my laboratory, as I have often offered to show him the facts? Neither has he ever been present at any of the numerous demonstrations that I have given at the Physiological Society.

As I have pointed out, the Purkinje phenomenon is a photo-chemical phenomenon which is found with other photo-chemical substances, and therefore a duplex mechanism is not needed to

explain it. Mr. Parsons himself admits that Dr. Houstoun's work supports my view of a single mechanism. Mr. Parsons insinuates that those who have found the Purkinje phenomenon in the fovea, for instance, Hering, Hess, Garten, and others, are not competent observers, and do not use refined methods. This style of writing cannot be too severely condemned. I here challenge him to point out a defect in the method which I have devised and which is described in my book. How can Mr. Parsons possibly judge by reading a paper whether a particular method is the best. One advantage of attending International Congresses is that you learn from other men their opinions of the accuracy of different observers. I found that those of Hess were regarded with particular respect, and in this I quite agree. I have never found him wrong yet. His work on the recurrent image in the fovea is a particularly pretty piece of work. He shows that the recurrent image (stated by others to be absent in the fovea) appears bent and was therefore retarded. Both von Kries and Hess demonstrated their methods to me and it was quite obvious that retardation would explain von Kries's results.

Mr. Parsons states with regard to my discovery of channels in the retina, "The anatomical evidence in favour of such channels is of little value." Why should it be? Anatomical points must be settled by anatomical evidence. It may be as well to relate how I discovered them. I was examining the outer side of a monkey's retina with Mr. Devereux Marshall, a pathologist of the eye of considerable experience. I pointed out the canals to him and asked him if by any chance they could be artefacts; his reply was, "Most emphatically no." I then said we ought to be able to see them entoptically with our own eyes, and I there and then devised a method and saw in whirling currents the exact pattern of the canals of the monkey's retina. Anything more conclusive than this could hardly be imagined, and the fact of the canals explains so much of the work of Purkinje.

I would suggest to anyone who on reading my book finds that he cannot see some of the phenomena, that he asks another observer, but on no account tells him what he is supposed to see. I find that women are particularly good at seeing these phenomena if they are not biased one way or the other.

In order to ascertain certain facts and test certain methods, I have visited Paris, Utrecht, Groningen, Amsterdam, Upsala, Berlin, Leipzig, Freiburg, Vienna, Bern, Budapest, etc., whereas Mr. Parsons would not even cross the road to see my facts, and yet tries to pose as an authority.

I must here repeat that it is absolutely necessary to see experiments before an opinion can be given upon them. In the time occupied by a research recorded in a few lines in my book, I could have written a voluminous compilation.

Some years ago one of our ablest physiologists advised me to go to Prof. Asher, of Bern, whom he considered the best qualified man in Europe to give an opinion on the theory of vision. I did so, and discussed the subject in minute detail. The following is an extract of a letter received from him March 11, 1920: "Your theory as to rods and cones, and their position in a photo-chemical theory seems to me very suggestive. I have always found a great difficulty in reconciling the facts of parallelism between the physical properties of the visual purple and the vision during dark adaptation with the physiological and psychological improbability of the Parinaud-Kries theory. Your views seem a way to avoid the difficulties."

The most conclusive fact against the duplicity theory is that when a colourless spectrum has been obtained, with further dark adaptation colour re-appears; in fact Burch stated that with complete dark adaptation there was no photo-chromatic interval. Spectral coloured light appeared as coloured immediately it was visible as light.

THREE CASES OF TOTAL REMOVAL OF IRIS

BY

D. J. WOOD,

CAPE TOWN, SOUTH AFRICA.

P. A., a fitter, came to me some years ago saying that he had been struck in the left eye by a piece of iron. The A.C. was full of blood, and there was a fine cut, not more than $\frac{1}{8}$ inch long, just outside the cornea. The presumption was that the piece of iron was in the eye. A skiagram was negative, but I knew that the chip was small and thin from the very small hole. Accordingly I tried with the only magnet I then possessed, a small one, to get the fragment out. Using a fine terminal, introduced through the wound, I made three attempts, but failed. I decided to do no more, and expected the eye to do badly. To my surprise it remained quiet, and a week later the blood had more than half gone, but no iris was visible. It had in fact been entirely removed through the tiny wound and there was no foreign body in the eye at all. The most unexpected thing was that when I saw him some months later I found that I had escaped damaging the lens, and that even the vitreous was almost clear. Vision was $\frac{6}{9}$. There was no trace of iris, and the whole lens and the tips of the ciliary processes could be seen all round.

A. B. M., a carpenter, came to me two years ago with the history that the night previously he had fallen against some wire netting and hurt his eye. It was very red, the A.C. was full of blood, and there was a scar above with iris pigment in it. I