PERSPECTIVE

The emerging roles of topical non-steroidal anti-inflammatory agents in ophthalmology

Peter Koay

Non-steroidal anti-inflammatory drugs (NSAIDs) are a heterogeneous group of compounds which in chemical terms belong to different structural classes. By definition, NSAIDs do not include a steroid nucleus derived biosynthetically from cholesterol in their chemical structures. Although the term NSAIDs can be expanded to include other classes of compounds such as immunosuppressives and cytokine modulators,¹ this discussion will be limited to commercially available topical ophthalmic drugs generally regarded as NSAIDs. Topical ophthalmic drugs under discussion are of the following chemical classes: indoles, phenylalkanoic acids, phenylacetic acids, and pyrazolones. The main topical ophthalmic drug of the indole chemical class is indomethacin. This was the first topically administered NSAID to be used widely in ophthalmic clinical studies. Phenylalkanoic acids are water soluble and are readily formulated as topical ophthalmic solutions such as flurbiprofen 0.03% (Ocufen), ketorolac tromethamine 0.5% (Acular), and suprofen 1% (Profenal). The main topical ophthalmic phenylacetic acid derivative is diclofenac 1% (Voltarol) (Table 1). NSAIDs are also known as cyclo-oxygenase inhibitors based on their mode of action, and are important mediators or modulators of ocular inflammatory reactions.²

Arachidonic acid metabolites in ocular inflammation

Activation of phospholipase A₂, following tissue insults, breaks down cell membrane phospholipids to arachidonic acid. This forms the substrate for further reactions mainly by the cyclo-oxygenase and the lipoxygenase pathways (Fig 1). Other pathways, such as the cytochrome P₄₅₀ pathway, are also involved in the production of biologically active lipids. The resulting arachidonic acid metabolites, along with other chemical mediators, interact to cause inflammation. Although arachidonic acid metabolites are associated with pro-inflammatory actions, there are a few studies to suggest that some of these metabolites also have anti-inflammatory actions.²

The main products of the cyclo-oxygenase (CO) pathway are prostaglandins (PGs). In the rabbit model, PGs

appear in the aqueous after ocular paracentesis, thermal or mechanical injury which cause a rise in both intraocular pressure and protein concentration in the aqueous humour. The principal source of PGs in the aqueous humour is probably derived from the iris-ciliary body. PGs cause miosis, increase in vascular permeability of the blood-ocular barriers, and changes in intraocular pressure. Furthermore, PGs are known to possess chemokinetic activity and can serve as mediators in both humoral and cellular phases of the inflammatory response.²

The main products of the lipoxygenase (LO) pathway are leukotrienes (LT) such as leukotriene B₄ and peptidoleukotrienes. Leukotriene B₄ causes aggregation of polymorphonuclear cells (PMN), PMN chemotaxis in vivo, PMN chemokinesis, exudation of plasma, translocation of calcium, and stimulation of phospholipase A₂. Peptidoleukotrienes result in bronchoconstriction, secretion of mucus, plasma leakage from post capillary venules, vasoconstriction of arterioles, and miosis.²

Inflammation is complex involving a large number of mediators and products. NSAIDs inhibit the cyclo-oxygenase enzyme (Fig 1) but not the lipoxygenase enzyme. Thus, NSAIDs inhibit the biosynthesis of PGs but not LTs. Inhibition of PG biosynthesis inhibits intraoperative miosis during cataract surgery, reduces the vascular permeability of the blood-ocular barrier, and modifies inflammation. It should be noted, however, that NSAIDs are not capable of inhibiting the actions of PGs once formed.³

Glucocorticoids inhibit the phospholipase A₂ enzyme (Fig 1), thus inhibiting the biosynthesis of arachidonic acid itself. This results in the inhibition of the biosynthesis of both PGs and LTs. Other effects of glucocorticoids not directly involved in the arachidonic pathway include: effects on white cells (inhibit migration of macrophages, neutropenia, lymphocytopenia, eosinopenia, monocytopenia, neutrophilic leucocytosis); inhibition of degranulation of neutrophils, mast cells, basophils; lysosome stabilisation; suppression of lymphokines; reduction of capillary permeability; and suppression of vasodilatation.⁴

Ocular penetration and distribution of topical ophthalmic NSAIDs

Indomethacin is soluble in alcohol but not in water and was initially formulated for ophthalmic clinical studies using a sesame oil vehicle which proved to be irritating. An aqueous suspension of indomethacin 1% is now available and shaking the bottle is recommended.⁵ Indomethacin has been shown to significantly decrease levels of prostaglandin E₂ in the human cornea.⁶ In an animal model, topical 1% indomethacin suspension resulted in high concentrations in the cornea, but low concentrations in the iris, the ciliary body, the lens, and the choroid.⁷

Table 1 Commercially available topical ophthalmic non-steroidal anti-inflammatory drugs

Chemical class	Generic name	Trade name (Manufacturer)
Indoles	Indomethacin	Indomethacin 0.5% and 1% in castor oil eyedrops (Moorfields Eye Hospital) Indocid ophthalmic suspension (Merck, Sharp & Dohme)
Phenylalkanoic acids	Flurbiprofen Ketorolac Suprofen	Ocufen (Allergan) Acular (Allergan) Profenal (Alcon)
Phenylacetic acids	Diclofenac	Voltaren/Voltarol (Ciba Vision Ophthalmics)
Pyrazolones	Oxyphenbutazone	Tanderil 10% (Geigy)

Figure 1 A simplified schematic representation of the enzymatic pathways of arachidonic acid metabolism. Site of action of glucocorticoids (A) and NSAIDs (B).

Topical administration of indomethacin appears to produce better penetration into the eye than oral administration.⁸

Commercially available topical NSAIDs generally have good ocular penetration. Topical use of flurbiprofen or ketorolac will result in the distribution of the drug throughout the conjunctiva, cornea, aqueous, sclera, iris, ciliary body, and choroid-retina in rabbits.⁹⁻¹⁰ Diclofenac applied topically readily penetrates the cornea and into the aqueous.¹¹ Systemic absorption of topical NSAIDs is minimal as demonstrated by the plasma levels of diclofenac following ocular instillation (two drops of diclofenac to each eye) being below the limit of quantification (10 ng/ml) over a 4 hour period.¹²

Potential advantages of topical NSAIDs over corticosteroids

The main advantage of topical NSAIDs is that the undesirable effects of topical corticosteroids are avoided. These include decreased immunological response to infection,¹³ cataract formation,¹⁴ and steroid-induced raised intraocular pressure (IOP).¹⁵ A study of steroid responsive patients revealed that flurbiprofen has no effect upon the elevation of IOP compared with dexamethasone.¹⁶ Topical diclofenac 0.1% had no significant effect on IOP following cataract surgery compared with placebo.¹⁷ Furthermore, the use of NSAIDs such as flurbiprofen sodium or diclofenac sodium does not exacerbate acute herpetic keratitis or prolong viral shedding.¹⁸

In animal models, low therapeutic doses of cortisone have been noted to have no significant effect on epithelial regeneration. However, large doses of cortisone retard epithelial and endothelial regeneration. Large doses also

retard the formation of fibrinous coagulum, cellular infiltration, and fibroblastic repair.¹⁹ Srinivasan, in a series of experiments involving mechanical corneal injury, demonstrated in a rabbit model that corticosteroids (prednisolone acetate 1%, dexamethasone 0.1%, and fluorometholone 1%) appear to inhibit re-epithelialisation by mainly conjunctival epithelial cells following complete epithelial denudation. These corticosteroids do not inhibit re-epithelialisation by mainly corneal epithelial cells after partial epithelial denudation. NSAIDs (indomethacin 0.5% and flurbiprofen 0.01%) in contrast, do not inhibit the re-epithelialisation by either corneal or conjunctival epithelial cells.²⁰ Topical diclofenac sodium 0.1% does not delay re-epithelialisation following excimer laser treatment in rabbit corneas.²¹ However, Hersh *et al* in their study of epithelial scrape wound healing in rabbits demonstrated a delay in epithelial healing for the first 3 days when NSAIDs (flurbiprofen sodium 0.03%, diclofenac sodium 0.1%) or steroids (prednisolone acetate 1%) were used compared with placebo vehicle.²²

Corticosteroids can result in decreased stromal corneal wound healing.²³ One rabbit study involving full thickness corneal wounds demonstrated that treatment with flurbiprofen sodium or diclofenac sodium actually resulted in stronger scars than untreated controls. Corneal wounds treated with prednisolone acetate resulted in the weakest scar.²⁴ In another rabbit study, the effects of equipotent anti-inflammatory doses of flurbiprofen and prednisolone on the wound bursting pressure of 4 mm corneal incisions were compared. The results of this study suggest that flurbiprofen and prednisolone do not differ in their effects on corneal wound healing.²⁵ Topical flurbiprofen 0.03% and prednisolone acetate 1% show similar astigmatic decay

curves following 6.5 mm scleral pocket cataract surgery with no clinical or statistical difference in the astigmatic results 4 months after surgery.²⁶ Although the advantage of NSAIDs over corticosteroids regarding corneal epithelial healing and wound strength is far from certain, the findings might favour the use of topical NSAIDs.

Use of topical NSAIDs in reducing postoperative inflammation

Postoperative inflammation may be evaluated by clinical biomicroscopy examination and by anterior fluorophotometry. The latter assumes a correlation between inflammation and breakdown of the blood-aqueous barrier as measured by accumulation of fluorescein in the anterior chamber. Fluorophotometry allows for quantification, better reproducibility, and greater objectivity in the evaluation of postoperative inflammation.

Topical ketorolac tromethamine 0.5% is effective in reducing clinical and anterior fluorophotometric postoperative inflammation when compared with placebo^{27 28} and dexamethasone 0.1%^{29 30} in extracapsular cataract extraction. Topical diclofenac sodium 0.1% was favoured over placebo at 4, 8, and 15 days after cataract surgery for clinical assessments of anterior chamber cells, flare, conjunctival erythema, and ciliary flush.³¹ Kraff *et al.*³² in a study without concomitant steroidal anti-inflammatory agents, showed significantly better 'summed flare and anterior chamber cells score' at 2 to 5 days and 7 to 9 days after cataract surgery in patients receiving diclofenac sodium 0.1% than in those on placebos. Brennan *et al.*³³ found no difference in postoperative inflammation (evaluated by slit-lamp assessment of cell and flare, and objectively by using the Kowa FC-1000 laser cell and flare meter) between diclofenac sodium 0.1% (Voltaren) and prednisolone acetate 1% (Pred forte) in patients undergoing phacoemulsification and posterior chamber intraocular lens implant. Furthermore, topical diclofenac sodium 0.1% has been shown to be superior to prednisolone sodium phosphate 1% in reducing anterior fluorophotometric postoperative inflammation.³⁴ The effects of NSAIDs in postoperative inflammation following laser trabeculoplasty³⁵⁻³⁷ and cyclocryotherapy³⁸ are under investigation.

Effectiveness of topical NSAIDs on symptoms of allergic conjunctivitis

Seasonal allergic conjunctivitis is characterised by itching, burning, tearing, and lid oedema in response to exposure to airborne allergens. Although histamine release is recognised as a cause of itching, other mediators such as prostaglandins, leukotrienes, and platelet activating factors are involved. NSAIDs act by interrupting the synthesis of prostaglandins which may lower the threshold of histamine induced itching and influence the nerve responsiveness to itching. In a double masked randomised trial in patients with seasonal allergic conjunctivitis, although placebo treated eyes experienced some relief due to the dilution of antigens, ketorolac tromethamine 0.5% has been shown to significantly reduce the severity of the associated ocular itching.³⁹ A study comparing the effects of diclofenac sodium 0.1% and ketorolac tromethamine 0.5% showed similar efficacy in treating ocular itching and bulbar injection.⁴⁰ Diclofenac 0.1% is effective in the treatment of episcleritis, hay fever conjunctivitis, phlyctenular conjunctivitis, and corneal limbal ulcers, replacing or in conjunction with corticosteroids.⁴¹ Oxyphenbutazone (Tanderil eye ointment)⁴² has been shown to be ineffective for allergic conjunctivitis and has the added problem of some patients being intolerant of the ointment base.^{43 44}

Inhibition of intraoperative miosis with topical NSAIDs

The use of preoperative topical parasympatholytic and sympathomimetic agents with intraoperative sympathomimetics⁴⁵ is effective in keeping the pupil dilated during cataract surgery. Topical flurbiprofen 0.03%, suprofen 1%, diclofenac sodium 0.1%, and indomethacin 1% appear to have a small statistical effect on inhibiting intraoperative miosis.^{4 46-48} This may or may not be clinically significant depending on surgical technique and viscoelastic technique in use.

Prevention and treatment of pseudophakic or aphakic cystoid macular oedema with topical NSAIDs

Although the exact aetiology of cystoid macular oedema (CMO) has not been established, ocular inflammation involving prostaglandins has been implicated.^{49 50} Prostaglandins cause increased capillary leakage, breakdown of the blood-aqueous barrier, and perifoveal capillary dilatation. Other possible factors affecting the incidence of CMO include the type of surgery (for example, phacoemulsification, intracapsular or extracapsular cataract surgery), primary capsulotomy, phototoxicity, and the role of other autocooids (for example, bradykinin, serotonin).⁵¹ The natural history of spontaneous resolution in CMO complicates the interpretation of the efficacy of modifying agents. In the study of CMO, it is useful to distinguish angiographic CMO from clinically significant CMO that includes a reduction of the measurable variables of vision. Angiographic evidence of CMO occurs in as many as 50-70% of postoperative cataract patients with only approximately 1% of patients suffering significant loss of vision.⁵²

Prophylactic use of topical indomethacin 1% with concurrent corticosteroids is effective in reducing angiographic CMO in the short term (within 7 months) but did not, in early studies (with one exception⁵³), demonstrate a beneficial effect on visual acuity.^{54 55} The use of topical diclofenac five times daily until discharge and subsequently three times daily for 6 months with concurrent corticosteroids has been shown to significantly reduce the incidence of angiographic CMO in intracapsular cataract surgery.⁵⁶ Treatment with topical ketorolac tromethamine 0.5% without concurrent steroids before and after cataract surgery is effective in reducing postoperative angiographic CMO but produces no significant difference in the visual acuity results.⁵⁷

One recent study demonstrated that flurbiprofen 0.03% and indomethacin 1% (with concurrent corticosteroids in approximately 88% of the patients in each group) reduces the incidence of both angiographic CMO and clinical CMO (defined as reduced Snellen visual acuity and contrast sensitivity) from day 21 to day 60 after extracapsular cataract surgery. These topical NSAIDs were used four times daily 2 days before surgery, half hourly for 2 hours before surgery, and four times daily after surgery for 3 months.⁵⁸ The use of flurbiprofen 0.03% appears to significantly improve contrast sensitivity with or without CMO at 12 cycles per degree. In the presence of CMO, patients using flurbiprofen generally had higher contrast sensitivity scores than patients using the placebo vehicle.⁵⁹

Flach *et al.*⁶⁰ showed that there was improvement in distance visual acuity in patients with angiographically proved chronic CMO (visual acuity less than or equal to 20/40 for 6 months) treated with ketorolac tromethamine 0.5% when compared with placebo. In a study involving 30 eyes with chronic CMO after cataract surgery, Peterson *et al.*

were able to demonstrate 80% improvement of three lines or more when using topical indomethacin 1%.⁶¹ Furthermore, 53% of these patients demonstrated an 'on/off' phenomenon induced by the initiation and cessation of treatment documented by visual acuity measurements and fluorescein angiography. Treatment of confirmed symptomatic CMO with fenoprofen 1% resulted in visual improvement that was not statistically significant.⁶²

In conclusion, topical NSAIDs help to maintain the blood-aqueous barrier after ocular surgery. Although the number of cases of clinically significant CMO may be small, the consequences for the patient can be profound.⁶³ Topical NSAIDs may reduce this risk and improve the visual outcome.

Analgesic effects of topical NSAIDs

Following excimer photorefractive keratectomy (PRK) patients may experience significant ocular pain and discomfort until corneal re-epithelialisation. Hyperaesthesia may occur following excimer laser ablation.⁶⁴ Studies on topical diclofenac sodium show that it reduces corneal sensitivity,⁶⁵ has an anaesthetic effect,⁶⁶ and decreases discomfort after excimer laser treatment.⁶⁷ Sher *et al*⁶⁸ have reported, in a clinical series of 16 diclofenac treated patients and 16 placebo patients, that topical diclofenac appears to significantly reduce the ocular pain following excimer PRK. Diclofenac treated patients rarely experience the early post-PRK peak in pain and have less pain overall. In addition, such patients have significantly less post-laser photophobia, burning, and stinging symptoms. Significantly fewer patients prescribed topical diclofenac require oral narcotics. Topical ketorolac tromethamine 0.5% (Acular) has also been reported as an analgesic.^{69–71} There is no statistical difference in the effectiveness of topical ketorolac compared with topical diclofenac on discomfort following radial keratotomy.⁷² Painful ocular conditions such as episcleritis and corneal limbal ulcers may benefit not only from the corticosteroid sparing effect of diclofenac sodium 0.1% but also from its analgesic properties.³⁰ Indomethacin does not appear to have any significant analgesic activity.⁷³

Effects of topical NSAIDs following excimer laser treatment

The cornea has a lesser capacity to synthesise cyclo-oxygenase and lipoxygenase products from arachidonic acid than the conjunctiva and anterior uvea.⁷⁴ Nevertheless, following the application of 193 nm excimer laser on rabbit corneas, prostaglandin E₂ (PGE₂) production (mediated by the cyclo-oxygenase pathway) has been noted to be rapid and sustained with leucocyte infiltration in the cornea without significant change in the levels of leukotriene B₄ (mediated by the lipoxygenase pathway).⁷⁵ In another experiment, 193 nm excimer laser irradiation caused a greater increase in PGE₂ production in rabbit corneal tissue (8.6-fold increase in PGE₂ compared with unoperated cornea) than keratectomy with a micro-keratome to the same depth.⁷⁶ These observations suggest that the inflammatory responses following excimer laser irradiation are mediated primarily by prostaglandins.

In mammalian cell cultures, ultraviolet A light stimulates cyclo-oxygenase metabolism and produces high levels of PGE₂⁷⁷ which acts as a chemoattractant for polymorphonuclear leucocytes (PMNs).⁷⁸ Topical diclofenac sodium 0.1% was shown to reduce the local release of PGE₂ and the migration of PMNs in the rabbit cornea after 193 nm excimer laser ablation.⁷⁹ Diclofenac was also

shown to inhibit the chemokinetic activity (speed of PMN migration) of human PMNs elicited *in vitro*.⁸⁰ In another study, postoperative topical diclofenac sodium treatment resulted in a significant decrease in prostaglandin E₂ levels and a significant increase in corneal leucocytes at 10 hours. Treatment with fluorometholone, a corticosteroid, did not significantly alter prostaglandin E₂ levels but depressed leucocyte ingress.⁵⁸ PMNs in tear fluid of rabbits subjected to corneal de-epithelialisation are inhibited by topically applied flurbiprofen 0.01%, indomethacin 0.5%, and aspirin 0.5%.⁸¹

PMNs play a central role in the inflammatory process. In a study involving experimental keratitis, the ability of a NSAID (suprofen) and a corticosteroid (prednisolone acetate) to suppress PMN invasion of the rabbit cornea was evaluated. Suprofen therapy initiated immediately after induction of inflammation was ineffective and was only efficacious if therapy was actually started 48 hours before corneal injury. In contrast, prednisolone acetate therapy was effective when initiated immediately after induction of inflammation. Prednisolone acetate therapy, in addition, produced a marked increase in therapeutic effect when started 48 hours before the insult. Combined treatment of suprofen and prednisolone acetate was significantly more effective than with either drug alone.⁸² In a small clinical trial, concurrent use of corticosteroids and NSAIDs appeared to reduce myopic regression for 1 year after PRK and the effect was more pronounced with flurbiprofen sodium than diclofenac sodium.⁸³

The use of NSAIDs without corticosteroids following excimer PRK has resulted in cases of corneal infiltrates. Theoretical inhibition of the cyclo-oxygenase pathway for prostaglandin generation may result in increased formation of leukotrienes recognised as chemotactic for PMNs^{84–86} and this may encourage PMN infiltrate formation in the earlier healing stage. It has been suggested that patients taking NSAIDs alone may experience more corneal haze and regression of effect.⁸⁷ However, in a study comparing the use of a topical NSAID (diclofenac sodium 0.1%) and corticosteroids (dexamethasone 0.1%) 20 consecutive eyes of 10 patients (4 to 9 dioptres myopia intended correction) showed no statistical difference in haze or regression effect.⁸⁸ Experimental work on rabbits treated with 193 nm excimer laser myopic PRK to correct 5 dioptres, showed less corneal haze in the group receiving post-laser topical diclofenac 0.1% and fluorometholone 0.1%.⁸⁹ The usefulness of specific NSAIDs in excimer PRK with regard to corneal haze and regression is still uncertain.

Adverse reactions, precautions, and interactions of topical NSAIDs

Frequently reported adverse reactions of NSAIDs include transient burning, stinging, and minor signs of ocular irritation on instillation. Hypersensitivity reactions with itching, reddening, photosensitivity, and keratitis punctata may occur. These reactions may be seen especially in patients with asthma,⁹⁰ urticaria, or acute rhinitis and patients who have symptoms precipitated by drugs with prostaglandin synthetase inhibiting activity (for example, acetylsalicylic acid).^{91–92} There have been reports that topical flurbiprofen may cause an increase in bleeding tendency of ocular tissues in conjunction with surgery.⁸⁸ Although there have been no reported adverse events, there is a theoretical possibility that patients receiving other medications which prolong bleeding time, or with known haemostatic defects, may experience exacerbation of bleeding with topical flurbiprofen and diclofenac.^{87–88} It is advisable that patients with a history

of herpes simplex keratitis using NSAIDs should be monitored. As there are no adequate studies in pregnant women, topical NSAIDs should be avoided in these patients.

Prostaglandins such as the topical PGF_{2α} isopropylester series can cause reductions of intraocular pressure by enhancing uveoscleral outflow in laboratory and clinical trials.⁹³⁻⁹⁶ NSAIDs which block prostaglandin synthesis may thus possibly interfere with the hypotensive effects of prostaglandins. In a rabbit model, topical flurbiprofen 0.03% can interfere with the ocular hypotensive efficacy of topical epinephrine (adrenaline) 1% but not with the hypotensive efficacy of topical carbonic anhydrase inhibitors.⁹⁷⁻⁹⁸ Further clinical studies are required.

The future of topical NSAIDs

In cataract surgery, a single topical agent with multiple effects is potentially attractive for routine use. NSAIDs can inhibit intraoperative miosis and prostaglandin release, act as a postoperative anti-inflammatory agent with analgesic properties, and reduce the risk of cystoid macula oedema. However, the widespread use of NSAIDs is currently hindered by their relatively high cost, especially with unit dose packaging and prolonged treatment regimens. This situation will hopefully improve as more NSAIDs become commercially available.

P KOAY

Darlington Memorial Hospital,
Darlington DL3 6HX

- Rosenbaum JT. Towards cytokine insight in sight. *Br J Ophthalmol* 1995; 79: 970-1.
- Bhattacharjee P. The role of arachidonate metabolites in ocular inflammation. *The ocular effects of prostaglandins and other eicosanoids*. New York: Alan R Liss, 1989: 211-27.
- Van Huse H. Topical treatment of anterior ocular diseases with diclofenac Na eye drops. *Klin Monatsbl Augenheilkd* 1986; 188: 615-9.
- Flach AJ. Cyclo-oxygenase inhibitors in ophthalmology. *Surv Ophthalmol* 1992; 36: 259-84.
- Srobel J. Concentration of locally applied indomethacin eye drops for the inhibition of prostaglandin synthesis in the eye. *Ophthalmology* 1991; 202: 43-7.
- Frucht-Pery J, Zauberman H. The effect of indomethacin on prostaglandin E₂ in human cornea and conjunctiva. *Arch Ophthalmol* 1992; 110: 343-5.
- Spies BM, Mathis GA, Franson KL, Leber A. Kinetics of uptake and effects of topical indomethacin application on protein concentration in the aqueous humor of dogs. *Am J Vet Res* 1991; 52: 1159-63.
- Sanders DR, Kraff MC, Goldstick B. Aqueous penetration of oral and topical indomethacin in humans. *Arch Ophthalmol* 1983; 101: 1614-6.
- Anderson JA, Chen CC, Vita JB. Disposition of topical flurbiprofen in normal and aphakic rabbit eyes. *Arch Ophthalmol* 1982; 100: 642-5.
- Ling TL, Comb DL. Ocular bioavailability and tissue distribution of ketorolac tromethamine in rabbits. *J Pharm Sci* 1987; 76: 289-94.
- Goa KL, Chriss P. Ocular diclofenac: a review of its pharmacology and clinical use in cataract surgery and potential in other inflammatory ocular conditions. *Drugs Aging* 1992; 2: 478-9.
- Voltaire ophthalmic prescribing information 11/93, CS665633.
- Fraser-Smith EB, Matthews TR. Effect of ketorolac on *Pseudomonas aeruginosa* ocular infection in rabbits. *J Ocular Pharmacol* 1988; 4: 101-9.
- Kana JS, Wiederholt M. Corticosteroid and the active ion transport of the isolated human lens. *Ann Ophthalmol* 1984; 16: 1034-9.
- Hic J, Gigon S, Leuenberger PM. Comparison of the anti-inflammatory effect of dexamethasone and diclofenac ophthalmic preparations. *Klin Monatsbl Augenheilkd* 1984; 184: 494-8.
- Geiser DK, Hodapp E, Goldberg I, Kass MA, Becker B. Flurbiprofen and intraocular pressure. *Ann Ophthalmol* 1981; 13: 831-3.
- Strelow SA, Sherwood MB, Broncato LJB, Napier A, Driebe WT, Guy JR, et al. The effect of diclofenac sodium ophthalmic solution on intraocular pressure following cataract extraction. *Ophthalmic Surg* 1992; 23: 170-5.
- Trousdale MD, Barlow WE, McGuigan LJB. Assessment of diclofenac on herpes keratitis in rabbit eyes. *Arch Ophthalmol* 1989; 107: 1664-6.
- Ashton N, Cook C. Effect of cortisone on healing of corneal wounds. *Br J Ophthalmol* 1951; 35: 708.
- Srinivasan BD. Corneal reepithelialization and anti-inflammatory agents [Review]. *Trans Am Ophthalmol Soc* 1982; 80: 758-822.
- Loya N, Bassage S, Vyas S, del Cerro M, Park SB, Aquavella JV. Topical diclofenac following excimer laser: effect on corneal sensitivity and wound healing in rabbits. *J Refract Corneal Surg* 1994; 10: 423-7.
- Hersh PS, Rice BA, Baer JC, Wells PA. Topical nonsteroidal agents and corneal wound healing. *Arch Ophthalmol* 1990; 108: 577-83.
- Henahan JF. Wounds destabilized by steroids more than by incision sutures. *Ophthalmology Times* 1992; 8.
- McCarey BE, Napalkov JA, Phippen PA, Reaves TA. Corneal wound healing strength with topical anti-inflammatory drugs. *Invest Ophthalmol Vis Sci* 1993; 34: 1321.
- Miller D, Gruenberg P, Miller R, Bergamini MVW. Topical flurbiprofen or prednisolone: effect on corneal wound healing in rabbits. *Arch Ophthalmol* 1981; 99: 681-2.
- Masket S. Comparison of the effect of topical corticosteroid and non-steroidals on postoperative corneal astigmatism. *J Cataract Refract Surg* 1990; 16: 715-8.
- Flach AJ, Lavelle CJ, Olander KW, Retzlaff JA, Sorenson LW. The effect of ketorolac tromethamine solution 0.5% in reducing postsurgical inflammation after cataract extraction and intraocular lens implantation. *Ophthalmology* 1988; 95: 1279-84.
- Flach AJ, Graham J, Kruger L, Stegman RC, Tanenbaum L. Quantitative assessment of postsurgical breakdown of the blood aqueous barrier following administration of ketorolac tromethamine solution: a double-masked, paired comparison with vehicle-placebo solution study. *Arch Ophthalmol* 1988; 106: 344-7.
- Flach AJ, Jaffe NS, Akers WA. The effect of ketorolac tromethamine in reducing postoperative inflammation: a double-masked parallel comparison with dexamethasone. *Ann Ophthalmol* 1989; 21: 407-11.
- Flach AJ, Kraff MC, Sanders DR, Tanenbaum L. The quantitative effect of 0.5% ketorolac tromethamine solution and dexamethasone sodium phosphate solution on the post-surgical blood-ocular barrier. *Arch Ophthalmol* 1988; 106: 480-3.
- Vickers FF, McGuigan LJB, Ford C, Wysowsky H, Mellars K, Koester J, et al. The effect of diclofenac sodium ophthalmic on the treatment of post-operative inflammation. *Invest Ophthalmol Vis Sci* 1991; 32: 793.
- Kraff MC, Martin RG, Nuemann AC, Weinstein AJ. Efficacy of diclofenac sodium ophthalmic solution versus placebo in reducing inflammation following cataract extraction and posterior chamber lens implantation. *J Cataract Refract Surg* 1994; 20: 138-44.
- Brennan KM, Brown RM, Roberts CW. A comparison of topical non-steroidal anti-inflammatory drugs to steroids for control of post cataract inflammation. *Insight* 1993; 18: 8-9, 11.
- Kraff MC, Sanders DR, McGuigan L, Raanan MG. Inhibition of blood-ocular humor barrier breakdown with diclofenac. A fluorophotometric study. *Arch Ophthalmol* 1990; 108: 380-3.
- Hotchkiss ML, Robin AL, Pollack IP, Quigley HA. Nonsteroidal anti-inflammatory agent after argon laser trabeculoplasty. A trial with flurbiprofen and indomethacin. *Ophthalmology* 1984; 91: 969-76.
- Weinreb RN, Robin AL, Baerveldt G, Drake MV, Blumenthal M, Wilensky J. Flurbiprofen pre-treatment in argon laser trabeculoplasty for primary open-angle glaucoma. *Arch Ophthalmol* 1984; 102: 1629-32.
- Goethals M, Missotten L. Efficacy and safety of indomethacin 0.1% versus flurbiprofen 0.03% eye drops in inflammation after argon laser trabeculoplasty. *Doc Ophthalmol* 1994; 85: 287-93.
- Hurvitz LM, Spaeth GL, Zakhour I, Mahmood E, Murray G. A comparison of the effect of flurbiprofen, dexamethasone and placebo on cyclo-oxygenase-induced inflammation. *Ophthalmic Surg* 1984; 15: 394-9.
- Garry R. Ketorolac tromethamine: a non-steroidal anti-inflammatory agent for ophthalmic use in the management of ocular itching associated with seasonal allergic conjunctivitis. *Allergan Technical Report* 1993.
- Tauber J, Abelson M, Ostrov C, Laibovitz R, Raizman M, Betts J, et al. A multicenter comparison of diclofenac sodium 0.1% to ketorolac tromethamine 0.5% in patients with acute seasonal allergic conjunctivitis. *Invest Ophthalmol Vis Sci* 1994; 35 (suppl): 1291.
- Van Huse H. Topical treatment of anterior ocular diseases with diclofenac Na eye drops. *Klin Monatsbl Augenheilkd* 1986; 188: 615-9.
- Murphy JE. Drug profile: Tanderil eye ointment. *J Int Med Res* 1973; 1: 136-40.
- Rushford WAI, Holman RM, Pulvertaft TB. Tanderil/Chloramphenicol eye ointment in the treatment of the 'Red eye' seen in general practice. *J Int Med Res* 1978; 6: 141-3.
- Emery DG, Smith VH, Wilkinson P. A trial of oxyphenbutazone eye ointment in general practice. *Practitioner* 1973; Dec: 3-7.
- Gimbel HV. The effect of treatment with topical nonsteroidal anti-inflammatory drugs with and without intra-operative epinephrine on the maintenance of pupillary mydriasis during cataract surgery. *Ophthalmology* 1989; 96: 585-8.
- Bonomi L, Totolo G, Marchini G, Rigotti P, Ghini M. Prevention of trauma induced miosis during cataract extraction by diclofenac sodium. *New Trends in Ophthalmology* 1987; 2: 512-8.
- Goa KL, Chriss P. Ocular diclofenac: a review of its pharmacology and clinical use in cataract surgery and potential in other inflammatory ocular conditions. *Drugs Aging* 1992; 2: 480.
- Psilas K, Kalogeropoulos C, Loucatzicos E, Asproudis I, Petroustos G. The effect of indomethacin, diclofenac and flurbiprofen on the maintenance of mydriasis during extracapsular cataract extraction. *Doc Ophthalmol* 1992; 81: 293-300.
- Norton AL, Brown WJ, Carson M. Pathogenesis of aphakic macular edema. *Am J Ophthalmol* 1975; 80: 86-101.
- Obstbaum SA, Galin MA. Cystoid macular edema and ocular inflammation. The corneo-retinal inflammatory syndrome. *Trans Ophthalmol Soc UK* 1979; 99: 187-91.
- Kraff MC, Sanders DR, Jampol LM, Lieberman HL. Factors affecting pseudophakic cystoid macular edema: five randomised trials. *J Am Intraocular Implant Soc* 1985; 11: 380-5.
- Abelson MB, Parver L. How to handle CME in pseudophakia. *Rev Ophthalmol* 1994; Oct: 138-9.
- Miyake K, Sakamura S, Muira H. Long-term follow-up study on the prevention of aphakic cystoid macular edema by topical indomethacin. *Br J Ophthalmol* 1980; 64: 324-8.
- Yannuzzi LA, Landau AN, Turtz AL. Incidence of aphakic cystoid macular edema with the use of topical indomethacin. *Ophthalmology* 1981; 88: 947-54.
- Kraff MC, Sanders DR, Jampol LM, Peyman GA, Lieberman HL. Prophylaxis of pseudophakic cystoid macular edema with topical indomethacin. *Ophthalmology* 1982; 89: 885-90.
- Quentin CD, Behrens-Baumann W, Gaus W. Prophylactic treatment of cystoid macula oedema with diclofenac eye drops in intracapsular cataract extraction with the Choyce Mark IX anterior chamber lens. *Fortschr Ophthalmol* 1989; 86: 546-9.
- Flach AJ, Stegman RC, Graham J, Kruger L, Kruger LP. Prophylaxis of aphakic cystoid macular edema without corticosteroids. *Ophthalmology* 1990; 97: 1253-8.
- Solomon LD. Efficacy of topical flurbiprofen and indomethacin in preventing pseudophakic cystoid macular edema. *J Cataract Refract Surg* 1995; 21: 73-81.
- Ginsburg AP, Cheetham JK, DeGryse RE, Abelson M. Effects of flurbiprofen and indomethacin on acute cystoid macular edema after cataract

- surgery: functional vision and contrast sensitivity. *J Cataract Refract Surg* 1995; 21: 82–92.
- 60 Flach AJ, Dolan BJ, Irvine AR. Effectiveness of ketorolac tromethamine 0.5% ophthalmic solution for chronic aphakic and pseudophakic cystoid macular oedema. *Am J Ophthalmol* 1987; 103: 479–86.
 - 61 Peterson M, Yoshizumi MO, Hepler R, Mondino B, Krieger A. Topical indomethacin in the treatment of chronic cystoid macular edema. *Graefes Arch Clin Exp Ophthalmol* 1992; 230: 401–5.
 - 62 Burnett J, Tessler H, Isenberg S, Tso MO. Double-masked trial of fenopropfen sodium: treatment of chronic aphakic cystoid macular edema. *Ophthalmic Surg* 1983; 14: 150–2.
 - 63 Drolsum L, Haaskjold E. Causes of decreased visual acuity after cataract extraction. *J Cataract Refract Surg* 1995; 21: 59–63.
 - 64 Ishikawa T, del Cerro M, Liang FQ, Kim JC, Aquavella JV. Hypersensitivity following excimer laser ablation through the corneal epithelium. *Refract Corneal Surg* 1992; 8: 466–74.
 - 65 Szerenyi K, Sorken K, Garbus JJ, Lee M, Mc Donnell PJ. Decrease in normal human corneal sensitivity with topical diclofenac sodium. *Am J Ophthalmol* 1994; 118: 312–5.
 - 66 Kaidman GW. Diclofenac and its effect on corneal sensation. *Arch Ophthalmol* 1995; 113: 262.
 - 67 Harr D. PRK: no pain, major gain. *Ocular Surg News* 1992; 21: 73–4.
 - 68 Sher NA, Frantz JM, Talley A, Parker P, Lane SS, Ostrov C, et al. Topical diclofenac in the treatment of ocular pain after excimer photorefractive keratectomy. *Refract Corneal Surg* 1993; 9: 425–36.
 - 69 Ekdahl J. Promising use of non-steroidals with PRK. *Eyecare Technology* 1994; 4: 58–105.
 - 70 Stein R, Stein HA, Cheskes A, Symons S. Photorefractive keratectomy and post-operative pain. *Am J Ophthalmol* 1994; 117: 403–5.
 - 71 Donnenfeld ED, Selkin BA, Perry HD, Moadel K, Selkin GT, Cohen AJ, et al. Controlled evaluation a bandage contact lens and a topical non-steroidal anti-inflammatory drug in treating traumatic corneal abrasions. *Ophthalmology* 1995; 102: 979–84.
 - 72 Epstein RL, Laurence EP. Relative effectiveness of topical ketorolac and topical diclofenac on discomfort after radial keratotomy. *J Cataract Refract Surg* 1995; 21: 156–9.
 - 73 Hoh H. Local anaesthetic effect and subjective tolerance of 0.5% levobunolol in normal eyes. *Klin Monatsbl Augenheilkd* 1990; 1: 20–6.
 - 74 Kulkarni PS, Srinivasan BD. Cyclooxygenase and lipoxygenase pathways in anterior uvea and conjunctiva [Review]. *Progr Clin Biol Res* 1989; 312: 39–52.
 - 75 Phillips AF, Szerenyi K, Campos M, Krueger RR, McDonnell PJ. Arachidonic acid metabolites after excimer laser corneal surgery. *Arch Ophthalmol* 1993; 111: 1273–8.
 - 76 Szerenyi KD, Campos M, McDonnell PJ. Prostaglandin E2 production after lamellar keratectomy and photorefractive keratectomy. *J Refract Corneal Surg* 1994; 10: 413–6.
 - 77 Hanson DL, DeLeo VA. Long wave ultraviolet radiation stimulates arachidonic acid release and cyclooxygenase activity in mammalian cell culture. *Photochem Photobiol* 1989; 49: 423–30.
 - 78 Levine JD, Gooding J, Donatoni P, Borden L, Goetzl EJ. The role of the polymorphonuclear leukocyte in hyperalgesia. *J Neurosci* 1985; 5: 3025–9.
 - 79 Szerenyi K, Wang XW, Lee M, Mc Donnell PJ. Topical diclofenac treatment prior to excimer laser photorefractive keratectomy in rabbits. *Refract Corneal Surg* 1993; 9: 437–42.
 - 80 Perianin A, Gougerot-Pocidalo MA, Giroud JP, Hakim J. Diclofenac sodium, a negative chemokinetic factor for neutrophil locomotion. *Biochem Pharmacol* 1985; 34: 3433–8.
 - 81 Srinivasan BD, Kulkarni PS. Polymorphonuclear leukocyte response. Inhibition following corneal epithelial denudation by steroidal and non-steroidal anti-inflammatory agents. *Arch Ophthalmol* 1981; 99: 1085–9.
 - 82 Leibowitz HM, Ryan WJ, Kupferman A, De Santis L. Effect of concurrent topical corticosteroid and NSAID therapy of experimental keratitis. *Invest Ophthalmol Vis Sci* 1986; 27: 1226–9.
 - 83 Arshinoff S, D'Addario D, Sadler C, Bilotta R, Johnson TM. Use of topical nonsteroidal anti-inflammatory drugs in excimer laser photorefractive keratectomy. *J Cataract Refract Surg* 1994; 20 (suppl): 216–22.
 - 84 Napoli SA, Helm C, Insler MS, Ensley HE, Pretus HA, Feigen LP. External ocular inflammatory effects of lipoxygenase enzyme products. *Ann Ophthalmol* 1990; 22: 30–4.
 - 85 Palmer RM, Stepney RJ, Higgs GA, Eakins KE. Chemokinetic activity of arachidonic acid lipogenase products on leukocytes of different species. *Prostaglandins* 1980; 20: 411–7.
 - 86 Stjernschantz J. Autocoids and neuropeptides. In: Sears ML, ed. *Pharmacology of the eye*. New York: Springer-Verlag, 1984: 311.
 - 87 Johnson D, Aston J. Issues and answers in excimer laser PRK. *Rev Ophthalmol* 1994; 7: 84–93.
 - 88 Ferrari M. Use of topical non-steroidal anti-inflammatory drugs after photorefractive keratectomy. *J Refract Corneal Surg* 1994; 10 (2 suppl): S287–9.
 - 89 Nassaralla BA, Szerenyi K, Wang XW, Al Reaves T, McDonnell PJ. Effect of diclofenac on corneal haze after photorefractive keratectomy in rabbits. *Ophthalmology* 1995; 102: 469–74.
 - 90 Frew A. Selected side effects: non-steroidal anti-inflammatory drugs and asthma. *Prescribers' J* 1994; 34: 2.
 - 91 Voltarol ophtha. ABPI data sheet compendium 1995–6: 416–7.
 - 92 Ocufen. ABPI data sheet compendium 1995–6: 74.
 - 93 Serle JB. Pharmacological advances in the treatment of glaucoma. *Drugs Aging* 1994; 5: 157–70.
 - 94 Bito LZ, Stjernschantz J, Resul B, Miranda OC, Basu S. The ocular effects of prostaglandins and the therapeutic potential of a new PGF2 alpha analogue, PhXA41 (latanoprost), for glaucoma management. *J Lipid Mediators* 1993; 6: 535–43.
 - 95 Gum CG, Kingsbury S, Whitley RD, Garcia A, Gelatt KN. Effect of topical prostaglandin PGA2. PGA2 isopropyl ester, and PGF2 alpha isopropyl ester on intraocular pressure in normotensive and glaucomatous canine eyes. *J Ocular Pharmacol* 1991; 7: 107–16.
 - 96 Ando Y, Matsunami C, Yamamoto T, Kitazawa Y. Safety and ocular hypotensive effect of new prostaglandin analogue, S-1033, in normal volunteers. *Acta Soc Ophthalmol Jap* 1994; 98: 1109–14.
 - 97 Sugrue MF, O'Neill-Davis L. The effect of cyclooxygenase inhibition on the ocular hypotensive action of topical carbonic anhydrase inhibitors in rabbits. *J Ocular Pharmacol* 1991; 7: 201–11.
 - 98 Trusopt Ophthalmic Solution 2% (dorzolamide hydrochloride). Merck Sharpe & Dohme Ltd, Data sheet 1995.