PERSPECTIVE

Ocular involvement in sarcoidosis

Aniki Rothova

Sarcoidosis is a chronic multisystemic granulomatous disorder thought to result from an exaggerated cellular immune response to a variety of self antigens or non-self antigens.¹ The aetiology of sarcoidosis is unknown, which might be in part related to the diverse manifestations of the disease and the absence of approved diagnostic criteria. Although specific inhalation antigens have been put forward as possible triggers, no study has yet proved a consistent relation with a causative agent.² Genetic factors might also be involved since familial incidence in certain populations as high as 19% has been noted and, further, specific HLA associations have been connected with disease susceptibility and outcome—for example, HLA-B8 and DR-3 with Löfgren's syndrome (acute systemic disease with fever, polyarthritides, erythema nodosum, and favourable prognosis).^{3–4} The infectious agents, specifically mycobacteria and more recently herpesvirus 8, have also been implicated.^{5–6} Sarcoidosis has only sporadically been reported in AIDS and has occasionally been described during the immune recovery with the highly active antiretroviral therapy.^{7–8} The evidence that disease might be of environmental or infectious origin is supported by occupational clustering (medical nurses, fire fighters) and by the transmission of the disease from transplanted organs.^{9–11} The genetic and environmental influences cannot be adequately evaluated in small samples of patients; a large study addressing the phenotyping differences and various environmental data is lacking however. So far, only interesting disease patterns in families and inconclusive allelic associations for different populations have been reported. Therefore, sarcoidosis might represent an abnormal immune response directed against one or more possible antigens in an individual with a hereditary or acquired abnormality of the immune system.

Epidemiology

Sarcoidosis occurs worldwide but is predominant in certain ethnic and racial groups (for example, US blacks, Scandinavian and Irish white people).¹² Black patients present typically with more severe and acute disease whereas the white patients usually have asymptomatic and chronic disease.^{13–14} The frequency of sarcoidosis is reported to be low in various parts of the world^{14–15}; it is not known whether this low frequency of sarcoidosis is genuine or whether it represents an underdiagnosis owing to the frequent occurrence of subclinical course, similarity with other diseases, or absence of firm diagnostic criteria.

Pathogenesis

Sarcoidosis is characterised by the formation of non-caseating granulomas in affected organs. Three types of agents are known to cause epithelioid cell granulomas—infectious organisms (bacteria and fungi), products of plants and animals (pollen, proteins), and metallic compounds (beryllium, zirconium). In sarcoid granulomas, the presence of mycobacterial DNA was identified by

molecular techniques and a causal role in the aetiology of sarcoidosis was proposed.^{16–17} Conflicting results have been obtained from the studies assessing the aetiological role of mycobacterial infection in the pathogenesis of sarcoidosis and the discrepancies were attributed to different sensitivities of the diagnostic procedures.^{18–19} The nested polymerase chain reaction (PCR), a procedure more sensitive than the standard PCR protocol, has failed to identify sequences specific for *Mycobacterium tuberculosis* complex in sarcoidosis, whereas positive results were found in biopsies from patients with active tuberculosis.^{20–21} Polymerase chain reactions of 123 bp fragment of IS6110 have shown the absence of *M. tuberculosis* DNA in lymph nodes and lung biopsies from patients with sarcoidosis and suggest that *M. tuberculosis* does not have a pathogenic role in sarcoidosis in most patients.²² By molecular techniques, atypical mycobacteria, *Corynebacterium acnes*, and *Propionibacterium* DNA have also been identified in sarcoid granulomas and a link with sarcoidosis has been proposed.^{23–24} Whether these bacteria induce some cases of sarcoidosis by an allergic mechanism has not yet been elucidated.

Presumably, in response to antigen exposure, exuberant cellular immune response occurs in target organs, and leads to the development of granulomas. Granulomas consist of epithelioid and multinucleated giant cells (aggregated macrophages), abundant CD4 positive T cells and, usually in the periphery of the granuloma, to some extent CD8 positive T cells and B cells may be encountered.^{25–26} T lymphocytes in affected organs are of T helper 1 phenotype and produce interferon γ and interleukin 2, which subsequently lead to the production of tumour necrosis factor and interleukin 6 by macrophages, and thereby cause a cascade of inflammatory changes, culminating in fibrosis. Fibrosis of the granulomas induces organ destruction and loss of function. The very high ratios of T helper to T suppressor cells have been reported for various organs affected by sarcoidosis, including the eye.²⁶ Epithelioid cell granulomas are not pathognomonic for sarcoidosis, and other causes of granulomatous inflammation (for example, Wegener's disease) as well as the presence of infectious agents (fungi or mycobacteria) and foreign bodies (beryllium, organic dusts) should be sought.¹ Probably secondary laboratory abnormalities include depressed cellular immunity to other antigens (resulting in cutaneous anergy), hyperglobulinaemia caused by overstimulated B lymphocytes, and hypercalcaemia.

So far, the search for the associations between HLA related genes and sarcoidosis has revealed that the most common allele associated with the susceptibility to sarcoidosis has been HLA-B8.²⁷ The present studies, using the DNA sequence analysis rather than serological typing, support the hypothesis that the major genetic factors controlling the development of sarcoidosis are located within the DRB1 locus in the HLA class II region.^{28–29} Several studies on genetic polymorphism have shown the potential to predict the clinical course of the disease.²⁸ Although the

HLA class II genotyping was associated with disease outcome in Scandinavian patients (HLA-DR 14 and 15 were associated with chronic sarcoidosis and HLA-DR 17 with non-chronic forms of sarcoidosis), in other populations other alleles were responsible for the susceptibility and course of the disease.^{30 31}

Clinical manifestations

The course of sarcoidosis ranges from asymptomatic to severe and even lethal disease. The clinical manifestations and severity of the disease vary widely and are strongly associated with racial and ethnic factors: acute and more severe disease is typical of black patients, whereas asymptomatic and chronic disease is more frequent in whites.¹² In all races, however, mortality remains similar.³² Although the overall prognosis is relatively benign, many patients develop some organ dysfunction. Factors associated with poor outcome include black race, late onset, and disease persisting for longer than 6 months, involvement of more than three organs, and stage III of pulmonary disease (parenchymal infiltrates without adenopathy).^{1 33} Sarcoidosis affects people of all ages with the peak age of onset in the third decade.¹³

The disease affects predominantly the lungs and thoracic lymph nodes, skin, and eyes and might have either self limited or chronic course. Although the manifestations of sarcoidosis may be widespread, as a result of the frequent manifestations in lungs and lymph nodes, most patients will have some respiratory problems.¹³ The majority of patients have constitutional symptoms such as fever, malaise, fatigue, and weight loss. Frequently the disease is asymptomatic and discovered by chest radiography (hilar adenopathy) performed for unrelated causes. The appearance of the chest radiograph is usually used to classify a pulmonary sarcoid: stage 1 represents bilateral hilar adenopathy without parenchymal involvement, stage 2 bilateral adenopathy with parenchymal involvement; and stage 3 is pulmonary infiltrates without hilar adenopathy, including cystic changes.^{34 35} Specific manifestations of sarcoidosis include acute Löfgren's syndrome, which is a combination of erythema nodosum, arthritis, and hilar lymphadenopathy (sometimes associated with anterior uveitis) and Heerfordt's syndrome (uveoparotid fever), consisting of fever, parotid swelling, uveitis, and sometimes facial palsy.³⁶ Most frequently affected extrapulmonary organs are lymph nodes, eyes, and skin. Skin lesions are common and include, in addition to sarcoid granulomas, erythema nodosum. Erythema nodosum represents a reaction pattern of the skin to diverse causes (the spontaneously healing lesions are tender red nodules located usually on the lower extremities); the lesions are not granulomatous and therefore biopsy does not contribute to the diagnosis of sarcoidosis. The development of granulomas is regularly noted in scar tissues. CNS manifestations occur in 5%–12% of patients.¹ Renal insufficiency may develop secondarily to granulomatous nephritis or hypercalcaemia. Cor pulmonale may develop secondarily to severe pulmonary disease. Direct sarcoid infiltration of the heart muscle is seen in about 5% of patients with sarcoidosis.

Sarcoidosis in children is frequently of extrapulmonary type with frequent ocular as well as cutaneous and rheumatological manifestations and has a guarded prognosis.^{37 38} The triad—rash, uveitis, and arthritis—is characteristic for children under 4 years of age; more frequent is the occurrence of sarcoidosis in older children (8–15 years of age), which is characterised by almost universal lung involvement and manifestations in eyes, skin, liver, and spleen. Long term follow up illustrated a high frequency of blindness and severe multisystem (for example, heart and renal) involvement in children with onset of

sarcoidosis at a very early age.³⁷ Childhood sarcoidosis might be misdiagnosed because of its rarity and resemblance to juvenile chronic arthritis. Autosomal, dominantly inherited Blau syndrome (familial granulomatous inflammatory arthritis, uveitis and rash, and camptodactylia) is characterised by multiorgan inflammation and may strongly resemble sarcoidosis; however, the typical pulmonary involvement is absent.^{39–41}

Diagnostic methods

In the absence of the known causative agent, the diagnosis of sarcoidosis remains mainly a diagnosis of exclusion. Since there are no definitive diagnostic serological or radiological tests, the presence of non-caseating granulomas on tissue biopsy together with compatible clinical features is usually considered as a proof of the diagnosis of sarcoidosis. However, even in cases with tissue biopsy consistent with the presence of sarcoidosis, other possible causes of granulomatous diseases should be evaluated. Although biopsy is not pathognomonic, tissue examination is essential to differentiate sarcoidosis from infections or malignancies. Tissue biopsy might be obtained from lung, skin, lymph node, or other affected organs. Undirected biopsies yielded less positive results compared with those obtained from overtly affected organs; this was also reported for conjunctival granuloma or lacrimal gland biopsy.^{42–44} The analysis of fluids and cells obtained by bronchoalveolar lavage, including CD4/CD8 ratio, is not specific enough for the definitive diagnosis of sarcoidosis.⁴⁵ Chest x ray usually shows hilar adenopathy in combination with interstitial lung disorder. Chest computed tomography has been reported to be more sensitive than radiography.⁴⁶ Measurements of serum angiotensin converting enzyme (ACE) level reflect the disease activity; ACE levels fluctuate with corticosteroid use, but are not specific enough for diagnostic purposes.^{47 48} High levels of serum ACE were also reported for other diseases (for example, tuberculosis, leprosy, diabetes mellitus). Moreover, as serum ACE levels reflect the systemic burden of inflammation, normal serum ACE levels do not exclude the diagnosis of sarcoidosis, especially not in those with isolated ocular disease. The patients with presumed sarcoidosis (intraocular inflammation compatible with the diagnosis of sarcoidosis in conjunction with raised serum ACE levels) were similar to those with histologically proved sarcoidosis in terms of ocular presentations and clinical course.⁴⁹ The combination of raised serum ACE levels with abnormal gallium scanning was a specific and sensitive tool for diagnosing patients suspected of having ocular sarcoidosis who had normal chest radiographs.⁵⁰ Aqueous ACE levels of patients with sarcoid associated uveitis were increased, though their diagnostic value was not yet determined.⁵¹ Increased levels of ACE in cerebrospinal fluid were reported, but their diagnostic role is not clear.⁵² Gallium scan uptake, serum lysozyme levels, hypergammaglobulinaemia, and decreased delayed type hypersensitivity are not specific for sarcoidosis. Kveim–Siltzbach test, which uses human tissue obtained from patients with sarcoidosis to induce granuloma in a tested patient, has been abandoned. Although lung involvement occurs at some time in the course of the disease in nearly all cases with sarcoidosis, the sensitivities and specificities of various diagnostic procedures for patients with ocular sarcoidosis are not known. In patients with uveitis, sensitivity of raised serum ACE was 73% and specificity 83%.⁵⁰ In patients with active systemic sarcoidosis, gallium-67 scanning had a sensitivity of 95%, but a low specificity (68%), while the sensitivity of chest radiography was 80%, and of serum ACE 77%.⁵³

Treatment

The optimal therapy for sarcoidosis is not well defined; therapeutic decisions are dictated by the localisation of the disease and severity of organ involvement. The mainstay of treatment is corticosteroid therapy, which exhibits especially short term beneficial effects.³⁴ The controversy remains, however, concerning the efficacy of corticosteroids to alter the natural course of the disease.^{35–36–54} For refractory disease, methotrexate was reported to be the most promising drug and was recently reported to be effective in ocular involvement.^{54–56} The efficacy of antimalarial agents and cyclosporin has not been conclusively proved.^{57–59}

Ocular involvement

Ocular disease may be the initial manifestation in sarcoidosis and may progress to severe visual impairment or even blindness. No specific extraocular manifestations of sarcoidosis were associated with the development of ocular involvement or uveitis.^{60–61} Two peaks of incidence were reported for ocular sarcoidosis, the first at ages 20–30 years and the second at ages 50–60 years.⁶¹ Ocular involvement manifests in 25%–60% of patients with systemic sarcoidosis.⁶² This percentage depends on the population studied and the extent of ocular examinations. The most common ocular manifestations are uveitis (30%–70%) and conjunctival nodules (40%).^{62–66}

Uveitis is a frequent and early feature of sarcoidosis. More than 80% of uveitis cases manifested before or within 1 year after the onset of systemic disease. Uveitis preceded the non-ocular signs of sarcoidosis in about 30%.⁶¹ In large surveys of uveitis patients, the frequency of sarcoidosis was about 5%.⁶² Traditionally, the most common type of sarcoid associated uveitis was attributed to anterior uveitis. It is noteworthy that the high frequency of sarcoid associated anterior uveitis (70%–75%) was noted mainly in the studies where the majority of patients were black.^{62–63–66–68} In contrast, posterior type was the most common localisation of sarcoid uveitis in white patients (65%–83%), specifically in elderly female patients.^{49–61–62–65–69} The visual impact of sarcoid associated uveitis was severe, about 10% of patients developed blindness in at least one eye.⁷⁰ The main cause of visual loss was cystoid macular oedema.^{49–71} The poor visual prognosis was associated with the advanced age of the patients, black race, female sex, chronic systemic disease, and also with posterior segment involvement, peripheral punched out lesions, and the presence of cystoid macular oedema and glaucoma.^{69–71–73}

CLINICAL MANIFESTATIONS

Neurological manifestations of sarcoidosis are varied and were identified in 12% (71/554) of patients with sarcoido-

sis. Posterior segment eye involvement was frequently accompanied by CNS involvement.⁶² Neurological manifestations may include optic nerve disease, cranial nerve palsies, encephalopathy, and disorders of the hypothalamus and pituitary gland.^{74–75} Chiasmal syndromes and motility disorders may also occur. Facial nerve palsy was the most common but usually self limited manifestation of CNS involvement. Optic nerve involvement may be caused either by direct sarcoid tissue infiltration or compression by cerebral mass, and may lead to optic nerve atrophy.^{76–78} Usually, the involvement of the optic nerve is an indication for systemic treatment. The treatment of chronic neurological symptoms with methotrexate or cyclophosphamide was associated with a better therapeutic response than the treatment with corticosteroids alone.

Lids and orbit

Lacrimal gland or conjunctival involvement is common and usually asymptomatic, although extensive granulomas leading to diplopia or severe keratoconjunctivitis sicca may develop.^{79–80} Eyelid or conjunctival granulomas were repeatedly reported and usually react well to topical treatment.⁸¹ In resistant cases, an intralesional corticosteroid administration may help. Other manifestations in orbit and lids are not frequent and are usually unilateral.⁸² Orbital symptoms mimic other inflammatory syndromes manifesting in the orbit. Sarcoid induced inflammatory myositis may resemble Graves' ophthalmopathy. The association of sarcoidosis with autoimmune thyroid disease is well known and the coincidence of sarcoidosis with Graves' disease has also been noted.⁸³

Corneal involvement by granulomas is extremely rare.⁸⁴ Corneal band degeneration may develop as a consequence of long standing anterior uveitis.

Uveitis

Acute signs of uveitis as pain, photophobia, lacrimation, or redness might be absent, so that the patients with sarcoid associated "silent uveitis" may develop permanent ocular damage before the intraocular process is diagnosed and treatment initiated. In a series of 121 patients with biopsy proved sarcoidosis, 29 patients were diagnosed with uveitis. Of these 29, 10 were diagnosed by an ophthalmologist in the absence of ocular complaints at the time of initial ophthalmological screening.⁶¹

Classic sarcoid associated anterior uveitis may either present as acute iridocyclitis, which is mostly seen in Löfgren's syndrome, or as a chronic granulomatous uveitis with keratic precipitates, which may vary from cellular to large "mutton fat" type. In the chronic type of the disease, the granulomatous nodules of the iris (Fig 1A) and in the

Figure 1 Anterior segment granulomas in sarcoidosis. (A) Granulomatous iridocyclitis in sarcoidosis with Koeppe nodules visible on the pupillary border. (B) Multiple, extremely large granulomas in anterior chamber in a black patient with biopsy proved sarcoidosis. Keratic precipitates located on the lower part of corneal endothelium are also visible.

Figure 2 Retinal vasculitis in sarcoidosis. (A) Retinal vasculitis in a 22 year old white male patient with sarcoidosis visible as focal perivascular inflammatory infiltrates. (B) Fluorescein angiogram of this patient exhibits focal perivenous leakage in the affected areas. (C) Retinal vasculitis in a 44 year old white male patient with sarcoidosis. Late phase fluorescein angiogram. Note the multiple segmental perivascular leakage areas. (D) Sheathing of the peripheral retinal vessels surrounded by haemorrhages in a 24 year old black female patient with sarcoidosis. The retinal abnormalities normalised after periocular corticosteroid administration.

anterior chamber angle are sometimes seen. These are usually whitish and range in size from small to large masses (Fig 1B). Granulomas located on trabecular meshwork (sometimes associated with elevated eye pressure) were noted in 49% of Japanese patients.⁸⁵ Chronic anterior uveitis may lead to secondary cataract, glaucoma, and cystoid macular oedema; corneal band keratopathy may also develop.

Intermediate uveitis with vitritis and genuine snow banking may be occasionally encountered. More frequent is the presence of vitritis with peripheral vasculitis and snowball infiltrates. This type of intermediate uveitis may precede more severe posterior segment changes. Snowball opacities in the vitreous cavity are characteristic but not exclusive for sarcoidosis and the differentiation from idiopathic pars planitis and other types of intermediate uveitis might be difficult.

Characteristic fundusoscopic findings in posterior segment involvement include periphlebitis, sometimes subclinical and only visible on fluorescein angiography (Fig 2), associated with typical segmental cuffing or more extensive sheathing and perivenous exudates, which are usually indicated as “candle wax drippings” (Fig 2).^{86–88} The term “taches de bougie” was originally used by Franceschetti in 1949, to indicate multiple small round chorioretinal lesions, which were located in the peripheral retina and were not associated with retinal vessels⁸⁶; this original term was probably first translated as “candle wax drippings” and (with the original description already forgotten) subsequently used to indicate the different characteristic features of sarcoidosis—namely, perivenous exudates.

Choroidal lesions may be multifocal and located anywhere in the fundus. Round punched out lesions (the

original “taches de bougies”) are frequently seen in the peripheral retina; recently the clinical entity of peripheral multifocal chorioretinitis was described in elderly white females, characterised by the presence of intraocular inflammation and multiple (>10) punched out lesions in the peripheral retina (Fig 3).⁷³ This entity was predominantly seen in elderly white female patients and was associated with the frequent occurrence of cystoid macular oedema (72%) and visual loss (42%). The association of acute posterior multifocal placoid pigment epitheliopathy (AMPPE) and sarcoidosis has also been noted.⁸⁹ Optic disc leakage may be frequently observed on fluorescein angiography and might be secondary to uveitis. Differentiation from genuine optic nerve disease is required because of the therapeutic consequences. Supposedly choroidal granuloma may be observed as an elevated white mass (Fig 4).^{90–92} The association of posterior segment and neurological involvement in sarcoidosis has been reported to be as high as 27%.⁶² The presentation as haemorrhagic retinopathy resembling Eales’ disease has been described as well as branch and central retinal vein occlusions, development of capillary non-perfusion, and subsequent neovascularisations, which may react well to anti-inflammatory treatment (Figs 2D and 5).^{93–98} The development of retinal pigment detachment has also been noted. Optociliary shunts, dilated collateral veins on the optic nerve head, connecting the central retinal vein to the peripapillary choroidal venous plexus, have been reported, and may be misdiagnosed as neovascularisation of the optic disc (Fig 6).⁷⁸ Arterial macroaneurysms, occurring in elderly female patients with peripheral multifocal chorioretinitis have been described, and were associated with severe cardiovascular disease (Fig 7).⁹⁹ The most frequent complications of

Figure 3 Multiple chorioretinal punched out lesions in the peripheral fundus in a 64 year old white female patient with sarcoidosis (peripheral multifocal chorioretinitis). Note the confluence of the lesions in the far periphery.

posterior segment involvement in sarcoidosis included cystoid macular oedema (76%), cataract (49%), glaucoma (36%), retinal ischaemia (16%), and neovascularisations (11%).⁷¹ Fluorescein angiography is a standard technique for monitoring the posterior segment activity; recently indocyanine green (ICG) angiographic findings in patients with posterior segment sarcoidosis demonstrated the widespread extent of choroidal involvement. ICG angiography may therefore represent an additional tool for monitoring the disease spread and activity.¹⁰⁰

Differential diagnosis is wide and sarcoidosis should be looked for in almost all forms of intraocular inflammation. Differentiation from other granulomatous diseases such as tuberculosis should be attempted. Patients with the clinical features of intermediate uveitis (pars planitis), multifocal chorioretinitis, Behçet's disease, syphilis, toxoplasmosis,

Figure 5 Peripheral area of capillary non-perfusion with impending development of the neovascular membrane in ocular sarcoidosis. Late phase fluorescein angiogram.

and intraocular lymphoma should also be assessed for sarcoidosis. Birdshot chorioretinopathy may strongly resemble sarcoidosis; the true association has also been noted.^{101–103} Retinal abnormalities from sickle cell disease and other haemoglobinopathies in young black people may also be confused with ocular sarcoidosis.

TREATMENT OF OCULAR SARCOIDOSIS

Corticosteroids are the mainstay of therapy for ocular sarcoidosis and are given preferably in topical or periocular administration. Cycloplegics are frequently needed to prevent synechiae. The indication for systemic treatment in ocular disease includes optic neuritis and severe or topical treatment resistant posterior uveitis. The initial dosage depends on the disease severity, but often high dosage is required to induce the remission or quiescent stage of

Figure 4 Posterior segment granulomas in sarcoidosis. (A) Peripheral choroidal nodule surrounded by several small chorioretinal lesions in a 26 year old white male patient with sarcoidosis. (B) Whitish chorioretinal nodules nasal to optic nerve head in a 55 year old white female patient with sarcoidosis. (C) Scarification of a large choroidal granuloma in 30 year old white male patient with sarcoidosis located in inferior part of the fundus and associated with secondary changes of retinal pigment epithelium.

Figure 6 Optic disc and peripapillary atrophy with abnormal vessels (optociliary shunts) of the optic nerve head in a patient with juvenile onset sarcoidosis. Focal chorioretinal lesions in different stages of inflammatory activity were visible throughout the fundus and in the peripapillary area.

Figure 7 Arterial macroaneurysms in a patient with sarcoidosis. Late phase fluorescein angiogram.

intraocular inflammation; thereafter slow tapering, and prolonged low dose treatment is usually administered to maintain the disease quiescence. For corticosteroid resistant patients, administration of low dose methotrexate therapy may be effective.^{55 56} There remains some debate over how to treat the patients with cystoid macular oedema; treatment will usually be initiated for visual acuity below 20/40, but possibly more early treatment is required to prevent future visual loss. The surgical intervention in ocular sarcoidosis should preferably be performed at a quiescent stage; with aggressive inflammation control before cataract surgery, IOL implantations were well tolerated and good visual results were achieved.¹⁰⁴ The major cause of the decreased visual acuity following cataract extractions was the pre-existing glaucomatous damage and posterior segment involvement, especially the presence of cystoid macular oedema. Neovascularisation may regress with systemic anti-inflammatory treatment, though cases with retinal ischaemia, persistent neovascularisation, or vitreous haemorrhage benefit from laser treatment.¹⁰⁵ Vitrectomy may be essential for the differentiation of sarcoidosis from intraocular malignancy, and for cases with persistent vitreous haemorrhage or resistant vitreous opacities. The effect of the removal of the vitreous on intraocular inflammation and on cystoid macular oedema has not yet been proved.

Conclusions

Ocular sarcoidosis may present with a wide variety of ocular symptoms in all parts of the eye and may be associated

with chronic and progressive intraocular inflammation leading to visual deterioration. The diagnosis may be difficult owing to the absence of diagnostic criteria and the variety of presentations. Although isolated anterior uveitis is more frequently seen in black patients, the (more severe) posterior segment involvement is encountered in white patients. Visual loss is associated with posterior segment involvement and is caused by cystoid macular oedema in the majority of cases. Treatment, which has not changed during the past decades, consists mostly of symptomatic administration of corticosteroids. An evaluation of the efficacy of other therapeutic alternatives for ocular sarcoidosis is needed.

A ROTHOVA

Department of Ophthalmology, FC Donders Institute, University Medical Centre, Utrecht, PO Box 85 500, 3508 GA Utrecht, Netherlands

- 1 Newman LS, Rose CS, Maier LA. Sarcoidosis. *N Engl J Med* 1997;336:1223–34.
- 2 Bresnitz EA, Stolley PD, Israel HL, et al. Possible risk factors for sarcoidosis: a case-control study. *Ann NY Acad Sci* 1986;465:632–42.
- 3 Gardner J, Kennedy HG, Hamblin A, et al. HLA associations in sarcoidosis: a study of two ethnic groups. *Thorax* 1984;39:19–22.
- 4 Brennan NJ, Crean P, Long JP. High prevalence of familial sarcoidosis in an Irish population. *Thorax* 1984;34:14–18.
- 5 El-Zaatari FA, Graham DY, Samuelsson K. Detection of Mycobacterium avium complex in cerebrospinal fluid of a sarcoid patient by specific polymerase chain reaction assays. *Scand J Infect Dis* 1997;29:202–4.
- 6 Di Alberti L, Piattelli A, Favia G, et al. Human herpesvirus 8 variants in sarcoid tissues. *Lancet* 1997;350:1655–61.
- 7 Amin DN, Sperber K, Brown LK, et al. Positive Kveim test in patients with coexisting sarcoidosis and human immunodeficiency virus infection. *Chest* 1992;101:1454–6.
- 8 Mirmirani P, Maurer TA, Herndier B, et al. Sarcoidosis in a patient with AIDS: a manifestation of immune restoration syndrome. *J Am Acad Dermatol* 1999;41:285–6.
- 9 Edmonstone WM. Sarcoidosis in nurses: is there an association? *Thorax* 1988;43:342–3.
- 10 Burke WMJ, Keogh A, Maloney PJ, et al. Transmission of sarcoidosis via cardiac transplantation. *Lancet* 1990;336:1579.
- 11 Kern DG, Neill MA, Wrenn DS. Investigation of a unique time-space cluster of sarcoidosis in firefighters. *Am Rev Respir Dis* 1993;148:974–80.
- 12 Rybicki BA, Major M, Popovich J. Racial differences in sarcoidosis incidence: a 5-year study in a health maintenance organization. *Am J Epidemiol* 1995;145:234–41.
- 13 Maycock RL, Bertrand P, Morrison CE, et al. Manifestations of sarcoidosis. Analysis of 145 patients, with a review of nine series selected from the literature. *Am J Med* 1963;35:67–89.
- 14 Siltzbach LE, Geraint James D, Neville E, et al. Course and prognosis of sarcoidosis around the world. *Am J Med* 1974;57:847–52.
- 15 Gupta SK, Mitra K, Chatterjee S. Sarcoidosis in India. *Br J Dis Chest* 1985;79:275–83.
- 16 Saboor SA, Johnson NM, McFadden J. Detection of mycobacterial DNA in sarcoidosis and tuberculosis with polymerase chain reaction. *Lancet* 1992;339:1012–5.
- 17 Fiddler HM, Rook GA, Johnson NM, et al. Mycobacterium tuberculosis DNA in tissue affected by sarcoidosis. *BMJ* 1993;306:546–9.
- 18 Vokurka M, Lecossier D, Du Bois RM, et al. Absence of DNA from mycobacteria of the M tuberculosis complex in sarcoidosis. *Am J Respir Crit Care Med* 1997;156:1000–3.
- 19 Kon OM, Du Bois RM. Mycobacteria and sarcoidosis. *Thorax* 1997;52:S47–51.
- 20 Hance AJ. The role of mycobacteria in the pathogenesis of sarcoidosis. *Sem Respir Infect* 1998;13:197–205.
- 21 Vago L, Barberis M, Gori A, et al. Nested polymerase chain reaction for Mycobacterium tuberculosis IS6110 sequence on formalin-fixed paraffin-embedded tissues with granulomatous diseases for rapid diagnosis of tuberculosis. *Am J Clin Pathol* 1998;109:411–5.
- 22 Wilsher ML, Menzies RE, Croxson MC. Mycobacterium tuberculosis DNA in tissues affected by sarcoidosis. *Thorax* 1998;53:871–4.
- 23 Ishige I, Usui Y, Takemura T, et al. Quantitative PCR of mycobacterial and propionibacterial DNA in lymph nodes of Japanese patients with sarcoidosis. *Lancet* 1999;354:120–3.
- 24 Popper HH. Epithelioid cell granulomatosis of the lung: new insights and concept. *Sarcoidosis Vasculitis and Diffuse Lung Diseases* 1999;16:32–46.
- 25 Moller DR. Systemic sarcoidosis. In: Jeffers JD, Shein LA, eds. *Fishman's pulmonary diseases and disorders*. New York: McGraw-Hill, 1998:1055–68.
- 26 Chan CH, Wetzig RP, Palestine AG, et al. Immunohistopathology of ocular sarcoidosis. *Arch Ophthalmol* 1987;105:1398–402.
- 27 Smith MJ, Turton CW, Mitchell DN, et al. Association of HLA B8 with spontaneous resolution in sarcoidosis. *Thorax* 1981;36:296–8.
- 28 Rybicki BA, Maliarik MJ, Major M, et al. Genetics of sarcoidosis. *Clin Chest Med* 1997;18:707–17.
- 29 Ishihara M, Ohno S. Genetic influences on sarcoidosis. *Eye* 1997;11:155–61.
- 30 Berlin M, Fogdell-Hahn A, Olerup O, et al. HLA-DR predicts the prognosis in Scandinavian patients with pulmonary sarcoidosis. *Am J Respir Crit Care Med* 1997;156:1601–5.
- 31 Maliarik MJ, Chen KM, Major ML, et al. Analysis of HLA-DPB1 polymorphisms in African-Americans with sarcoidosis. *Am J Respir Crit Care Med* 1998;158:111–14.
- 32 Keller AZ. Hospital, age, racial, occupational, geographical, clinical and survivorship characteristics in the epidemiology of sarcoidosis. *Am J Epidemiol* 1971;94:222–30.

- 33 Takada K, Ina Y, Noda M. The clinical course and prognosis of patients with severe, moderate or mild sarcoidosis. *J Clin Epidemiol* 1993;46:359–66.
- 34 Hunninghake GW, Gilbert S, Pueringer R. Outcome of the treatment for sarcoidosis. *Am J Resp Crit Care Med* 1994;149:893–8.
- 35 Gottlieb JE, Israel HL, Steiner RM. Outcome in sarcoidosis. The relationship of relapse to corticosteroid therapy. *Chest* 1997;111:623–31.
- 36 Bruins Slot WJ. Ziekte van Besnier Boeck en febris uveoparotidea. *Ned T Geneesk* 1936;80:2859–63.
- 37 Fink CW, Cimaz R. Early onset sarcoidosis: not a benign disease. *J Rheumatol* 1997;24:174–7.
- 38 Hoover DL, Khan JA, Giangiacomo J. Pediatric ocular sarcoidosis. *Surv Ophthalmol* 1986;30:215–28.
- 39 Chadarevian de JP, Raphael SA, Murphy GF. Histologic, ultrastructural, and immunocytochemical features of the granulomas seen in a child with the syndrome of familial granulomatous arthritis, uveitis, and rash. *Arch Pathol Lab Med* 1993;117:1050–2.
- 40 Tromp G, Kuivaniemi H, Raphael S, et al. Genetic linkage of familial granulomatous inflammatory arthritis, skin rash, and uveitis to chromosome 16. *Am J Hum Genet* 1996;59:1097–107.
- 41 James GD. A comparison of Blau's syndrome and sarcoidosis. *Sarcoidosis* 1994;11:100–1.
- 42 Spaide RF, Ward DL. Conjunctival biopsy in the diagnosis of sarcoidosis. *Br J Ophthalmol* 1990;74:469–71.
- 43 Karma A. Diagnostic aspects of ocular sarcoidosis. *Sarcoidosis* 1994;11:58–60.
- 44 Weinreb RN. Diagnosing sarcoidosis by transconjunctival biopsy of the lacrimal gland. *Am J Ophthalmol* 1984;97:573–6.
- 45 Greening AP, Nunn P, Dobson N, et al. Pulmonary sarcoidosis: alterations in bronchoalveolar lymphocytes and T cell subsets. *Thorax* 1985;40:278–83.
- 46 Kosmorsky GS, Meisler DM, Rice TW, et al. Chest computed tomography and mediastinoscopy in the diagnosis of sarcoidosis-associated uveitis. *Am J Ophthalmol* 1998;126:132–4.
- 47 Roulston JE, O'Malley GI, Douglas JG. Effects of prednisolone on angiotensin converting enzyme activity. *Thorax* 1984;39:356–60.
- 48 Baarsma GS, La Hey E, Glasius E, et al. The predictive value of serum angiotensin converting enzyme and lysozyme levels in the diagnosis of ocular sarcoidosis. *Am J Ophthalmol* 1987;104:211–17.
- 49 Stavrou P, Linton S, Young DW, et al. Clinical diagnosis of ocular sarcoidosis. *Eye* 1997;11:365–70.
- 50 Power WJ, Rodriguez A, Pedroze-Seres M. The value of combined serum angiotensin-converting enzyme and gallium scan in diagnosing ocular sarcoidosis. *Ophthalmology* 1995;102:2007–11.
- 51 Weinreb RN, Sandman R, Ryder MI, et al. Angiotensin-converting enzyme activity in human aqueous humor. *Arch Ophthalmol* 1985;103:34–6.
- 52 Oksanen V. New cerebrospinal fluid, neurophysiological and neuroradiological examinations in the diagnosis and follow-up of neurosarcoidosis. *Sarcoidosis* 1987;4:105–10.
- 53 Köhn H, Klech H, Mostbeck A, et al. ⁶⁷Ga scanning for assessment of disease activity and therapy decisions in pulmonary sarcoidosis in comparison to chest radiography, serum ACE and blood T-lymphocytes. *Eur J Nucl Med* 1982;7:413–416.
- 54 Lower EE, Baughman RP. Prolonged use of methotrexate for sarcoidosis. *Arch Intern Med* 1995;155:846–51.
- 55 Dev S, McCallum RM, Jaffe GJ. Methotrexate treatment for sarcoid-associated panuveitis. *Ophthalmology* 1999;106:111–8.
- 56 Shetty AK, Zganjar BE, Ellis GS, et al. Low-dose methotrexate in the treatment of severe juvenile rheumatoid arthritis and sarcoid iritis. *J Pediatr Ophthalmol Strabismus* 1999;36:125–8.
- 57 Wallace DJ. The use of chloroquine and hydroxychloroquine for non-infectious conditions other than rheumatoid arthritis or lupus: a critical review. *Lupus* 1996;5:S59–64.
- 58 Pia G, Pascalis L, Aresu GA, et al. Evaluation of the efficacy and toxicity of the cyclosporine A-flucortolone-methotrexate combination in the treatment of sarcoidosis. *Sarcoidosis Vasculitis and Diffuse Lung Diseases* 1996;13:146–52.
- 59 Baltzan M, Mehta S, Kirkham TH, et al. Randomized trial of prolonged chloroquine therapy in advanced pulmonary sarcoidosis. *Am J Resp Crit Care Med* 1999;160:192–7.
- 60 Kaye GB, Salmon JF, Isaacs S. Risk factors for uveitis in sarcoidosis. *S Afr Med J* 1993;83:342–4.
- 61 Rothova A, Alberts C, Glasius E, et al. Risk factors for ocular sarcoidosis. *Doc Ophthalmol* 1989;72:287–96.
- 62 Hunter DG, Foster CS. Ocular manifestations of sarcoidosis. In: Albert DM, Jakobiec FA, eds. *Principles and practice of ophthalmology*. Philadelphia: WB Saunders, 1994:443–50.
- 63 Obenaus CD, Shaw HE, Sydnor CF, et al. Sarcoidosis and its ophthalmic manifestations. *Am J Ophthalmol* 1978;86:648–55.
- 64 James GD, Anderson R, Langley D, et al. Ocular sarcoidosis. *Br J Ophthalmol* 1964;48:461–70.
- 65 Karma A. Ophthalmic changes in sarcoidosis. *Acta Ophthalmol (Suppl)* 1979;141:1–94.
- 66 Jabs DA, Johns CJ. Ocular involvement in chronic sarcoidosis. *Am J Ophthalmol* 1986;102:297–301.
- 67 Rothova A, Buitenhuis HJ, Meenken C, et al. Uveitis and systemic disease. *Br J Ophthalmol* 1992;76:137–41.
- 68 Dresner MS, Brecher R, Henkind P. Ophthalmology consultation in the diagnosis and treatment of sarcoidosis. *Arch Intern Med* 1986;146:301–4.
- 69 Dana MR, Merayo-Llves J, Schaumberg DA, et al. Prognosticators for visual outcome in sarcoid uveitis. *Ophthalmology* 1996;103:1848–53.
- 70 Rothova A, Suttrop-van Schulten MSA, Treffers WF, et al. Causes and frequency of blindness in patients with intraocular inflammatory disease. *Br J Ophthalmol* 1996;80:332–6.
- 71 Rothova A. Posterior segment involvement in sarcoidosis. In: Ohno S, Aoki K, Usui M, Uchio E, eds. *Uveitis today*. Amsterdam: Elsevier, 1998:207–10.
- 72 Silver MR, Messner LV. Sarcoidosis and its ocular manifestations. *J Am Optom Assoc* 1994;65:321–7.
- 73 Lardenoye CWTA, Van der Lelij A, Loos W de, et al. Peripheral multifocal chorioretinitis: a distinct clinical entity? *Ophthalmology* 1997;104:1820–6.
- 74 Stern BJ, Krumholz A, Johns C. Sarcoidosis and its neurological manifestations. *Arch Neurol* 1985;42:909–16.
- 75 Lower EE, Broderick JP, Brott TG, et al. Diagnosis and management of neurological sarcoidosis. *Arch Intern Med* 1997;157:1864–8.
- 76 Beardsley ThL, Brown SVL, Sydnor CF. Eleven cases of sarcoidosis of the optic nerve. *Am J Ophthalmol* 1984;1:62–77.
- 77 Karma A, Mustonen EMA. Optic nerve involvement in sarcoidosis. *Neuro-ophthalmology* 1985;5:231–46.
- 78 Mansour AM. Sarcoid optic disc edema and opticociliary shunts. *J Clin Neuro-ophthalmol* 1986;6:47–52.
- 79 Diestelmeier MR, Sausker WF, Pierson DL. Sarcoidosis manifesting as eyelid swelling. *Arch Dermatol* 1982;118:356–7.
- 80 Sacher M, Lanzieri CF, Sobel LI. Computed tomography of bilateral lacrimal gland sarcoidosis. *J Comp Ass Tomograph* 1984;8:213–5.
- 81 Hall JG, Cohen KL. Sarcoidosis of the eyelid skin. *Am J Ophthalmol* 1995;119:100–1.
- 82 Collision JMT, Miller NR, Green WR. Involvement of orbital tissues by sarcoid. *Am J Ophthalmol* 1986;102:302–7.
- 83 Papadopoulos KI, Hornblad Y, Liljeblad H, et al. High frequency of endocrine autoimmunity in patients with sarcoidosis. *Eur J Endocrinol* 1996;134:331–6.
- 84 Lennarson P, Barney NP. Interstitial keratitis as presenting ophthalmic sign of sarcoidosis in a child. *J Pediatr Ophthalmol Strabismus* 1995;32:194–6.
- 85 Mizuno K, Takahashi J. Sarcoid cycloplegia. *Ophthalmology* 1996;93:511–7.
- 86 Francheschetti A, Babel J. La chorio-rétinite en “taches de bougie”, manifestation de la maladie de Besnier-Boeck. *Ophthalmologica* 1949;118:701–10.
- 87 Spalton DJ, Sanders MD. Fundus changes in histologically confirmed sarcoidosis. *Br J Ophthalmol* 1981;65:348–58.
- 88 Sanders MD, Shilling JS. Retinal, choroidal, and optic disc involvement in sarcoidosis. *Trans Ophthalmol Soc UK* 1976;96:140–4.
- 89 Dick DJ, Newman PK, Richardson J, et al. Acute posterior multifocal placoid pigment epitheliopathy and sarcoidosis. *Br J Ophthalmol* 1988;72:74–7.
- 90 Campo RV, Aaberg TM. Choroidal granuloma in sarcoidosis. *Am J Ophthalmol* 1984;97:419–27.
- 91 Marcus DF, Bovino JA, Burton TC. Sarcoid granuloma of the choroid. *Ophthalmology* 1982;89:1326–30.
- 92 Olk RJ, Lipmann MJ, Cundigg HC. Solitary choroidal mass as the presenting sign in systemic sarcoidosis. *Br J Ophthalmol* 1983;67:826–9.
- 93 Gragoudas ES, Regan CDJ. Peripapillary subretinal neovascularization in presumed sarcoidosis. *Arch Ophthalmol* 1981;99:1194–7.
- 94 DeRosa AJ, Margo CE, Orlick ME. Hemorrhagic retinopathy as the presenting manifestation of sarcoidosis. *Retina* 1995;15:422–7.
- 95 Ohara K, Okubo A, Sasaki H. Branch retinal vein occlusion in a child with ocular sarcoidosis. *Am J Ophthalmol* 1995;119:806–7.
- 96 Kimmel AS, McCarthy MJ, Blodi CF, et al. Branch retinal vein occlusion in sarcoidosis. *Am J Ophthalmol* 1989;107:561–2.
- 97 Duker JS, Brown GC, McNamara JA. Proliferative sarcoid retinopathy. *Ophthalmology* 1988;95:1680–6.
- 98 Dees C, Arnold JJ, Forrester JV, et al. Immunosuppressive treatment of choroidal neovascularisation associated with endogenous posterior uveitis. *Arch Ophthalmol* 1998;116:1456–61.
- 99 Rothova A, Lardenoye CWTA. Arterial macroaneurysms in peripheral multifocal chorioretinitis associated with sarcoidosis. *Ophthalmology* 1998;105:1393–7.
- 100 Wolfensberger TJ, Herborst CP. Indocyanine green angiographic features in ocular sarcoidosis. *Ophthalmology* 1999;106:285–9.
- 101 Priem HA, Oosterhuis JA. Birdshot chorioretinopathy. Clinical characteristics and evolution. *Br J Ophthalmol* 1988;72:646–59.
- 102 Vrabec TR, Augsburger JJ, Fischer DH. Taches de Bougie. *Ophthalmology* 1995;102:1712–21.
- 103 Brod RD. Presumed sarcoid choroidopathy mimicking birdshot retinopathy. *Am J Ophthalmol* 1990;109:357–8.
- 104 Akova YA, Foster CS. Cataract surgery in patients with sarcoidosis-associated uveitis. *Ophthalmology* 1994;101:473–9.
- 105 Hirose S, Ohno S. Argon laser treatment of retinal neovascularisation associated with sarcoidosis. *Jpn J Ophthalmol* 1984;28:356–61.