

Prevalence of retinopathy at diagnosis among type 2 diabetic patients attending a diabetic centre in south India

Mohan Rema, Raj Deepa, Viswanathan Mohan

Abstract

Aim—To assess the prevalence of retinopathy in newly diagnosed south Indian type 2 diabetic patients attending a diabetic centre.

Methods—448 consecutive newly diagnosed type 2 diabetic patients were recruited. Four field retinal colour photography was performed and graded using a modified form of the Early Treatment Diabetic Retinopathy Study grading system.

Results—Of the 438 patients with assessable photographs, 32 (7.3%, 95% confidence interval 5.0 to 10.2) had retinopathy. None of the variables tested showed a significant association with retinopathy on univariate or multivariate logistic regression analysis.

Conclusion—The overall prevalence of retinopathy at diagnosis among clinic based south Indian patients with type 2 diabetes appears to be lower than that reported among Europeans.

(*Br J Ophthalmol* 2000;**84**:1058–1060)

Diabetic retinopathy is one of the leading causes of blindness in the western world but its frequency varies in different ethnic groups.^{1–7} As the prevalence of diabetes is high among Asian Indians,^{8,9} studies on diabetic complications are of considerable clinical significance. We report on the prevalence of diabetic retinopathy in newly diagnosed south Indian type 2 diabetic patients.

Patients and methods

The study group comprised 448 consecutive newly diagnosed type 2 diabetic patients attending the MV Diabetes Specialities Centre at Chennai (formerly Madras) in southern India. This centre provides treatment to approximately 15% of the diabetic population of Chennai. About 95% of patients seen at the centre are drawn from southern India (Tamil Nadu, Andhrapradesh, Karnataka, and Kerala). An ongoing epidemiological study has shown that the clinical profile of the patients seen at the centre is fairly representative of the diabetic population of Chennai.¹⁰ Type 2 diabetes mellitus was diagnosed using the criteria of the WHO study group on diabetes.¹¹ Clinical examination included recording of height, weight, and calculation of body mass index (BMI). The blood pressure was recorded in the right arm while seated.

Best corrected Snellen's acuity was recorded and analysis performed using the Snellen's reciprocal.¹² Slit lamp examination was performed to document any abnormality in the anterior segment. Intraocular pressure was measured using a Schiottz tonometer before dilating the pupils. There was, however, no contraindication for dilatation in any patient.

Retinopathy was assessed after adequate mydriasis by colour photography. Four field colour photography of the retina was taken of both eyes using a 50VT Topcon retinal camera using 35 mm colour transparencies. The four fields taken were stereoscopic pictures of the macula, superior temporal, nasal, and inferior temporal quadrants.¹³ Four fields were chosen to reduce the costs and to improve compliance. The nasal field centred on the nasal margin of the disc to assess the disc, superior and inferior nasal fields in one photograph. Photographs were coded and assessed in a masked manner for the presence and severity of retinopathy and quality of photographs. The photographs were graded twice by the same observer 4 weeks apart and the mean of the two readings was taken. A modified version of the Early Treatment Diabetic Retinopathy Study (ETDRS) grading system^{14–16} was used to grade the photographs. A simplified description of the retinopathy levels allocated is shown in Table 1.

The biochemical parameters (plasma glucose, serum cholesterol, and serum triglycerides) were measured using commercial kits (Boehringer Mannheim, Germany) on Ciba Corning Express Plus Autoanalyser (Medfield, USA). Glycosylated haemoglobin (HbA_{1c}) was measured using a machine Variant (Bio Rad, Hercules, CA, USA). Proteinuria was estimated by the sulphosalicylic acid method.¹⁷

Table 1 Levels of retinopathy

Level	Definition
10	No retinopathy
20	≥1 Microaneurysm
30	Microaneurysms + minimal haemorrhages and cotton wool spots
31–40	Microaneurysms, haemorrhages, cotton wool spots, and/or venous changes in one quadrant and/or intraretinal microvascular abnormalities and/or hard exudates
41–50	Microaneurysms, haemorrhages more than level 40 + venous changes in two or more quadrants and/or intraretinal microvascular abnormalities and/or cotton wool spots
>50	Severe preproliferative retinopathy with venous beading, intraretinal microvascular abnormalities and numerous cotton wool spots
60	Proliferative retinopathy
70	Proliferative retinopathy and high risk characteristics

Madras Diabetes Research Foundation, 35 Conran Smith Road, Gopalapuram, Chennai 600 086, India
M Rema
R Deepa
V Mohan

Correspondence to:
Dr M Rema
drema@md2.vsnl.net.in

Accepted 3 May 2000

Table 2 Prevalence of retinopathy by photographic assessment

Retinopathy level	No (n=438)	%
10	406	92.7
20	23	5.3
30	4	0.9
31-50	4	0.9
≥60	1	0.2

Statistical analysis was carried out using the SPSS package (version 4.0.1, Chicago). Student's *t* tests, χ^2 tests, or Fisher's exact tests were employed as appropriate. Univariate and multiple logistic regression analyses were carried out using retinopathy as the dependent variable and age, body mass index, systolic BP, diastolic BP, glycosylated haemoglobin, serum cholesterol, triglycerides, and smoking as independent variables.

Results

The study population consisted of 448 patients with a mean age of 48 (SD 14) years; 69% of the patients were men with a mean age of 46 (12) years and BMI 24.1 (3.5) kg/m² while the women were of mean age 47 (17) years with BMI 26.2 (5.0) kg/m².

Of the 448 patients photographed the photographs were of good quality in 398 (88.8%), in 40 patients (8.9%) they were satisfactory, and in three (0.7%) they were unreadable. In seven patients the photographs could not be taken for various reasons: camera failure (n = 2), mature cataract (n = 2), or inability to cooperate (n = 3). Of the 438 patients with assessable photographs, the photographs of both eyes could be graded in 430 patients (98.2%); 1.1% of the nasal photographs, 0.9% of the superior temporal photographs, and 0.9% of the inferior temporal photographs could not be graded. In these patients grading was based on the remaining fields.

Table 2 presents the prevalence of retinopathy. Overall, 32 patients (7.3%, 95% confidence interval (CI) 5.0 to 10.2) had diabetic retinopathy. One patient with retinopathy and 28 patients without retinopathy had reduced vision mainly from non-diabetic causes such as cataract or central retinal vein occlusion. The two patients with mature cataract subsequently underwent cataract extraction and neither had diabetic retinopathy.

The subjects with retinopathy had a lower BMI (mean 23.7 kg/m², 95% CI 23.7 to 24.7) than the non-retinopathy group (mean 25.0 kg/m², 95% CI 24.6 to 25.5), *p* = 0.031. There was no significant difference between any of the other clinical or biochemical parameters in those with and those without retinopathy.

None of the variables studied was found to be associated with diabetic retinopathy either on univariate or multiple logistic regression analysis.

Discussion

In Europeans the prevalence of retinopathy at the time of diagnosis of type 2 diabetes varies¹⁸⁻²⁴ and in the United Kingdom Prospective Diabetes Study (UKPDS) 35% of type 2

diabetic subjects were reported to have retinopathy at diagnosis.²³ This is considerably higher than that seen in our study (7.3%) and another from the same city but without retinal photography (6.9%).²⁴ It therefore appears that the prevalence of retinopathy at the time of diagnosis of diabetes in our patients is lower than that reported in the UK²³ and in the USA¹⁸⁻²¹ but is similar to that reported from Australia.¹⁸ One possible explanation is that there could be a longer period of undiagnosed (asymptomatic) type 2 diabetes in the UK. The mean age of the patients at the time of diagnosis in our study was considerably lower (men 45 (12) years, women 48 (12) years) than in the UKPDS study (men 52.4 (9) years, women 53.5 (8.8) years).²³ Another explanation could be that the time at which patients seek medical attention may vary between India and the UK as a result of the different health systems.

The limitations of this study are that it is clinic based and the sample size is small. Although referral bias cannot be excluded, this is unlikely to affect our results significantly because the patient profile at our centre is similar to that in an ongoing epidemiological study.¹⁰ The strength of this study is that it is based on retinal photography and standard grading techniques and, indeed, is one of the first studies of retinopathy using these techniques in south Asia.

In conclusion, using retinal photography we have found that the prevalence of diabetic retinopathy at diagnosis of diabetes in this south Indian clinic based study is 7.3% which appears to be lower than that reported in European subjects.

The authors thank Mr Prem for the retinal photography, Mr A K Mathai for statistical help, and Mrs G Malarvizhi for secretarial help in typing the manuscript.

- Klien R, Klein BE, Moss SE. Visual impairment in diabetes. *Ophthalmology* 1984;**91**:1-9.
- Vision problems in the United States. Data analysis: definitions, data sources, detailed data tables, analysis, interpretation. New York: National Society to Prevent Blindness, 1989:1-46.
- Caird FI, Pirie E, Ramsell TG. The natural history of diabetic retinopathy. In: *Diabetes and the eye*. Oxford: Blackwells, 1968:72-100.
- Dorf E, Ballantyne EF, Bennett PH, et al. Retinopathy in Pima Indians. Relationships to glucose level, duration of diabetes, age at diagnosis of diabetes and age at examination in a population with a high prevalence of diabetes mellitus. *Diabetes* 1976;**25**:554-60.
- Ballard DJ, Melton LJ, Dwyer MS, et al. Risk factors for diabetic retinopathy: a population based study in Rochester, Minnesota. *Diabetes Care* 1986;**9**:334-42.
- Rema M, Ponnaiya M, Mohan V. Prevalence of retinopathy in non insulin dependent diabetes mellitus at a diabetes centre in southern India. *Diabetes Res Clin Pract* 1996;**34**:29-36.
- Dandona L, Dandona R, Naduvilath, TJ, et al. Population based assessment of diabetic retinopathy in an urban population in southern India. *Br J Ophthalmol* 1999;**83**:937-40.
- Ramachandran A, Jali MV, Mohan V, et al. High prevalence of diabetes in an urban population in south India. *BMJ* 1988;**297**:587-90.
- King H, Auberti RE, Herman WH. Global burden of diabetes, 1995-2025: prevalence, numerical estimates and projections. *Diabetes Care* 1998;**21**:1414-31.
- Mohan V, Ravikumar RK, Shanthi Rani CS, et al. Intimal medial thickness of the carotid artery in South Indian diabetic and non-diabetic subjects: the Chennai Urban Population Study (CUPS). *Diabetologia* 2000;**43**:494-9.
- World Health Organization. *Diabetes mellitus. Report of a WHO study group*. Technical Report Series No 727. Geneva: WHO, 1985.
- Ferris FL, Kassoff A, Bresnick GH, et al. New visual acuity charts for clinical research. *Am J Ophthalmol* 1982;**94**:91-6.
- Moss SE, Meuer SM, Klein R, et al. Are seven standard photographic fields necessary for classification of diabetic retinopathy? *Invest Ophthalmol Vis Sci* 1989;**30**:823-8.
- Classification of diabetic retinopathy: detailed grading of stereo-color photographs. In: *Early treatment of diabetic*

- retinopathy study manual of operations*. Baltimore, MD: Early Treatment of Diabetic Retinopathy Study Coordinating Centre, Department of Epidemiology and Preventive Medicine, University of Maryland, 1985:18.
- 15 Early Treatment of Diabetic Retinopathy Study Research Group. Grading diabetic retinopathy from stereoscopic colour fundus photographs: an extension of the modified Airlie House classification. ETDRS Report. *Ophthalmology* 1991;**98**:786–806.
 - 16 Klein BEK, Davis MD, Segal P, *et al*. Assessment of severity and progression of diabetic retinopathy. *Ophthalmology* 1984;**91**:10–17.
 - 17 Varley H, Gowenlock AH, Bell M. *Practical clinical biochemistry*. London: Heinmann, 1980:600–1.
 - 18 Harris MI, Klein R, Welborn TA, *et al*. Onset of NIDDM occurs at least 4–7 years before clinical diagnosis. *Diabetes Care* 1992;**15**:815–9.
 - 19 Klein R, Klein BEK, Moss SE, *et al*. The Wisconsin Epidemiologic study of Diabetic Retinopathy. III. Prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years. *Arch Ophthalmol* 1984;**102**:527–32.
 - 20 Owens DR, Volund E, Jones D, *et al*. Retinopathy in newly presenting non insulin dependent (type 2) diabetic patients. *Diabetes Res Clin Pract* 1988;**9**:59–65.
 - 21 Klein R, Klein BEK, Moss SE, *et al*. The Beaver Dam study: retinopathy in adults with newly discovered and previously diagnosed diabetes mellitus. *Ophthalmology* 1992;**99**:58–62.
 - 22 Nagi DK, Pettitt DJ, Bennett PH, *et al*. Diabetic retinopathy assessed by fundus photography in Pima Indians with impaired glucose tolerance and NIDDM. *Diabetic Med* 1997;**14**:449–56.
 - 23 Kohner EM, Aldington SJ, Stratton IM, *et al*. Diabetic retinopathy at diagnosis of non insulin dependent diabetes mellitus and associated risk factors. United Kingdom Prospective Diabetes Study 30. *Arch Ophthalmol* 1998;**116**:297–303.
 - 24 Ramachandran A, Snehalatha C, Vijay V, *et al*. Diabetic retinopathy at the time of diagnosis of NIDDM in South Indian subjects. *Diabetes Res Clin Pract* 1996;**32**:111–4.



Prevalence of retinopathy at diagnosis among type 2 diabetic patients attending a diabetic centre in south India

Mohan Rema, Raj Deepa and Viswanathan Mohan

Br J Ophthalmol 2000 84: 1058-1060

doi: 10.1136/bjo.84.9.1058

Updated information and services can be found at:

<http://bjo.bmj.com/content/84/9/1058.full.html>

References

These include:

This article cites 17 articles, 6 of which can be accessed free at:

<http://bjo.bmj.com/content/84/9/1058.full.html#ref-list-1>

Article cited in:

<http://bjo.bmj.com/content/84/9/1058.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Retina](#) (1207 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>