

Latent TGF β binding protein-1 and fibrillin-1 in human capsular opacification and in cultured lens epithelial cells

Shizuya Saika, Takeshi Miyamoto, Takeshi Tanaka, Iku Ishida, Yoshitaka Ohnishi, Akira Ooshima

Department of Ophthalmology, Wakayama Medical University School of Medicine, Wakayama, Japan
S Saika
T Miyamoto
T Tanaka
I Ishida
Y Ohnishi

Department of Pathology
A Ooshima

Correspondence to:
Shizuya Saika, MD,
Department of Ophthalmology, Wakayama Medical University School of Medicine, 811-1 Kimiidera, Wakayama, 641-8509, Japan
shizuya@wakayama-med.ac.jp

Accepted for publication
10 May 2001

Table 1 Summary of the cases and the results

Case No	Age*	Sex	Indication for	
			Duration†	IOL removal
1	65	F	-	CCC‡
2	72	M	-	CCC
3	58	M	-	CCC
4	72	F	-	Lens dislocation
5	61	F	-	Enucleation
6	55	F	-	CCC
7	28	M	6 days	IOL dislocation
8	64	M	10 days	PVR
9	58	F	14 days	PVR
10	68	F	0.65 y	PVR
11	75	F	0.75 y	PVR
12	51	M	1 y	Malignant glaucoma
13	53	M	3 y	PVR
14	75	M	2.7 y	IOL dislocation
15	78	M	4 y	PVR
16	80	M	4 y	PVR
17	76	M	4.6 y	IOL dislocation
18	68	M	5 y	IOL dislocation
19	57	M	5.3 y	IOL dislocation
20	77	F	5.4 y	PVR
21	77	F	6.2 y	IOL dislocation
22	60	M	8 y	IOL dislocation
23	68	M	9 y	IOL dislocation
24	31	M	10 y	IOL dislocation
25	67	M	UK	PVR

*Age at the removal of the intraocular lens (IOL); †duration between implantation and explantation of the IOL.

y = year(s); ‡CCC = continuous circular capsulorhexis; PVR = IOL removal to facilitate inspection of the fundus during vitrectomy for proliferative vitreoretinopathy; capsulotomy, obtained by anterior capsulotomy during cataract surgery; - = negative; + = positive; ++ = strongly positive; +/- = negative overall with focal positivity; +/- = positive overall but focally negative.

Abstract

Background/aim—It was previously reported that collagenous extracellular matrix (ECM) in human capsular opacification contained isoforms of transforming growth factor β (TGF β). In the present study, the authors performed immunohistochemistry to examine whether ECM in human capsular opacification and in cultures of bovine lens epithelial cells (LECs) contained latent TGF β binding protein-1 (LTBP-1), TGF β 1 latency associated peptide (β 1-LAP), and fibrillin-1, a suspected ligand of LTBP-1 as well as a component of the extracellular microfibrillar apparatus. The aim of the study was to further clarify the mechanism of TGF β 1 deposition in ECM of capsular opacification.

Methods—Human capsular opacification specimens and uninjured lens capsules, as well as cultured bovine LECs, were processed for immunohistochemistry using antibodies against LTBP-1, β 1-LAP, fibrillin-1, and collagen type I.

Results—LTBP-1, β 1-LAP, and fibrillin-1 all were localised to the ECM in human capsular opacification. Uninjured lens epithelium stained for β 1-LAP, but not for

LTBP-1 and fibrillin-1. ECM deposited in confluent LEC cultures stained for LTBP-1, β 1-LAP, and fibrillin-1, while cultures with only sparse cellularity were unstained for LTBP-1 or fibrillin-1.

Conclusions—LECs upregulate LTBP-1 and fibrillin-1 during postoperative healing. LTBP-1, β 1-LAP, and fibrillin-1 colocalised to the ECM in capsular opacification and in confluent LEC cultures. TGF β 1 is considered to deposit in ECM in the large latent form. ECM secreted by LEC may function as a scavenger or repository of TGF β .

(Br J Ophthalmol 2001;85:1362-1366)

Capsular opacification is the most common complication of implantation of an intraocular lens (IOL). Lens epithelial cells (LECs) are responsible for this occurrence, depositing extracellular matrix (ECM) on the capsule.¹⁻³ LECs may transform to a mesenchymal-like phenotype and produce ECM components including collagens, the major components of capsular opacification, in some conditions.⁴⁻⁵ LEC proliferation in association with ECM production is considered part of the wound healing process in the crystalline lens.

Transforming growth factor β (TGF β) is pivotal in regulating cell behaviour during wound repair. Three isoforms (β 1, β 2, and β 3) are known in mammals.^{6,7} Each isoform of TGF β shows similar biological activities in vivo—that is, positive or negative regulation of cell growth, differentiation, and/or ECM production.⁸ Aqueous humour contains abundant TGF β 2.⁹ Moreover, healing LEC and ECM accumulation in capsular opacification reportedly contain TGF β isoforms.¹⁰⁻¹¹ Macrophages adhering to IOL are also immunoreactive for TGF β isoforms.¹² These isoforms are secreted as inactive complexes including small and large latent forms. Small latent TGF β is composed of active TGF β plus TGF β latency associated peptide (β 1-LAP); the large latent form consists of the small latent form of TGF β plus latent TGF β binding protein (LTBP). The ECM reportedly acts as a scavenger or repository for TGF β in various tissues.¹³⁻¹⁵ Fibrillin-1, a component of the extracellular microfibrillar apparatus, reportedly can bind to LTBP-1.^{16,17} These two proteins share a number of features including multiple EGF-like repeats, the RGD sequence, and an eight cysteine repeat motif unique to this family of proteins.^{16,17} Immunolocalisation of LTBP-1 and fibrillin-1, in capsular opacification has not

Table 2 Primary antibodies used

Antibody	Animal	Dilution	Source
Anti-LTBP-1	rabbit	×2000	Pharmingen ¹
Anti-β1-LAP	mouse*	×100	R & D system ²
Anti-fibrillin-1	goat	×200	Santa Cruz ³
Anti-bovine fibrillin-1	rabbit	×200	LSL ⁴
Anti-bovine collagen type I	rabbit	×200	LSL

LTBP-1 = latent TGFβ1 binding protein-1; β1-LAP = TGFβ1 latency associated peptide; TGFβ = transforming growth factor β; *mouse monoclonal.

1, San Diego, CA; 2, Minneapolis, MN; 3, Santa Cruz Biotechnology, Santa Cruz, CA, USA; 4, Tokyo, Japan.

been fully examined, although TGFβ isoforms are located to LECs and/or ECM there.¹¹ Fibrillin-1 is not a component of the normal lens, but may serve a connector between LTBP-1 and ECM in capsular opacification similarly to that in ECM of other tissues. We therefore examined immunohistochemically localisation and protein expression patterns of LTBP-1, β1-LAP, and fibrillin-1 in human




Figure 1 Localisation pattern of large transforming growth factor β binding protein-1 (LTBP-1) in human posterior capsule opacification. In (A) (case 1) and (B) (case 8), lens epithelium of an uninjured lens capsule (A, arrowheads) and a capsule 10 days after cataract surgery (B, arrowheads) lack LTBP-1. In (C) (case 17), extracellular matrix accumulation (asterisk) between the anterior lens capsule and the anterior surface of the intraocular lens shows immunoreactivity for LTBP-1. Lens cells are negative for this protein (arrowheads). In (D) (case 19), lens cells (arrowheads) and regenerated lenticular fibres of Sommering's ring (asterisk) unstained for LTBP-1. AC = anterior capsule; IOL = intraocular lens. Indirect immunostaining; bar = 20 μm.




Figure 2 Localisation pattern of β1 latency associated peptide (β1-LAP) in human posterior capsule opacification. In (A) (case 1), lens epithelial cells (arrowheads) of an uninjured lens capsule exhibit immunoreactivity for β1-LAP. In (B) (case 25), extracellular matrix accumulation (asterisk) around the edge of the capsulotomy (yellow arrow) shows weak immunoreactivity for β1-LAP, while lens cells (arrowheads) exhibit marked immunoreactivity. In (C) (case 17), extracellular matrix accumulation (asterisk) between the intraocular lens optic and the anterior capsule as well as lens cells (arrowheads) show strong β1-LAP immunoreactivity. AC = anterior capsule; IOL = intraocular lens. Indirect immunostaining; bar = 20 μm.

specimens with capsular opacification to further clarify the mechanism of TGFβ1 deposition in ECM of capsular opacification. Furthermore, we determined that cultured bovine LECs produce and deposit LTBP-1, β1-LAP, and fibrillin-1 in their ECM.

Materials and methods

CLINICAL SPECIMENS

Specimens obtained from 25 Japanese patients with capsular opacification were included in this study (Table 1). Three circular sections of the anterior capsule with LECs without opacification of the anterior subcapsule were obtained by circular capsulotomy during phacoextraction in patients with posterior subcapsular cataracts. Specimens from nine patients (cases 1–9) were obtained at the department of ophthalmology of Wakayama Medical University Hospital, while specimens from 16 patients (cases 10–25) were provided from the implant data system committee of the Japanese Society of Cataract and Refractive Surgery. The specimens were embedded in paraffin after being fixed in 10% formalin. Tissue sections 5 μm thick were cut, deparaffinised, and processed for immunohistochemistry as described below.

CELL CULTURE

Lens capsules were obtained from five bovine eyes. The tissues were explanted for outgrowth of LECs in Eagle's medium supplemented with antibiotics, antimycotics, and 10% fetal calf serum.¹⁸ After one passage, the cells (5×10^4 /ml/well) were seeded on chamber slides (Nunc, Naperville, IL, USA) and incubated for 3, 10, or 14 days. Three chambers were prepared for each culture condition. Cells then were dried, fixed in cold acetone, and processed for immunohistochemistry.

IMMUNOHISTOCHEMISTRY

Immunohistochemistry was performed as previously described.¹⁹ Deparaffinised sections and fixed cultured cells were treated with phosphate buffered saline (PBS) containing 3% skim milk and 5% calf serum to decrease non-specific binding of antibodies. Specimens then were allowed to react with the primary antibodies listed in Table 2 for 12 hours at 4°C. After washing in PBS, specimens were treated with peroxidase conjugated secondary antibodies for 4 hours at 4°C as previously reported.^{5 11 12} After a colour reaction with 3,3'-diaminobenzidine, specimens were embedded in balsam and observed by light microscopy. Negative controls were prepared by omission of the primary antibodies.

Results

SPECIMENS OF UNINJURED LENS CAPSULE AND CAPSULAR OPACIFICATION

LTBP-1 was immunohistochemically detected in ECM accumulation in the lesion of human fibrous capsular opacification, but not in uninjured LECs (Fig 1). Immunoreactivity for β1-LAP was marked in the cytoplasm of both uninjured LECs and healing lens cells in the fibrotic capsular opacification (Fig 2). The ECM of fibrotic capsules also was also reactive




Figure 3 Fibrillin-1 as a component of fibrous extracellular matrix in human posterior capsule opacification. In (A) (case 2), uninjured lens epithelium (arrowheads) lacks immunoreactivity for fibrillin-1. In (B) (case 18), extracellular matrix accumulation (asterisk) around the edge of anterior capsulotomy (yellow arrow) shows marked immunoreactivity for fibrillin-1. In (C) (case 17), weak immunoreactivity for fibrillin-1 (arrows) is detected surrounding proliferating lens epithelial cells, which lack immunoreactivity. AC = anterior capsule; IOL = intraocular lens. Indirect immunostaining; bar = 20 μ m.

for β 1-LAP (Fig 2). The regenerated lenticular structure of Sommering's ring and also healing lens cells were unstained for these two proteins (Fig 1D). LECs of uninjured lenses were negative for fibrillin-1, but LECs in healing capsular opacification were surrounded by fibrillin-1 deposition. Fibrillin-1 was immunohistochemically detected in ECM accumulation

on fibrous capsular opacification (Fig 3). Similar results were obtained in fibrillin-1 immunohistochemistry with two primary antibodies of different sources. No immunoreactivity was observed in negative control staining (not illustrated). There is no difference of the expression patterns of these three proteins among specimens of uninjured lens cells of various ages in given specimens.

LEC CULTURE

In sparsely cellular preconfluent cultures at day 3, almost no immunoreactivity for LTBP-1 was detected in LECs (Fig 4A). In early confluence, a small amount of LTBP-1 was detected extracellularly in a very fine fibrillar pattern, while in mature confluent cultures at day 14 immunoreactivity for LTBP-1 appeared as an organised extracellular network throughout the culture (Fig 4B and C). Immunoreactivity for β 1-LAP was detected in cells in sparsely cellular cultures and early confluent cultures (Fig 4D and E), although its immunoreactivity seemed weaker compared with that of *in vivo* LECs. Mature confluent cultures of LECs exhibited extracellular deposition of β 1-LAP (Fig 4F). Sparsely cellular LEC cultures did not show fibrillin-1 protein expression (Fig 4H), while LECs in early confluent cultures at day 10 were reactive for fibrillin-1 in a fine fibrillar pattern in the ECM (Fig 4I). The cells




Figure 4 Immunolocalisation of LTBP-1, β 1-LAP, and fibrillin-1 in the extracellular matrix accumulation in bovine lens epithelial cell cultures. In sparse, preconfluent, cultures at day 3, almost no immunoreactivity for LTBP-1 was detected in LECs (A). In early confluence, a small amount of LTBP-1 was detected extracellularly in a very fine fibrillar pattern (B), while in mature confluent cultures at day 14, immunoreactivity for LTBP-1 appeared to form an organised network throughout the culture (C). Immunoreactivity for β 1-LAP was detected in the cells of a sparsely cellular culture (D) and an early confluent culture (E). The mature confluent culture of LECs exhibited extracellular deposition of β 1-LAP (F). In sparsely cellular cultures, LECs did not express fibrillin-1 protein (H). In early confluent cultures at day 10, fibrillin-1 was detected in a fine fibrillar pattern in fibroblast ECM; the cells themselves were unstained (I). Mature confluent LEC cultures showed a condensed fibrillin-1 structure (J). No specific immunoreactivity was observed in negative controls with anti-goat IgG (G) or anti-rabbit IgG but no primary antibody (K). LTBP-1 = large transforming growth factor β binding protein-1; β 1-LAP = β 1-latenancy associated peptide; LEC = lens epithelial cells. Indirect immunostaining; bar = 50 μ m.

themselves remained unstained. Mature confluent LEC cultures showed a condensed fibrillar structure of fibrillin-1 in their ECM, remaining the cell cytoplasm negative (Fig 4J). Fibrillin-1 staining patterns did not completely overlap with immunoreactivity for LTBP-1 or β 1-LAP. At all time points during culture, collagen type I was detected in the cytoplasm of LECs in a fine granular pattern, but not in the extracellular space (not illustrated). No specific immunoreactivity was observed in negative controls with anti-goat IgG or anti-rabbit IgG, but no primary antibody (Fig 4G and K).

Discussion

In the present study, immunostaining patterns for LTBP-1, β 1-LAP, and fibrillin-1 suggested that TGF β 1 and fibrillin-1 were colocalised to the ECM in human capsular opacification. Uninjured LECs were positive for β 1-LAP but negative for LTBP-1 and fibrillin-1, indicating that in vivo LECs upregulate LTBP-1 and fibrillin-1 during healing after surgery. Fibrillin-1 is considered to trap TGF β 1 through binding to LTBP-1. We previously reported that myofibroblastic transformation of in vivo LECs during lens wound healing is associated with expression of various ECM macromolecules including collagens.⁵ LTBP-1 and fibrillin-1 are considered to be upregulated similarly to other ECM components expressed by LECs in association with myofibroblastic transformation, in that cuboidal LECs in a specimen 10 days postoperatively or in Sommering's ring were negative for LTBP-1. Alternatively, ECM surrounding postoperative LECs may provide some signal(s) for expression of LTBP-1 (or for conversion from the small latent form to the large latent form of TGF β 1). Indeed, fibronectin reportedly regulates the epithelial mesenchymal conversion in cells including LECs, as shown by an α smooth muscle actin expression pattern.^{20 21} Similar mechanisms may work in upregulation of LTBP-1. Fibrillin-containing microfibrillar apparatus is considered to play an important part in ECM organisation during lens tissue repair, as occurs in other tissues.^{22 23} Localisation of LTBP-1 and β 1-LAP to fibrous capsular opacification suggests that it functions in storing or scavenging latent TGF β in this ECM compartment.

We further confirmed by cell culture that bovine LECs deposit LTBP-1 and β 1-LAP as well as fibrillin-1 in the ECM. Deposition of LTBP-1, β 1-LAP, and fibrillin-1 also has been demonstrated in other cell types in culture.¹³⁻¹⁷ In sparsely cellular LEC cultures, LTBP-1 and β 1-LAP were not detected in the cell layer, but in confluent LEC culture they were observed in the ECM in a fibrillar network or a granular pattern. Cell signals induced by confluence apparently stimulate deposition of LTBP-1, β 1-LAP, and fibrillin-1 into the extracellular space. In mature confluent cultures, fibrillin-1 containing structures did not completely overlap with immunoreactivity for LTBP-1 and β 1-LAP, indicating that LTBP-1 and β 1-LAP are deposited in areas where no fibrillin-1 is present in addition to being bound to

fibrillin-1. Other depositing mechanism may also be present. In this stage of cultures, however, collagen type I, the major ECM component in fibrous capsular opacification, was detected only in cytoplasm, not in the ECM on the cells; thus, LTBP-1 and β 1-LAP are not associated with type I collagen. Secretion of β 1-LAP and fibrillin-1 by cultured LECs might not be similar to that in in vivo LECs; cytoplasmic immunoreactivity for both proteins were marked in the cytoplasm of healing LECs in vivo, but weak or not seen in cell culture. Further study is needed to clarify the regulation mechanism of secretion of these proteins.

LECs proliferate following cataract surgery even though they are exposed to TGF β s that have a growth inhibitory effect on them.²⁴⁻²⁹ ECM molecules may regulate (enhance) post-operative proliferation of LECs in vivo.³⁰ We have reported that lens epithelial cells in capsular opacification express matrix metalloproteinases and their inhibitors.³¹ Degradation activity of fibrillin-1 containing ECM may modulate LEC growth by regulating release of TGF β from the matrix.

In conclusion, ECM accumulation surrounding LECs in vivo postoperatively and those in vitro deposits TGF β 1 presumably through binding LTBP-1 to fibrillin-1, and may modulate LEC behaviour not only by providing cell adhesion scaffolds but also by storage or scavenging of this growth factor.

Grant support: Supported in part by Wakayama Medical Award for Young Researchers, 1999.

The authors thank members of the IOL Implant Data System Committee of the Japanese Society of Cataract and Refractive Surgery for their cooperation.

- Apple DJ, Solomon KD, Tetz, *et al.* Posterior capsule opacification. *Surv Ophthalmol* 1992;37:73-116.
- Saika S, Ohmi S, Kanagawa R, *et al.* Lens epithelial cell outgrowth and matrix formation on intraocular lenses in rabbit eyes. *J Cataract Refract Surg* 1996;22:835-40.
- Meacock WR, Spalton DJ, Stanford MR. Role of cytokines in the pathogenesis of posterior capsule opacification. *Br J Ophthalmol* 2000;84:332-6.
- Ramaekers FCS, Osborn M, Weber K, *et al.* Identification of the cytoskeletal proteins in lens-forming cells, a special epithelial cell type. *Exp Cell Res* 1998;127:309-27.
- Saika S, Kawashima Y, Miyamoto T, *et al.* Immunolocalization of prolyl 4-hydroxylase subunits, α -smooth muscle actin, and extracellular matrix components in human lens capsules with lens implants. *Exp Eye Res* 1998;66:283-94.
- Massague J. The transforming growth factor- β family. *Annu Rev Cell Biol* 1990;6:597-641.
- Miller DA, Pelton RW, Derynck R, *et al.* Transforming growth factor β : a family of growth regulatory peptides. *Ann NY Acad Sci* 1990;593:208-17.
- Levine JH, Moses HL, Gold LI, *et al.* Spatial and temporal patterns of immunoreactive transforming growth factor β 1, β 2, and β 3 during excisional wound repair. *Am J Pathol* 1993;143:368-80.
- Cousins SW, McCabe MM, Danielpour D, *et al.* Identification of transforming growth factor-beta as an immunosuppressive factor in aqueous humor. *Invest Ophthalmol Vis Sci* 1991;32:2201-11.
- Gordon-Thomson C, de Jongh RU, Hales AM, *et al.* Differential cataractogenic potency of TGF- β 1, - β 2, and - β 3 and their expression in the postnatal rat eye. *Invest Ophthalmol Vis Sci* 1998;39:1399-409.
- Saika S, Miyamoto T, Kawashima Y, *et al.* Immunolocalization of TGF- β 1, - β 2 and - β 3, and TGF- β receptors in human lens capsules with lens implants. *Graefes Arch Clin Exp Ophthalmol* 2000;238:283-4.
- Saika S, Miyamoto T, Okada Y, *et al.* Transforming growth factor- β isoform proteins in cells and matrix deposits on intraocular lenses. *J Cataract Refract Surg* 2000;26:709-15.
- Dallas SL, Miyazono K, Skerry TM, *et al.* Dual role for the latent transforming growth factor- β binding protein in storage of latent TGF- β in the extracellular matrix and as a structural matrix protein. *J Cell Biol* 1995;131:539-49.
- Nakajima Y, Miyazono K, Kato M, *et al.* Extracellular fibrillar structure of latent TGF β binding protein-1: role in

- TGF β -dependent endothelial-mesenchymal transformation during endocardial cushion tissue formation in mouse embryonic heart. *J Cell Biol* 1997;136:193-204.
- 15 Tripale J, Keski-Oja J. Growth factors in the extracellular matrix. *FASEB J* 1997;11:51-9.
 - 16 Raghunath M, Unsold C, Kubitscheck U, et al. The cutaneous microfibrillar apparatus contains latent transforming growth factor- β binding protein (LTBP-1) and is a repository for latent TGF- β . *J Invest Dermatol* 1998;111:559-64.
 - 17 Ashworth JA, Kielty CM, McLeod D. Fibrillin and the eye. *Br J Ophthalmol* 2000;84:1312-7.
 - 18 Saika S, Yamanaka O, Kawashima Y, et al. OPC-15161 suppresses the proliferation of Tenon's capsule fibroblasts and the production of type I collagen and fibronectin stimulated by TGF- β 1 in vitro. *Curr Eye Res* 1998;17:933-90.
 - 19 Saika S, Shiraishi A, Saika S, et al. Role of lumican in the corneal epithelium during wound healing. *J Biol Chem* 2000;275:2607-12.
 - 20 Magro G, Frassetto F, Travali S, Lanzafame S. Immunohistochemical expression and distribution of α 2 β 1, α 6 β 1, α 5 β 1 integrins and their extracellular ligands, type IV collagen, laminin and fibronectin in palmar fibromatosis. *Gen Diagn Pathol* 1997;143:203-8.
 - 21 Kurosaka D, Kato K, Oshima T, et al. Extracellular matrices influence alpha-smooth muscle actin expression in cultured porcine lens epithelial cells. *Curr Eye Res* 1999;19:260-3.
 - 22 Raghunath M, Tschodrich-Rotter M, Sasaki T, et al. Confocal laser scanning analysis of the association of fibulin-2 with fibrillin-1 and fibronectin define different stages of skin regeneration. *J Invest Dermatol* 1999;112:97-101.
 - 23 Watson REB, Griffiths CEM, Craven NM, et al. Fibrillin-rich macrofibrils are reduced in photoaged skin. Distribution at the dermal-epidermal junction. *J Invest Dermatol* 1999;112:782-7.
 - 24 Arteaga CL, Coffey Jr, RJ, Dugger TC, et al. Growth stimulation of human breast cancer cells with anti-transforming growth factor β antibodies: evidence for negative autocrine regulation by transforming growth factor β 1. *Cell Growth Diff* 1990;1:367-74.
 - 25 Kurosaka D, Nagamoto T. Inhibitory effect of TGF- β 2 in human aqueous humor on bovine lens epithelial cell proliferation. *Invest Ophthalmol Vis Sci* 1994;35:3408-12.
 - 26 Wang X-J, Greenhalgh DA, Bickenbach JR, et al. Expression of a dominant-negative type II transforming growth factor β (TGF- β) receptor in the epidermis of transgenic mice blocks TGF- β -mediated growth inhibition. *Proc Natl Acad Sci USA* 1997;94:2386-91.
 - 27 Rakic J-M, Galand A, Vrensen GFJM. Separation of fibers from the capsule enhances mitotic activity of human lens epithelium. *Exp Eye Res* 1997;64:67-72.
 - 28 Ashcroft GS, Yang X, Glick AB, et al. Mice lacking Smad3 show accelerated wound healing and an impaired local inflammatory response. *Nat Cell Biol* 1999;1:260-6.
 - 29 Saika S, Miyamoto T, Okada Y, et al. Smad translocation and growth suppression in lens epithelial cells during wound repair in mice. *Exp Eye Res* 2001;72:679-86.
 - 30 Oharazawa H, Ibaraki N, Lin LR, et al. The effects of extracellular matrix on cell attachment, proliferation and migration in a human lens epithelial cell line. *Exp Eye Res* 1999;69:603-10.
 - 31 Kawashima Y, Saika S, Miyamoto T, et al. Matrix metalloproteinases and tissue inhibitors of metalloproteinases of fibrous human lens capsules with intraocular lenses. *Curr Eye Res* 2001;21:962-7.



Latent TGF β binding protein-1 and fibrillin-1 in human capsular opacification and in cultured lens epithelial cells

Shizuya Saika, Takeshi Miyamoto, Takeshi Tanaka, et al.

Br J Ophthalmol 2001 85: 1362-1366

doi: 10.1136/bjo.85.11.1362

Updated information and services can be found at:

<http://bjo.bmj.com/content/85/11/1362.full.html>

These include:

References

This article cites 29 articles, 11 of which can be accessed free at:

<http://bjo.bmj.com/content/85/11/1362.full.html#ref-list-1>

Article cited in:

<http://bjo.bmj.com/content/85/11/1362.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>