

WORLD VIEW

Blindness and visual impairment in the Americas and the Caribbean

B Muñoz, S K West

Br J Ophthalmol 2002;**86**:498–504Series editors:
WV Good and S RuitSee end of article for
authors' affiliationsCorrespondence to:
Beatriz Munoz, Wilmer
Eye Institute, Room 118,
Johns Hopkins University,
600 N Wolfe Street,
Baltimore, MD 21287,
USAAccepted for publication
18 September 2001**Aim:** To summarise available data on the prevalence and causes of visual impairment and blindness in the Americas and the Caribbean.**Methods:** The published literature was searched in Medline and LILACS using the following key words: blindness, visual impairment, prevalence. Articles were reviewed, and the references of the articles were also searched for relevant articles, which were also reviewed.**Results:** Using the mortality in children under the age of 5 as an indicator, the overall prevalence of childhood blindness (in the under age 15 group) for the region was estimated at 0.45/1000, with the majority (67%) living in countries with mortality of children under age 5 above 30/1000 live births. Corneal opacities were more common in countries where the under 5 year mortality are above 30/1000 live births and retinopathy of prematurity (ROP) was an important cause in countries with intermediate death rates. For adults, overall blindness rates were not estimated because of the social, economic, and ethnic diversity in the region. The primary causes of visual loss in adults in the Americas were age related eye diseases, notably cataract and glaucoma in the African-American and Hispanic populations, and age related macular degeneration in the white population. Uncorrected refractive error was a significant cause of decreased vision across ages, ethnic groups, and countries.**Conclusion:** More data are needed on the magnitude and causes of visual loss for the Caribbean and Latin American countries. Rates of blindness and visual loss from available data within these countries are widely disparate. Prevention and control of avoidable blindness needs to be an ongoing focus in this region.

North, Central, and South America, including the Caribbean, cover a region characterised by extreme diversity. Geographically, the region includes the driest desert in the world (Atacama Desert in Chile, with annual precipitation less than 0.5 inches) and the wettest location outside Antarctica (Mt Waialeale, Hawaiian Islands, with annual precipitation of 37 feet). Population-wise, the region includes the two largest cities in the world, Mexico City with 26 million and Sao Paulo, Brazil, with 24 million, and at the same time island countries where most of the population is in the rural areas in small communities.

The extremes are evident as well in the variation in socioeconomic and health indicators for the populations in this region. Canada has one of the lowest infant death rates in the world, while Haiti has among the worst, over 10 times the rate in Canada.¹ Within the high income countries, there is also quite a disparity, along racial/education/income lines. For example, within the United States, African-Americans have over twice the infant mortality of the white population.

Thus, there should be little surprise at finding a large variation in blindness and visual impairment rates across countries in the Americas. The major surprise, however, is how little is known about the magnitude of the problem in various countries within the region. Data on blindness and visual loss in children are based on models derived from estimates correlated with child mortality. Except for very localised surveys in the United States, there are almost no population based data on adults. This paper summarises the literature, including reports where they could be obtained, on the prevalence and causes of visual loss for the Americas.

METHODS

The published literature was searched in Medline using the following key words: blindness, visual impairment, prevalence. All articles were reviewed, and the references of the

articles were also searched for relevant articles, which were also reviewed. A search of a database in Latin America, LILACS (Latin American and Caribbean Literature on Health Sciences), was also carried out using similar key words in Spanish and Portuguese. In addition, reports on blindness surveys in Latin America and the Caribbean were requested through Pan American Health Organization and the Prevention of Blindness representatives. Finally, people thought to have relevant data were approached for permission to cite their findings, if the data were not yet published.

Because of the absence of data on childhood blindness, we adopted the model of Gilbert and Foster,² which extrapolates an estimated prevalence rate according to the childhood mortality reported for the country. There are no good comparable models for estimating rates of adult blindness and visual impairment, so the paucity of data in this area is all too evident. There is social, economic, and ethnic diversity in most of these countries which will greatly affect the rates and causes of blindness and visual loss,³ so it is simply not possible to extrapolate rates from surveys in one country to surveys in another. Therefore, we present adult prevalence rates specific for the country (or area within the country) where the study was conducted.

RESULTS**Childhood blindness
Magnitude of blindness**

Information on the prevalence of blindness in children is scarce and difficult to obtain because of its relatively low occurrence. Population based studies that could yield results with reasonable precision are very expensive to conduct, so alternative ways to assess the magnitude of the problem have been used. Based on data from blind registers in Europe and the few available population based surveys from Asia and

Table 1 Magnitude of blindness in children age 0–15, estimated as a function of under 5 years mortality (1999)

Region	Mortality for children under 5 years/1000 live births	Estimated prevalence of blindness/1000	Countries	Estimated population under 15 years (in millions)	Estimated number of blind
North America	30 and under	0.3	Canada, USA	67.0	20 100
Central America	30 and under	0.3	Costa Rica, Panama	2.1	630
	31–94	0.6	Belize, El Salvador, Guatemala, Honduras, Mexico, Nicaragua	44.7	26 820
South America	30 and under	0.3	Argentina, Chile, Colombia, Paraguay, Uruguay, Venezuela	38.7	11 610
	31–94	0.6	Bolivia, Brazil, Ecuador, Guyana, Peru	62.8	37 680
Caribbean	30 and under	0.3	Aruba, Bahamas, Barbados, Cayman Island, Cuba, Dominica, Grenada, Jamaica, Netherlands Antilles, Puerto Rico, St Kitts Nevis, St Lucia, St Vincent, Trinidad Tobago, Virgin Islands	5.1	1530
	31–94	0.6	Dominican Republic	2.7	1620
	95–170	0.9	Haiti	3.0	2700
Total		0.45	All countries	226.1	102 690

Africa that included children, Foster *et al* proposed an algorithm to estimate the magnitude of blindness in children age 0–15 using the mortality in children under 5 as an indicator.² Table 1 shows estimates of the magnitude of prevalence of childhood blindness for the Americas and the Caribbean by applying Foster's paradigm to population and mortality data from the year 1999.^{1–4}

Approximately 100 000 out of the 226 million children under 15 years of age were blind in the Americas and the Caribbean. The overall prevalence was 0.45/1000, the majority of them (67%) living in countries with under 5 death rates above 30/1000 live births.

Causes

As is the case for the estimation of the magnitude of childhood blindness, accurate data (population based) on the distribution of causes of blindness and severe visual impairment in childhood are not available. An alternative approach is to examine the causes in an identifiable group of blind children. Blind registers and schools for the blind have served as the primary source to look at the aetiology, although these two sources are subject to selection and survival bias. Recognising they are likely to under-represent the poor, and blind children with other disabilities, they could provide a starting point to determine the main causes.⁵ To be able to compare results across studies, the World Health Organization (WHO) recommends a standard methodology for reporting causes of childhood blindness based on two criteria—*anatomical site of the abnormality and underlying aetiology*.⁶ However, only the more recent studies have used this approach making it difficult to summarise all available data using the recommended criteria. In this review we present causes using the anatomical site whenever possible. The four more frequently reported causes of blindness and severe visual impairment for schools for the blind throughout countries from the regions are shown in Table 2. The observed distributions agree with the patterns described by others in terms of the specific causes being associated with different levels of economic development.^{2, 5, 17}

Corneal opacities were more common in countries where the under 5 year death rates are above 30 per 1000 live births. In Bolivia, Dominican Republic, and Peru around 20% of the blindness is attributable to corneal opacities in contrast with none or very few cases in the United States, Chile, and Argentina (Table 2). Although it was difficult to establish the aetiology of the corneal opacity, evidence from studies elsewhere suggests that the main causes are likely to be corneal ulceration after measles, herpes simplex infection, vitamin A deficiency, and ophthalmia neonatorum.^{16, 18}

Retinopathy of prematurity (ROP) was an important cause of blindness in countries with intermediate mortality where the survival of low birthweight infants has increased but technology for neonatal care is problematic. ROP as a cause of

blindness was rare in Bolivia, Dominican Republic, and Peru, very probably because of the lower survival rate of premature babies^{2, 10} (Table 2). More than one third of the blind or severely visually impaired children attending blind schools in Argentina, Cuba, and Paraguay had ROP as a primary cause of visual loss. In contrast, a lower frequency was observed in studies from the United States (8%–19%), probably the result of improvements in intensive neonatal care.

Cataract in childhood as a cause of blindness was reported from all countries, with a lower frequency in the United States, where only one of the four studies reported any childhood cataract as a frequent cause. The most frequent underlying causes of childhood cataract are congenital acquired rubella and genetic disease.¹⁹

Adult blindness and visual impairment Magnitude

The prevalence of blindness and visual impairment based on population surveys, using standard acuity test that have been published in the past three decades from the Americas and the Caribbean are shown in Table 3. The definition of blindness and visual impairment varied across studies; when available, both the WHO blindness definition (visual acuity 20/400 or worse) and legal blindness in the United States (visual acuity 20/200 or worse) were presented. Only a few studies also incorporated blindness based on visual field loss alone, and we did not include those data. Some studies reported presenting acuity (with habitual correction) in the better eye instead of best corrected acuity. Consequently, the prevalence of blindness and visual impairment for those studies tended to be higher because they included uncorrected refractive error.

Blindness and visual impairment estimates for North America came from the United States, because of the absence of data from Canada, except based on self report.^{37, 38} Within the United States, the aetiology and prevalence of blindness varied by ethnic group, location (rural/urban), and year of the survey, with more recent studies reporting lower rates. Temporal trends that affect blindness rates include increasing demand for cataract surgery to correct visual loss, using techniques with much better visual rehabilitation.³⁹

Studies carried out before 1990 reported blindness rates (best corrected visual acuity 20/200 or worse) in white people age 40 and older ranging from 0.5% in Beaver Dam to 1.1% in the rural community of Mud Creek Valley.^{30, 31} For black people, the Baltimore Eye Survey estimated a blindness rate of 1.5%.²⁹ In 1994, the Salisbury Eye Evaluation examined black and white people, and the reported prevalence of blindness among ages 65–84 was 1.7% and 0.5% respectively. A recent study of Mexican-Americans age 40 and older estimated blindness rates at 0.3%, although most of the population was less than age 60. Age adjusted comparisons showed rates in Mexican-Americans midway between white and black people.³⁵

Table 2 Causes of childhood blindness as reported from schools for the blind

Country	Year	Location	No	Four most frequent causes of blindness and severe visual impairment				%
				Category 1	Category 2	Category 3	Category 4	
Argentina ⁷	1993	School for the blind	573	ROP	Other retina	Optic nerve	Cataract	8
Bolivia ⁸	1988	School for the blind	78	Cornea	Retina	Cataract	Optic nerve	10
Brazil ⁹	1998	School for the blind	267	Glaucoma	Cataract	Chorioretinitis	Optic atrophy	12
Chile ⁷	1992	10 schools for the blind	94	Other retina	ROP	Optic nerve	Cataract	9
Colombia ¹⁰	1991–6	Three schools for the blind*	70	ROP	—	—	—	—
Cuba ¹⁰	1991–6	All school for the blind 2 regions*	51	ROP	—	—	—	—
Dominican Republic ⁷	1992	School for the blind	142	Cataract	Cornea	Glaucoma	Other retina	10
Ecuador ¹⁰	1991–6	All school for the blind*	73	ROP	—	—	—	—
Guatemala ¹⁰	1991–6	All school for the blind*	108	Cataract	Optic nerve	Glaucoma	Retina	15
Jamaica ¹¹	1991–6	School for the blind	36	ROP	—	—	—	—
Paraguay ¹⁰	1990	All school for the blind*	202	Cornea	Glaucoma	Cataract	Retinal degeneration	12
Peru ¹²	1986	School for the blind	220	Cataract	Other retina	Optic nerve	ROP	9
Uruguay ⁷	1981–95	Low vision clinic	762	Cortical visual impairment	ROP	Optic atrophy	nystagmus	7
USA ¹³	1996–7	Alabama school for the blind	123	Optic atrophy	Cataract	Albinism	Congenital malformations	12
USA ¹⁴	1998	20 schools for the blind	2,553	Cortical	ROP	Optic nerve	Albinism	7
USA ¹⁶	1998	School for visually impaired	62	ROP	Optic atrophy	Retinitis pigmentosa	Optic nerve hypoplasia	13

*Other causes of blindness not reported.

In Central and South America, the studies from Brazil, Chile, and Peru suggested a higher prevalence of blindness than the prevalence reported among Hispanics in studies in North America.^{21–23} Comparing two surveys, both carried out in the late 1980s using similar age groups (40 and older), 2.9% of the population had bilateral best corrected acuity 20/200 or worse in Peru²³ compared to 0.9% and 1.9% for white and black people in the Baltimore Eye Survey.^{28–29}

The only data on blindness prevalence available for the Caribbean comes from the Barbados Eye Study.²⁰ In this predominantly black population, age adjusted blindness rates were twice as high as the rates reported for black people in the Baltimore Eye Survey (3.0% v 1.5%).

Causes of visual loss

As Table 3 shows, the primary causes of visual loss in adults in the Americas were age related eye diseases, notably cataract and glaucoma in the African-American and Hispanic populations, and age related macular degeneration in the white population. In younger age groups, and in high risk populations, diabetic retinopathy was a leading cause of visual loss.

The excess risk of blindness in the Latino-American surveys was probably the result of unoperated cataract. An estimated 72%–74% of the blindness in Chimbote, Peru and Campinas, Brazil²³ was cataract related compared to the highest proportion reported for the United States, 27% in black people in the Baltimore Eye Survey. In another two studies in Chile²¹ and Brazil²² the proportion of blindness due to cataract was 55% and 62% respectively, also much higher than in North America.

In the Caribbean, data from the Barbados Eye Study showed that cataract and glaucoma were responsible for almost 60% of the blindness,²⁰ with each contributing equally. Prevalence of blindness from glaucoma was higher than the prevalence reported for black people in the United States. This high proportion of blindness due to glaucoma is consistent with the high prevalence of open angle glaucoma reported in the region.^{40–43}

Uncorrected refractive error is an important cause of visual loss in all ages, and one not captured by studies reporting rates of best corrected acuity loss. Table 4 shows the significance of uncorrected refractive error as a cause of visual impairment in several population groups. A significant number of those who have visual loss would not be classified as impaired with appropriate refractive correction. Studies in children in the United States, Chile, and Colombia suggested that almost half of the children with decreased visual acuity would no longer be impaired with appropriate refractive correction.^{44–46} Similarly, data from adults showed that close to 20% of the people aged 40 and older with presenting acuity 20/200 or worse improved to better than 20/200 after refraction.²⁸ In Proyecto VER, 73% of the individuals with presenting acuity worse than 20/40 improved after refraction.³⁵

There are populations, particularly in Central and South America, living in extremely impoverished and politically disenfranchised circumstances that continue to foster eye disease long since banished from most of the Americas. Six countries report onchocerciasis in localised areas affecting the poorest of the poor (Brazil, Colombia, Ecuador, Guatemala, Mexico, and Venezuela).^{48–49} The Onchocerciasis Elimination Program for the Americas (OEPA) has been active in these countries since 1990. According to the last report from the ninth Inter-American Conference on Onchocerciasis, semiannual treatments with ivermectin have been successful, accomplishing good to excellent coverage.⁴⁸ In fact, Colombia, Ecuador, and Mexico have treatment programmes that reportedly have reached full coverage.⁴⁸ However, distribution programmes in Guatemala and Brazil are performing below their annual treatment objectives, and in Venezuela the national programme has just completed plans for mass treatment

Table 3 Prevalence and causes of blindness and visual impairment in the adult population

Country	Year	Study design	Race	Age group	No	Type*	Visual acuity in the better eye	Prevalence	Primary causes
Barbados ²⁰	1987–92	Simple random sample	Black	40–84	4303	B	≤20/200 ≤20/400 <20/60→20/400	3.0% 1.7% 5.9%	Glaucoma, cataract, retinal/choroidal diseases
Chile ²¹	1988	Community based rural	Hispanic	≥60	664	B	≤20/200	4.4%	Cataract, glaucoma, retinal (not AMD)
Brazil ²² Sao Paulo	1986–7	Stratified random sample	Hispanic	≥65	120	P	≤20/400	6.8%	Cataract, myopic degeneration, glaucoma
Brazil ²³ Capinas	1986–7	Community based	Hispanic	≥50	7452	B	≤20/200	2.8%	74% Cataract was involved
Brazil ²⁴ Porto Alegre	1992–4	Community based	Hispanic	All ages	All residents neighbourhood	B	≤20/200	0.1%	Cataract, retinal, diabetic retinopathy
Peru ²³ Chimbote	1986–7	Community based	Hispanic	≥40	12193	B	≤20/200	2.9%	72% Cataract was involved
USA Framingham, MA ²⁵	1973–5	Community based	White	≥55 75–84	2631 394	B	≤20/200	0.6% 2.0%	Cataract, ARMD, glaucoma, diabetic retinopathy
National Health and Nutrition Examination ^{26,27}	1974–5	Multistage random sample	Black White	25–74	250 2660	P	≤20/200	0.2% 0.4%	Not reported
Hispanic Health and Nutrition Examination ^{26,27}	1982–4	Multistage random sample	Cuban Mexican Puerto Rican	25–74	396 1381 513	P	≤20/200	1.0% 0.9% 0.2%	Not reported
Baltimore, MD ^{28,29}	1985–8	Stratified random sample	White Black	≥40	2913 2395	B	≤20/400 ≤20/200 <20/60→20/200 ≤20/400 ≤20/200 <20/60→20/200	0.5% 0.9% 1.3% 0.9% 1.5% 1.9%	ARMD, cataract Cataract, glaucoma
Mud Creek Valley, KY ³⁰	1988	Community based	White	≥40	1136	B	≤20/200 <20/60→20/200	1.1% 1.5%	Cataract, ARMD, diabetic retinopathy, glaucoma
Beaver Dam, WI ³¹	1987–8	Community based	99% White	≥43	4897	B	≤20/200 <20/60→20/200	0.5% 0.8%	Not reported
3 communities in MA, CT, IO ³²	1990	Community based	94% White	≥71	5335	P	≤20/200 <20/60→20/200 <20/40→20/60	4.7% 14.4% 14.1%	Not reported
Salisbury, MD ^{33,34}	1994–5	Community based random sample	White Black	65–84	1853 666	B	≤20/400 ≤20/200 <20/60→20/200 ≤20/400 ≤20/200 <20/60→20/200	0.2% 0.5% 1.2% 0.8% 1.7% 2.6%	Cataract, ARMD, diabetic retinopathy Cataract, diabetic retinopathy, glaucoma
Tucson and Nogales, AZ ^{35,36}	1997–9	Community based random sample	Hispanic	≥40	4774	B	≤20/400 ≤20/200 <20/60→20/200	0.2% 0.3% 0.9%	Cataract, ARMD, diabetic retinopathy

*P = presenting acuity, B = best corrected.

Table 4 Proportion of decreased vision as a result of uncorrected refractive error

Country	Year	Type of survey	Race	Age group	Definition	% of decreased visual acuity due to uncorrected refractive error
Children						
Chile, La Florida ⁴⁴	1998	Cluster random sample	Hispanic	5–15	≤20/40	55
Colombia, Medellin ⁴⁵	1993	20% random sample from 88 000 children screened all public schools	Hispanic	5–14	Not specified	50
USA, California ⁴⁶	1979–88	First graders in a school district	(>50% Hispanic)	6–7	≤20/40	65
Adults						
Chile ⁴⁷	1990	Community screening for blindness	Hispanic	≥60	≤20/200	48
Brazil, ²² Sao Paulo	1986–7	Stratified random sample	Hispanic	≥65	≤20/25	49
Brazil, ²³ Campinas	1986–7	Community based	Hispanic	≥50	≤20/200	34
Brazil, ²⁴ Porto Alegre	1992–4	Community screening for blindness	Hispanic	All ages	≤20/200	44
Peru, ²³ Chimbote	1986–7	Community based	Hispanic	≥40	≤20/200	15
USA Baltimore, MD ²⁹	1985–8	Stratified random sample	White	≥40	≤20/200	21
			Black	≥40	≤20/200	18
USA, Salisbury, MD ³⁴	1994–5	Community based random sample	White	65–84	<20/40	34
			Black	65–84	<20/40	23
USA, Tucson and Nogales, AZ ³⁵	1997–9	Community based random sample	Hispanic	≥40	<20/40	73

programmes in all affected communities to be active by the year 2001.⁴⁹

In the past 15 years, there have been studies showing the existence of pockets of endemic trachoma in Brazil, Mexico, and Peru.^{50–64} In South America, the Amazonian region and the bordering states of northwest Brazil seem to be the most affected, with prevalence of active disease ranging from 5% to 42%.^{53 54 56 58} While trachoma has been found in urban areas in other states of Brazil,^{51 52 55} the low prevalence of active disease (<10%) in children, and no signs of scars and trichiasis in adults were reported, suggested blinding trachoma was not yet a problem.

In Central America, trachoma is endemic in pockets of southern Mexico and Guatemala.^{59–61} Two population based studies reported prevalence of active disease in children around 25%, and in some of the communities almost all examined adults (aged 40 and older) had signs of conjunctival scars. The area affected is mostly rural, inhabited by farmers of Mayan descent with limited access to public services including health care.

Trachoma was endemic until recently in the native Americans in the United States, with trachoma reported in 1982 at 1% among Navaho children.⁶⁵ Trachoma was reported as a cause of blindness in a recent review of the medical records from the Indian Health Services.⁶⁶ There may well be other indigenous populations with trachoma in the Americas, but surveys have not been carried out.

DISCUSSION

The region of the Americas encompasses an extremely diverse set of nations. Even within nations, the populations can range from well educated, wealthy members who have access to the most sophisticated eye care available, to very poor minority populations, isolated with little or no access to, or understanding of how to access, reasonable eye care. A simple summary of data on visual loss in the Americas does not do justice to this diversity.

First of all, it is clear that more data are needed on the magnitude and causes of visual loss, particularly in the Caribbean nations and other countries of central and southern America. Some surveys are currently planned, or the data collected but not yet available. Clearly, from the socioeconomic indicators and the estimates of childhood blindness, there will be pronounced differences within different countries. Such data are essential to plan and implement national programmes for blindness prevention. Where there may be populations at high

risk for onchocerciasis or trachoma, special surveys to identify these groups and national will to plan targeted interventions are a special priority.

Secondly, attention must be paid to the growing problem of ROP in children. Control of childhood blindness is a high priority set by the Vision 2020 agenda.⁶⁷ Although the total number of blind children is lower than that of adults, the personal and social costs are higher,⁶⁸ because of the life expectancy. In fact, the number of “blind years” due to childhood blindness is estimated to be similar to the number of “cataract blind years” in adults.¹⁷ The proportion of blindness due to ROP in the developed countries ranges from 6–18%.¹⁰ However, in middle income countries, there is an epidemic of ROP, due in part to the increased survival of low birthweight infants where there is suboptimal neonatal care.¹⁰

The problem of blindness due to cataract will continue to be an issue in the future, as the population ages and risk factors for cataract, such as diabetes, become even more prevalent. The rates of blindness and visual loss are higher in Latin America than in comparable age groups in the United States, due in large part to the proportion blind from cataract. The data from Chimbote and Campinas^{23 69} suggested that only two thirds of people who were offered cataract surgery were operated. No apparent reasons for refusal were reported, although fear of surgery, acceptance of visual loss as a consequence of old age, and poor outcomes were reasons in other studies.^{70–75} Data from another study in US based Hispanics suggested that economic barriers, such as no insurance, were barriers to having cataract surgery.⁷⁶ The estimated cataract surgery rate per million population per year is fivefold higher in North America compared to the rest of the Americas.^{77 78} It is clear that the barriers blocking access to cataract surgery need to be identified, and improvements instituted, in order to reduce the burden of blindness in Latin America.

One of the main causes of blindness in younger people is diabetic retinopathy. Prevalence of diabetes in the Hispanic and native American populations is very high; a recent study in a population of Mexican-Americans in the United States reported one in five people over age 40 had diabetes, and almost half had evidence of diabetic retinopathy.⁷⁹ This includes the 15% of people with diabetes who were not aware of their condition. Adequate control and early treatment of diabetes have been shown to be cost effective for controlling diabetic retinopathy, but this will continue to be a significant public health problem in populations where access to treatment is poor.

Finally, the problem of uncorrected refractive error as a cause of visual loss is starting to be recognised. While the

impact on quality of life of people with uncorrected refractive error is not as large as the impact from cataract or other ocular pathology,⁸⁰ it is easily avoidable. The size of the problem is underappreciated because many of the studies report visual loss only after best correction. Uncorrected refractive error is not only an issue in children, where visual screening ought to be part of preparation for schooling. Older people have significant uncorrected, or poorly corrected, refractive error as well.

In summary, there are few data on the rates of blindness and visual loss within countries in the Americas, and the data which do exist shows widely disparate rates, as expected. Although many countries have socioeconomic markers that place them as high to middle income countries, there are populations within these countries that do not enjoy the same freedom from visual loss as their compatriots. Prevention and control of avoidable blindness needs to be an ongoing focus in this region.

ACKNOWLEDGEMENTS

A research grant EY11283 from the National Eye Institute supported the project. Dr West is a Research to Prevent Blindness senior scientific investigator.

Authors' affiliations

B Muñoz, S K West, Dana Center, Wilmer Eye Institute, Johns Hopkins University, Baltimore, MD, USA

REFERENCES

- 1 **World Bank Group**. World development indicators data base. <http://devdata.worldbank.org/external/CPProfile.asp> (25 July 2001)
- 2 **Foster A**, Gilbert CE. Epidemiology of childhood blindness. *Eye* 1992;**6**:173–6.
- 3 **Viet HH**, Schwab IR. Social economic development in the prevention of global blindness. *Br J Ophthalmol* 2001;**85**:653–7.
- 4 **Statistics Division and Population Division of the United Nations Secretariat**. Indicators of young and elderly populations. <http://srch0.un.org/plweb/cgi> (26 July 2001).
- 5 **Gilbert CE**. Childhood blindness. In: Gordon JJ, Minassian DC, Weale R, eds. *The epidemiology of eye disease*. 1st ed. London: Chapman and Hall, 1998:181–207.
- 6 **Gilbert CE**, Foster A, Negrel D, et al. Childhood blindness: a new form for recording causes of visual loss in children. *Bull World Health Organ* 1993;**71**:485–9.
- 7 **Gilbert CE**, Canovas R, Canovas RK, et al. Causes of blindness and severe visual impairment in children in Chile. *Devel Med Child Neurol* 1994;**36**:326–33.
- 8 **Foster A**. Childhood blindness. *Eye* 1988;**2**(Suppl):S27–36.
- 9 **Leite LA**. Causas de cegueira entre os alunos em curso no Instituto Benjamin Constant no ano de 1996. *Rev bras Oftalmol* DATE??;**57**:619–623.
- 10 **Gilbert C**, Rahi J, Eckstein M, et al. Retinopathy of prematurity in middle-income countries. *Lancet* 1997;**350**:12–14.
- 11 **Moriarty BJ**. Childhood blindness in Jamaica. *Br J Ophthalmol* 1988;**72**:65–7.
- 12 **Rojas JR**, Lavado L, Echegaray L. Childhood blindness in Peru. *Ann Ophthalmol* 1990;**22**:423–5.
- 13 **Wilkinson ME**, Stewart IW. Iowa's pediatric low vision services. *J Am Optom Assoc* 1996;**67**:397–402.
- 14 **Decarlo DK**, Nowakowski R. Causes of visual impairment among students at the Alabama School for the Blind. *J Am Optom Assoc* 1999;**70**:647–52.
- 15 **Steinkuller PG**, Du L, Gilbert C, et al. Childhood blindness. *J AAPOS* 1999;**3**:26–32.
- 16 **Mets MB**. Childhood blindness and visual loss: an assessment at two institutions including a "new" cause. *Trans Am Ophthalmol Soc* 1999;**97**:653–96.
- 17 **Gilbert C**, Foster A. Childhood blindness in the context of Vision 2020—the right to sight. *Bull World Health Organ* 2001;**79**:227–32.
- 18 **Whitcher JP**, Srinivasan M, Upadhyay MP. Corneal blindness: a global perspective. *Bull World Health Organ* 2001;**79**:214–21.
- 19 **Gilbert C**, Rahi J, Eckstein M, et al. Hereditary disease as a cause of childhood blindness: regional variation. Results from blind school studies undertaken in countries of Latin America, Asia and Africa. *Ophthalmic Genet* 1995;**16**:1–10.
- 20 **Hyman L**, Wu SY, Connell AMS, et al. Prevalence and causes of visual impairment in the Barbados eye study. *Ophthalmology* 2001;**108**:1719–20.
- 21 **Vicencio C**, Verdaguier J, Lama G, et al. Estudio ocular en San Vicente de Tagua-Tagua. *Arch Chil Oftal* 1989;**2**:103–7.
- 22 **Medina NH**, Monteiro de Barros O, Muñoz EDH, et al. Morbidade ocular em idosos da cidade de Sao Paulo-SP, Brasil. *Arq Bras Oftal* 1993;**56**:276–83.
- 23 **Newton-Kara J**, Contreras F, Campos MA, et al. Screening and surgical intervention results from cataract-free-zone projects in Campinas, Brazil and Chimbote, Peru. *Int Ophthalmol* 1990;**X**:156–64.
- 24 **Esteves JF**, Domingues CG, Borges LPK, et al. Prevalence and causes of blindness in Porto Alegre, Brazil. *Arq Bras Oftal* 1996;**59**:244–7.
- 25 **Leibowitz HM**, Krueger DE, Maunder LR, et al. The Framingham eye study monograph. *Surv Ophthalmol* (Suppl) 1980.
- 26 **Lam BL**, Lee DJ, Gomez-Marin O. Prevalence of usual-corrected binocular distance visual acuity impairment in Hispanic and non-Hispanic adults. *Ophthalmic Epidemiol* 2000;**7**:73–83.
- 27 **Lee DJ**, Gomez-Marin O, Lam BL. Prevalence of uncorrected binocular distance visual acuity in Hispanic and non-Hispanic adults. *Ophthalmology* 1998;**105**:552–60.
- 28 **Tielsch JM**, Sommer A, Witt K, et al. Blindness and visual impairment in an American urban population. *Arch Ophthalmol* 1990;**108**:286–90.
- 29 **Sommer A**, Tielsch JM, Katz J, et al. Racial differences in the cause-specific prevalence of blindness in East Baltimore. *N Engl J Med* 1991;**325**:1412–17.
- 30 **Dana RM**, Tielsch JM, Enger C, et al. Visual impairment in a rural Appalachian community. *JAMA* 1990;**264**:2400–5.
- 31 **Klein R**, Klein BEK, Linton KLP, et al. The Beaver Dam Eye Study: visual acuity. *Ophthalmology* 1991;**98**:1310–15.
- 32 **Salive ME**, Guralnik J, Christen W, et al. Functional blindness and visual impairment in older adults from three communities. *Ophthalmology* 1992;**99**:1840–7.
- 33 **Rubin GS**, West SK, Muñoz B. A comprehensive assessment of visual impairment in a population of older Americans. The SEE study. Salisbury Eye Evaluation Project. *Invest Ophthalmol Vis Sci* 1997;**38**:557–68.
- 34 **Muñoz B**, West SK, Rubin GS, et al. The SEE Study Team. Causes of blindness and visual impairment in a population of older Americans. *Arch Ophthalmol* 2000;**118**:819–25.
- 35 **Muñoz B**, West SK, Rodriguez J, et al. Blindness, visual impairment and the problem of uncorrected refractive error in a Mexican-American population: proyecto ver. *Invest Ophthalmol Vis Sci* 2002;**43**:615–22.
- 36 **Rodriguez J**, Sanchez R, Muñoz B, et al. Causes of blindness and visual impairment in a population based sample of Hispanics. *Ophthalmology* (in press).
- 37 **Naeyaert KM**, Grace G. Prevalence and causes of blindness and visual impairment in Canada. *J Vis Imp* 1990:361–3.
- 38 **Hameed TK**, Hodge W, Buhmann R. An inventory of information on blindness and visual impairment in Canada. *Can J Ophthalmol* 2001;**36**:175–87.
- 39 **Brian G**, Taylor HR. Cataract blindness—challenges for the 21st century. *Bull World Health Organ* 2001;**79**:249–56.
- 40 **Mason RP**, Kosoko O, Wilson R, et al. National survey of the prevalence and risk factors of glaucoma in St Lucia, West Indies. Part I. Prevalence findings. *Ophthalmology* 1989;**96**:1363–8.
- 41 **Leske MC**, Connell AMS, Schachat AP, et al. The Barbados Eye Study group. The Barbados Eye Study group. Prevalence of open angle glaucoma. *Arch Ophthalmol* 1994;**112**:821–9.
- 42 **Leske MC**, Connell AM, Wu SY, et al. Incidence of open-angle glaucoma: the Barbados Eyes Studies. *Arch Ophthalmol* 2001;**119**:89–95.
- 43 **Alemañy-Martorell J**, Tejeiro-Fernandez A. Survey of blind people and main causes of blindness in members of the National Association for the Blind. *Rev cuba Oftalmol* 1994;**7**:68–76.
- 44 **Maul E**, Barroso S, Muñoz SR, et al. Refractive error study in children: results from La Florida, Chile. *Am J Ophthalmol* 2000;**129**:445–54.
- 45 **Rodriguez MDLA**, Castro-Gonzales M. Salud visual de escolares en Medellín, Antioquia, Colombia. *Bol Oficina Sanit Panam* 1995;**119**:11–14.
- 46 **Choi TB**, Lee DA, Oelrich FO, et al. A retrospective study of eye disease among first grade children in Los Angeles. *J Am Optom Assoc* 1995;**66**:484–8.
- 47 **Vicencio C**, Verdaguier JI, Verdaguier J, et al. Proyecto San Vicente de Tagua-Tagua zona libre de catarata. *Arch. Chileno oftalmologia* 1991;**48**:56–64.
- 48 **World Health Organization**. *Weekly epidemiological record*. Geneva: WHO, 19 January 2001;**76**:17–24.
- 49 **Richards FO**, Miri ES, Katabarwa M, et al. The Carter Center's assistance to river blindness control program: establishing treatment objectives and goals for monitoring ivermectin delivery systems on two continents. *Am J Trop Med* 2001;**65**:108–14.
- 50 **Medina NH**, Oliveira MB, Tobin S, et al. The prevalence of trachoma in preschool children in Olimpia, Guaraci and Cajobi, Sao Paulo, Brazil. *Trop Med Parasitol* 1992;**43**:121–3.
- 51 **Luna EJ**, Medina NH, Oliveira MB, et al. Epidemiology of trachoma in Bebedouro State of Sao Paulo, Brazil: prevalence and risk factors. *Int J Epidemiol* 1992;**21**:169–77.
- 52 **Nobrega MJ**, Bonomo PPO, Scarpi MJ, et al. Prevalence of trachoma in pre-school children in the suburb of Joinville, state of Santa Catarina, Brazil. *Arq bras Oftalmol* 1993;**53**:13–17.
- 53 **Morschbacher R**, Scarpi MJ. Prevalence of trachoma in Xingu indian park. *Arq bras oftalmol* 1996;**59**:83–7.
- 54 **Garrido C**, Guidugli T, Campos M. Laboratorial and clinical trachoma study in an indian population from the Brazilian Amazon. *Arq bras oftalmol* 1999;**62**:132–8.
- 55 **Caligaris LSA**. Occurrence of trachoma in preschool population in Paz da Bela Vista sub-district, Sao Paulo city, 1995. [Dissertation] Sao Paulo: Universidad de Sao Paulo, 1998.
- 56 **Carvalho RC**, Falcao R, Cohen J, et al. Prevalence of trachoma in school children in Manaus, Amazonas, Brazil. *Arq Bras oftalmol* 1997;**60**:243–7.

- 57 **Scarpi MJ**, Silva RJM, Ferreira IA, *et al.* Trachoma prevalence in Palmares, state of Pernambuco. *Arq bras oftalmol* 1990;**53**:171–4
- 58 **Scarpi MJ**, Plut RCA, Arruda HO. Prevalence of trachoma. In the village of Mocambo, State of Ceara, Brazil. *Arq bras oftalmol* 1989;**52**:177–9.
- 59 **Taylor HR**, Millan-Velasco F, Munoz EC, Ruvalcaba AM, Greene BM. Trachoma in Chiapas, Mexico. *Rev Int Trach* 1983;**60**:19–27.
- 60 **Taylor HR**, Millan-Velasco F, Sommer A. The ecology of trachoma: an epidemiological study of trachoma in Southern Mexico. *Bull World Health Organ* 1985;**63**:559–67.
- 61 **Wilson M**, Keyvan-Larijani E, Millan-Velasco F, *et al.* The epidemiology of trachoma in Chiapas (Mexico). *Rev Int Trach Pathol Ocul Trop Subtrop Sante Publique* 1987;**64**:159–74.
- 62 **Wong Cam C**, Nava San Roman A, Calderon Cabada M, *et al.* Ophthalmological findings in the Shipiba community of Santa Teresita: trachoma, cytologic demonstration. *Rev peru Oftalmol* 1986;**12**:14–16.
- 63 **Tenorio A**, Wong C, Tobaru L, *et al.* Trachoma in urban and urban-marginal population of Peru. *Arch Peru Oftalmol* 1987;**2**:26–7.
- 64 **Wong Cam C**, Nava A, Calderon M, *et al.* Afecciones oculares y causas de ceguera en la selva peruana. *Arch Peru Oftalmol* 1986;**1**:11–16.
- 65 **Friederich R**. Eye disease in the Navajo indians. *Ann Ophthalmol* 1982;**14**:38–40.
- 66 **Rearwin DT**, Tang JH, Hughes JW. Causes of blindness among Navajo Indians: an update. *J Am Optom Assoc* 1997;**68**:511–17.
- 67 **Resnikoff S**, Pararajasegaram R. Blindness prevention programmes: past, present, and future. *Bull World Health Organ* 2001;**79**:222–6.
- 68 **Rahi JS**, Gilbert CE, Foster A, *et al.* Measuring the burden of childhood blindness. *Br J Ophthalmol* 1999;**83**:387–88.
- 69 **Jose NK**, Contreras F, Campos MA, *et al.* Screening and surgical intervention results from cataract free zone projects in Campinas, Brazil and Chimbote, Peru. *Int Ophthalmol* 1990;**14**:155–64.
- 70 **Snellingen T**, Shrestha BR, Gharti MP, *et al.* Socioeconomic barriers to cataract surgery in Nepal: the South Asian cataract management study. *Br J Ophthalmol* 1998;**82**:1424–8.
- 71 **Rabiú MM**. Cataract blindness and barriers to uptake of cataract surgery in a rural community of northern Nigeria. *Br J Ophthalmol* 2001;**85**:776–80.
- 72 **Brilliant GE**, Lepkowski JM, Zurita *et al.* Social determinants of cataract surgery utilization in south India. *Arch Ophthalmol* 1991;**109**:584–9.
- 73 **Fletcher AE**, Donoghue M, Devavaram J, *et al.* Low uptake of eye services in rural India: a challenge for programs of blindness prevention. *Arch Ophthalmol* 1999;**117**:1393–9.
- 74 **Vaidyanathan K**, Limburg H, Foster A, *et al.* Changing trends in barriers to cataract surgery in India. *Bull World Health Organ* 1999;**77**:104–9.
- 75 **Courtright P**, Kanjaloti S, Lewallen S. Barriers to acceptance of cataract surgery among patients presenting to district hospitals in rural Malawi. *Trop Geogr Med* 1995;**47**:15–18.
- 76 **Broman AT**, Muñoz B, Rodriguez J, *et al.* Factors associated with cataract surgery in a population-based study of Hispanics: proyecto VER. *Invest Ophthalmol Vis Sci* 2001;**42**:s534.
- 77 **Apple DG**, Ram J, Foster A, *et al.* Elimination of cataract blindness: a global perspective entering the new millenium. *Surv Ophthalmol* 2000;**45**:S1–196.
- 78 **Foster A**. Cataract and “Vision 2020—the right to sight” initiative. *Br J Ophthalmol* 2001;**85**:635–9.
- 79 **West SK**, Klein R, Rodriguez J, *et al.* Diabetes and diabetic retinopathy in a Mexican-American population. *Diabetes Care* 2001;**24**:1204–9.
- 80 **Muñoz B**, Broman AT, Rodriguez J, *et al.* Self reported disability associated with glaucoma, cataract, and diabetic retinopathy in a population-based study of Hispanics. *Invest Ophthalmol Vis Sci* 2001;**42**:s711.



Have your say

eLetters

If you wish to comment on any article published in the *British Journal of Ophthalmology* you can send an eLetter using the eLetters link at the beginning of each article. Your response will be posted on *British Journal of Ophthalmology* online within a few days of receipt (subject to editorial screening).

www.bjophthalmol.com



Blindness and visual impairment in the Americas and the Caribbean

B Muñoz and S K West

Br J Ophthalmol 2002 86: 498-504

doi: 10.1136/bjo.86.5.498

Updated information and services can be found at:

<http://bjo.bmj.com/content/86/5/498.full.html>

References

These include:

This article cites 58 articles, 18 of which can be accessed free at:

<http://bjo.bmj.com/content/86/5/498.full.html#ref-list-1>

Article cited in:

<http://bjo.bmj.com/content/86/5/498.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>