

LABORATORY SCIENCE

Architecture of arachnoid trabeculae, pillars, and septa in the subarachnoid space of the human optic nerve: anatomy and clinical considerations

H E Killer, H R Laeng, J Flammer, P Groscurth

Br J Ophthalmol 2003;**87**:777–781

Aims: To describe the anatomy and the arrangement of the arachnoid trabeculae, pillars, and septa in the subarachnoid space of the human optic nerve and to consider their possible clinical relevance for cerebrospinal fluid dynamics and fluid pressure in the subarachnoid space of the human optic nerve.

Methods: Postmortem study with a total of 12 optic nerves harvested from nine subjects without ocular disease. All optic nerves used in this study were obtained no later than 7 hours after death, following qualified consent for necropsy. The study was performed with transmission (TEM) and scanning electron microscopy (SEM).

Results: The subarachnoid space of the human optic nerve contains a variety of trabeculae, septa, and stout pillars that are arranged between the arachnoid and the pia layers of the meninges of the nerve. They display a considerable numeric and structural variability depending on their location within the different portions of the optic nerve. In the bulbar segment (ampulla), adjacent to the globe, a dense and highly ramified meshwork of delicate trabeculae is arranged in a reticular fashion. Between the arachnoid trabeculae, interconnecting velum-like processes are observed. In the mid-orbital segment of the orbital portion, the subarachnoid space is subdivided, and can appear even loosely chambered by broad trabeculae and velum-like septa at some locations. In the intracanalicular segment additionally, few stout pillars and single round trabeculae are observed.

Conclusion: The subarachnoid space of the human optic nerve is not a homogeneous and anatomically empty chamber filled with cerebrospinal fluid, but it contains a complex system of arachnoid trabeculae and septa that divide the subarachnoid space. The trabeculae, septa, and pillars, as well as their arrangement described in this study, may have a role in the cerebrospinal fluid dynamics between the subarachnoid space of the optic nerve and the chiasmal cistern and may contribute to the understanding of the pathophysiology of asymmetric and unilateral papilloedema. All the structures described are of such delicate character that they can not even be visualised with high resolution magnetic resonance imaging (MRI).

See end of article for authors' affiliations

Correspondence to: H E Killer, Kantonsspital Aarau, Augenklinik, Aarau, Switzerland; Killer@ksa.ch

Accepted for publication 12 November 2002

The optic nerve is a white matter tract of the central nervous system that extends through the optic canal into the orbit. It is enveloped by the meninges and is surrounded by cerebrospinal fluid that enters the subarachnoid space via the chiasmal cistern.

Anatomically the nerve can be divided into orbital and intracanalicular portions. The widest part of the orbital portion is adjacent to the ocular globe and is called the bulbar segment. It is followed by the mid-orbital segment (Fig 1).

Although many studies on the cranial meninges are available, relatively few deal exclusively with the meninges of the optic nerve and the structures, such as trabeculae and septa, within its subarachnoid space. Anderson described arachnoid trabeculae stretching between the arachnoid layer and the pia layer in the subarachnoid space in the optic nerve of both humans and monkeys.¹ Hayreh observed dense fibrous adhesions between the arachnoid and the pia layer in the region of the optic canal, without rendering histological information concerning type or distribution of these structures in other regions of the subarachnoid space.^{2,3}

In our previous investigation on the meninges of the human optic nerve, which demonstrated lymphatic vessels in the dura mater, we noted a difference in the type and distribution of arachnoid trabeculae in the different portions of the optic nerve.⁴ In the present study we focused on the architecture of trabeculae and septa in the subarachnoid space as well as on their relations to the different portions of the optic nerve. Owing to the "cul de sac" anatomy of the optic nerve, the

cerebrospinal fluid dynamic in its subarachnoid space may be strongly influenced by the presence of trabeculae and septa. Knowledge on the precise anatomy of the structures as well as their arrangement in the subarachnoid space may help to solve questions concerning the pathophysiology of asymmetric and unilateral papilloedema.

MATERIAL AND METHODS

The 12 optic nerves from nine patients used in this study were removed no longer than 7 hours after death following qualified consent for necropsy. Access to the optic nerve and the globe was obtained through the orbital roof. Before removing the globe, the subarachnoid space of the intracranial segment of the optic nerve was carefully injected with fixative using a 26 gauge needle. Subsequently, the orbital portion of the optic nerve and the globe were carefully dissected in situ from the surrounding tissues and the specimens were fixed with 2.5% glutaraldehyde (0.1 M cacodylate buffer). In two cadavers the optic nerves and globes were removed with the intact sphenoid bone in order to examine the intracanalicular segment of the nerve.

Scanning electron microscopy (SEM)

The globe and optic nerve were fixed for at least 1 week in a solution of 2% glutaraldehyde (0.1 M cacodylate buffer). For examination of the intracanalicular segment the fixed specimens were decalcified with 0.5% HNO₃ for 24 hours.




Figure 1 Schematic drawing of the optic nerve demonstrating the location of the (a) bulbar segment (containing trabeculae), (b) mid-orbital segment (containing septae and pillars), and (c) canalicular portion (containing pillars). The bulbar segment and the mid-orbital segment together form the orbital portion (modified according to Liu and Kahn¹³).

Transverse sections of the bulbar, mid-orbital, and canalicular segment were then dehydrated in an acetone series, dried by the critical point method (CO₂), mounted on aluminium stubs, and sputtered with gold (15 nm). The specimens were studied with an SEM 505 (Philips, Eindhoven, Netherlands).

Transmission electron microscopy (TEM)

Small fragments (approximately 1 mm³) were cut from the bulbar and mid-orbital segment and postfixed for 1 day within 1% OsO₄ (0.1 M cacodylate buffer). Subsequently the specimens were dehydrated in a series of alcohol and embedded into Epon by routine procedure. Semithin sections (approximately 2 µm) were cut from each block and stained with toluidine blue to identify the meninges. Ultrathin sections (approximately 50 nm) were contrasted with uranyl acetate and lead citrate and studied with a CM 100 (Philips, Eindhoven, Netherlands).

RESULTS

Bulbar segment of the orbital optic nerve portion

The bulbar segment of the optic nerve showed a distinct enlargement of the subarachnoid space (ampulla) which was detectable even macroscopically. SEM examination revealed abundant round shaped arachnoid trabeculae bridging the subarachnoid space, which were anchored in the arachnoid and pia layers without broadening (Fig 2A). The trabeculae were often branched to form a delicate network (Fig 2B). The profile of the trabeculae varied between 5 µm and 7 µm. Thin lamellae of highly flat cells were located in a veil-like pattern between adjacent trabeculae. Occasionally, larger trabeculae containing one or two blood vessels were found within the trabecular network (Fig 2C).

Each trabecula was covered by flat leptomeningeal cells with smooth surface and slender processes (Fig 2D). The continuity of the cells was occasionally interrupted by intercellular gaps and fenestrations with a diameter between 0.2 µm and 1 µm. TEM revealed that each trabecula was surrounded by a complete sheath of leptomeningeal cells (Fig 3A). Adjacent trabeculae were connected by thin cytoplasmic bridges of leptomeningeal cells running freely through the subarachnoid space. The oval shaped nucleus of the leptomeningeal cells was rich in heterochromatin and protruded




Figure 2 SEM appearance of the subarachnoid space in the bulbar segment. (A) Overview of the subarachnoid space showing the complex network of trabeculae. The arrows point to veil-like cytoplasmic extensions between adjacent trabeculae (bar = 150 µm). (B, C) Delicate subarachnoid space network formed by branching trabeculae (bar = 50 µm). The arrow points to a trabeculum with a blood vessel. Note again the veil-like cytoplasmic extensions connecting adjacent trabeculae (bar = 2 µm). Surface of trabeculae covered by flattened cells with distinct intercellular clefts and fenestrations (bar = 0.2 µm).




Figure 3 TEM morphology of arachnoid trabeculae in the bulbar segment. The trabeculae are completely surrounded by arachnoid cells. (A) Note the thin cytoplasmic bridges (arrows) connecting adjacent trabeculae (bar = 2 μ m). (B) At higher magnification multiple layers of cytoplasmic extensions from arachnoid cells are detectable at the surface of a trabeculum. Note the desmosomes (arrows) between the arachnoid cells and the broad basal lamina-like extracellular matrix supporting the leptomenigeal cells (arrowheads) (bar = 0.25 μ m).

into the subarachnoid space. The leptomenigeal cells formed, with their slender processes, either single or multiple layers of surface lining cells (Fig 3B). In multiple layers the cells were attached to each other by well defined desmosomes. The size of the intercellular space varied distinctly and it was often widened forming small lacunae between leptomenigeal cells. The basal portion of the leptomenigeal cells was supported either by a fine fibrillar extracellular matrix resembling a basal lamina or by delicate collagenous fibrils in close contact with the cell membrane. The centre of the trabeculae was filled by densely packed collagenous fibrils which were arranged in small bundles (Fig 3A). Occasionally, single fibroblasts were detected within the trabeculae in close contact with the collagenous fibrils. Blood capillaries, lymphatic vessels, and nerve fibres were not present.

Mid-orbital segment of the orbital optic nerve portion

The subarachnoid space was distinctly smaller than in the bulbar segment. It contained numerous broad septa running in various directions (Fig 4A and B). They divided the subarachnoid space into chambers that were connected to each other by large perforations within the septae and by interseptal spaces. In addition, single round shaped pillars were detectable which, in contrast with trabeculae, ended with broadened ends at the pial and dural surface of the subarachnoid space and had a diameter of 10–30 μ m (Fig 4C). Both septa and pillars were covered by flattened cells with smooth surfaces displaying numerous intercellular fenestrations of varying size (Fig 4D). TEM morphology of leptomenigeal cells covering septa and pillars was comparable to that of trabeculae in the bulbar segment. They formed either single or multiple layers and were connected to each other by well defined desmosomes. The centre of the septa and pillars contained numerous collagenous fibrils and single fibroblasts but no blood vessels, lymphatics, or nerve fibres.

Intracanalicular portion of the optic nerve

Within the optic canal the subarachnoid space was continuous over large distances (Fig 5A). The centre of the canal demonstrated one or two large pillars with a size of approximately 0.5 mm containing one or two blood vessels. The other parts of the




Figure 4 SEM morphology of the subarachnoid space in the mid-orbital segment. (A) Overview of the subarachnoid space subdivided by well defined septa (arrows), N = optic nerve (bar = 150 μ m). (B) Arachnoid septum within the subarachnoid space; note the septal perforations connecting adjacent subarachnoid space chambers (bar = 25 μ m). (C) Pillar bridging the subarachnoid space and extending into the dura layer of the arachnoid (bar = 30 μ m). (D) Detail of a septum with typical intercellular clefts (bar = 0.5 μ m).

intracanalicular segment showed either single pillars with a diameter of approximately 25 μ m which expanded at the dural and pial attachment of the arachnoid layer (Fig 5B), or delicate round shaped and slightly curved trabeculae of approximately 5 μ m diameter (Fig 5C). At the orbital opening of the canal the trabeculae were more numerous running in parallel and bridging the subarachnoid space in oblique direction (Fig 5C). Again the surface of the pillars and trabeculae was covered by flat, smooth surfaced cells with small intercellular clefts. Occasionally single leucocytes were found closely attached to the leptomenigeal cells (Fig 5D).

DISCUSSION

The present study has used SEM and TEM to describe in detail for the first time both the fine anatomy of trabeculae, septa, and pillars in the subarachnoid space of the human optic nerve and their relation to location within the different portions of the nerve.

The morphology of the cranial and spinal meninges and of the subarachnoid space have been described in previous studies on dogs, cats, rats, and humans.^{1–8} In order to find appropriate terminology for the structures transversing the spinal subarachnoid space, Parkinson used the terms arachnoid septa, trabeculae, and “rough strands.”⁹ A classification of cranial arachnoid trabeculae in stout, columnar or sheet-like




Figure 5 SEM morphology of the subarachnoid space in the intracanalicular segment. (A) Overview of the subarachnoid space; A = ophthalmic artery, N = optic nerve (bar = 15 μm) (the optic nerve is turned 180°). (B) Pillars (arrows) and trabeculae (arrowheads) transversing the subarachnoid space (bar = 30 μm). (C) Trabeculae within the subarachnoid space found at the orbital opening of the optic canal (bar 30 μm). (D) Leucocyte at the surface of a pillar in the intracanalicular segment (bar = 5 μm).

types was suggested by Delmas.¹⁰ The same paper describes a close association of the trabeculae with blood vessels, a finding that also was confirmed by Alcolado.¹¹ Detailed information on anchorage of the fibroblasts that form the arachnoid trabeculae in the arachnoid and the pia layer is provided by Haines.¹² In contrast with the vast literature on the cranial and spinal meninges, there is a scant anatomical literature on the meninges of the optic nerve.

Arachnoid trabeculae as described in the cranial and spinal subarachnoid space have also been described in the subarachnoid space of the human optic nerve by Anderson, however without reference to different types and to distribution.¹ In the present study on human optic nerves we observed that the subarachnoid space of the optic nerve is widest in the bulbar segment of the intraorbital portion. There, a multitude of delicate arachnoid trabeculae are arranged in a reticular fashion. Some of the larger trabeculae contain blood vessels (Fig 2B) as described previously by Anderson.¹ In contrast with Hayreh's observation,^{2,3} we found the densest arrangement of trabeculae in the bulbar segment and not in the canalicular portion. In a paper dedicated to cerebrospinal fluid pressure in the subarachnoid space of the optic nerve, Liu suggested that the majority of trabeculae occur in the mid-orbital segment of the intraorbital portion.¹³ Neither Hayreh's nor Liu's study,

however, provide histological evidence that would support their assumptions. Concerning the architecture and location of the trabeculae and septa, our study shows that the subarachnoid space narrows in the mid-orbital segment where the delicate character of the trabeculae changes gradually into broader septa and stout pillars that subdivide the subarachnoid space into small compartments. Within the intracanalicular portion the subarachnoid space is extremely narrow. Only a few large pillars and trabeculae but no septa are observed. As described by other authors, free cells can be observed on the arachnoid surface and on the surface of the trabeculae.^{14,15}

In consideration of this anatomy the subarachnoid space of the optic nerve cannot be regarded as a homogeneous space filled with cerebrospinal fluid, but rather as a multichambered and subdivided tubular system with a blind end (cul de sac) behind the ocular globe.

Cerebrospinal fluid is mainly produced by the choroid plexus epithelium and the ependymal cells of the ventricular system from where it flows into interconnecting chambers—namely, the cisterns and the subarachnoid spaces of the central nervous system. Cerebrospinal fluid circulation and direction of flow have been studied with radioisotopes and other tracers injected into the cerebrospinal fluid. It is generally agreed that there is a bulk circulation of fluid from the site of origin to the site of absorption—that is, from the ventricles to the arachnoid villi in the cranial subarachnoid spaces. The mechanism by which cerebrospinal fluid is propelled and its circulatory route are not fully understood but are probably influenced by the release of newly produced cerebrospinal fluid, postural effects, the ventricular pulsations, and the pulse pressure of the vascular choroid plexus.^{16–18}

If the intracranial pressure rises because of an increase of cerebrospinal fluid, the fast and the slow axoplasmic transport slow down and papilloedema can be observed in the ocular fundus.^{3,19} Since the cerebrospinal fluid is assumed to communicate between the different cerebrospinal fluid compartments a homogeneous pressure is postulated in all of these cerebrospinal fluid chambers.

However, this anatomical study of the subarachnoid space of the human optic nerve, as well as clinical observations such as asymmetric and unilateral papilloedema, raise legitimate questions as to whether or not the pressure in the subarachnoid space can indeed be expected to be equivalent to the pressure in the ventricles, the cisterns, and the cranial subarachnoid spaces. Because of the influence of the measuring process itself pressure measurements in small and non-homogeneous compartments are technically difficult to perform and the measurements are not necessarily reliable (personal experience) although other groups published their results with more confidence.¹³

Our morphological study of trabeculae and septa within the subarachnoid space of the optic nerve provides strong anatomical evidence that the hydrodynamics between and within the different cerebrospinal fluid segments, especially within the subarachnoid space of the optic nerve, is more complex than within the ideal Bernoulli tube. Cerebrospinal fluid dynamics should therefore not simply be regarded as an undeflected continuum in a series of interconnecting chambers. Because the subarachnoid space ends blindly in the bulbar segment behind the globe, cerebrospinal fluid needs to reverse its direction of flow in order to return to the site of resorption. There is histological evidence for a cerebrospinal fluid outflow system into the dura of the optic nerve itself,⁴ however this cerebrospinal outflow pathway albeit helpful in maintaining the pressure within certain limits during movements of the globe and optic nerve, may not be sufficient to absorb enough cerebrospinal fluid during a rise in fluid pressure. The subarachnoid space of the optic nerve is at the lower end of the cerebrospinal fluid chambers and because of its cul de sac anatomy the pressure gradient points

unidirectionally from the chiasmatic cistern to the subarachnoid space of the optic nerve in the manner of a hydraulic pump. Pressure could therefore build up in this small anatomical compartment that ends blindly behind the globe.

Morgan pointed out that the optic disc is located between two pressure compartments—the globe and the subarachnoid space.²⁰ The pressure in the subarachnoid space largely determines the retrolaminar pressure.²⁰ While the optic disc protrudes into the globe in papilloedema, it is excavated in glaucoma. Both diseases are associated with loss of ganglion cell axons. As axonal transport is influenced by pressure gradients, a steady translaminar pressure is expected to be critical in order to allow optimal axonal transport.²¹ The retrolaminar pressure depends mainly on the cerebrospinal fluid pressure in the bulbar segment. The architecture of the trabecular meshwork is believed to be important for pressure homeostasis. Since the compression and distortion of the lamina cribrosa may contribute to the development of glaucomatous optic neuropathy, likewise the anatomy of the subarachnoid space and the local cerebrospinal fluid pressure may influence both the lamina cribrosa and the retrobulbar part of the optic nerve.^{22–25}

Because of the small size of the trabeculae and septa even high resolution magnetic resonance imaging is not useful in demonstrating their anatomy in normal or in diseased states of the optic nerve.^{24–25} We suggest that further studies should aim to demonstrate possible changes of the arachnoid trabeculae and septa in pathological conditions such as asymmetric and unilateral papilloedema and optic neuritis.^{26–31}

ACKNOWLEDGEMENTS

The authors thank W Moor (Institute of Pathology, Kantonsspital Aarau, Switzerland) for careful preparation of all the optic nerves used in this study as well as P Rosenbaum, MD (Albert Einstein College of Medicine, New York, USA), and Th Resink MD (University of Basle, Switzerland) for assisting with English language problems.

We also thank Mrs M Erni for excellent technical help for SEM and TEM preparation. Special thanks to Duane E Haines, (Department of Anatomy, University of Mississippi Medical Center, Jackson, MI, USA) for many helpful and inspiring comments on the manuscript.

Part of this work was supported by the Rudolf and Fridl Buck Foundation.

Authors' affiliations

H E Killer, Department of Ophthalmology, Kantonsspital Aarau, Switzerland

H R Laeng, Department of Pathology, Kantonsspital Aarau, Switzerland

J Flammer, Department of Ophthalmology, School of Medicine, University of Basle, Basle, Switzerland

P Groscurth, Institute of Anatomy, University of Zürich, Switzerland

REFERENCES

- 1 **Anderson DR**. Ultrastructure of meningeal sheaths. Normal human and monkey optic nerves. *Arch Ophthalmol* 1969;**82**:659–74.
- 2 **Hayreh S**. The sheath of the optic nerve. *Ophthalmologica* 1984;**189**:54–63.

- 3 **Hayreh S**. Pathogenesis of oedema of the optic disc. *Doc Ophthalmol* 1966;**24**:291–411.
- 4 **Killer HE**, Laeng HR, Groscurth P. Lymphatic capillaries in the meninges of the human optic nerve. *J Neuro-Ophthalmol* 1999;**19**:222–8.
- 5 **Andres K**. Über die Feinstruktur der Arachnoidea und Dura mater von Mammalien. *Zeitschrift für Zellforschung* 1967;**79**:272–95.
- 6 **Rascol M**, Izard JY. Arachnoid and subarachnoid spaces of the vault of the skull in man. *Acta Neuropath (Berl)* 1978;**41**:41–4.
- 7 **Cloyd MW**, Low F. Scanning electron microscopy of the subarachnoid space in the dog. *J Comp Neurol* 1974;**15**:325–68.
- 8 **Zenker W**. Hüllen des Zentralnervensystems, Ventrikelauskleidung und Liquor cerebrospinalis. In: *Benninghoff Anatomie, Urban & Schwarzenberg*. Band 2, 1994:s339–60.
- 9 **Parkinson D**. Human spinal arachnoid septa, trabeculae, and "rough strands". *Am J Anat* 1991;**192**:498–509.
- 10 **Delmas JA**, Low F. Scanning electron microscopy of the subarachnoid space in the dog. *J Comp Neurol* 1975;**161**:515–40.
- 11 **Alcolado R**, Weller RO, Parrisch EP, et al. The cranial arachnoid and pia mater in man: Anatomical and ultrastructural observations. *Neuropathol Applied Neurobiol* 1988;**14**:1–17.
- 12 **Haines DE**, Harkey HL, Al-Mefty O. The "subdural" space: a new look at an outdated concept. *J Neurosurg* 1993;**32**:111–20.
- 13 **Liu D**, Kahn M. Measurement on relationship of subarachnoid pressure of the optic nerve to intracranial pressure in fresh cadavers. *Am J Ophthalmol* 1993;**116**:548–56.
- 14 **Braun JS**, Kaissling B, Le Hir M, et al. Cellular components of the immune barrier in the spinal meninges and dorsal ganglia of the normal rat: immunohistochemical (MHC class II) and electron-microscopic observations. *Cell Tissue Res* 1993;**273**:209–17.
- 15 **Persky B**, Low F. Scanning electron microscopy of the subarachnoid space in the dog: inflammatory response after injection of defibrinated chicken erythrocytes. *The Anatomical Record* 1985;**212**:307–18.
- 16 **Milhorat TH**, ed. *Hydrocephalus and the cerebrospinal fluid*. Baltimore: Williams and Wilkins, 1972.
- 17 **Bito LZ**, Davson H. Local variations in cerebrospinal fluid composition and its relationship to the composition of the extracellular fluid of the cortex. *Exp Neurol* 1966;**14**:264–80.
- 18 **Wood JH**, ed. *Neurobiology of cerebrospinal fluid*. New York and London: Plenum Press, 1982.
- 19 **Sanders MD**. The Bowman lecture. Papilledema: the pendulum of progress. *Eye* 1997;**11**:267–94.
- 20 **Morgan WH**, Dao-Yi Y, Cooper RL, et al. The influence of the cerebrospinal fluid pressure on the lamina cribrosa tissue pressure gradient. *Invest Ophthalmol Vis Sci* 1995;**36**:1163–72.
- 21 **Levy NS**. Functional implications of axoplasmic transport. *Invest Ophthalmol Vis Sci* 1974;**13**:639–40.
- 22 **Fechtner RD**, Weinreb RN. Mechanisms of optic nerve damage in primary open angle glaucoma. *Surv Ophthalmol* 1994;**39**:23–42.
- 23 **Quigley HA**, Hohman RM, Addicks EM. Morphologic changes in the lamina cribrosa correlated with neural loss in open-angle glaucoma. *Am J Ophthalmol* 1983;**673**–91.
- 24 **Gass A**, Barker, et al. MRI of the optic nerve in benign intracranial hypertension. *Neuroradiol* 1996;**38**:769–73.
- 25 **Jenkins JR**. Optic hydrops: isolated nerve sheath dilation demonstrated by CT. *Am J Neuroradiol* 1987;**8**:867–70.
- 26 **Strominger MB**, Weiss GB, Mehler MF. Asymptomatic unilateral papilledema in pseudotumor cerebri. *J Clin Neuro-Ophthalmol* 1992;**12**:238–41.
- 27 **Sedwick LA**, Burde RM. Unilateral asymmetric optic disk swelling with intracranial abnormalities. *Am J Ophthalmol* 1983;**96**:484–7.
- 28 **Killer HE**, Flammer J. Unilateral papilledema caused by a fronto-temporo-parietal arachnoid cyst. *Am J Ophthalmol* 2001;**132**:589–91.
- 29 **Marcelis J**, Silberstein SD. Idiopathic intracranial hypertension without papilledema. *Arch Neurol* 1991;**48**:392–9.
- 30 **Green JP**, Newman NJ, Stowe Z. "Normal pressure" pseudotumor cerebri. *J Clin Neuro-Ophthalmol* 1996;**16**:241–6.
- 31 **Killer HE**, Mironow A, Flammer J. Optic neuritis with marked distension of the optic nerve sheath due to local fluid congestion. *Br J Ophthalmol* (in press).



Architecture of arachnoid trabeculae, pillars, and septa in the subarachnoid space of the human optic nerve: anatomy and clinical considerations

H E Killer, H R Laeng, J Flammer, et al.

Br J Ophthalmol 2003 87: 777-781

doi: 10.1136/bjo.87.6.777

Updated information and services can be found at:

<http://bjo.bmj.com/content/87/6/777.full.html>

These include:

References

This article cites 24 articles, 5 of which can be accessed free at:

<http://bjo.bmj.com/content/87/6/777.full.html#ref-list-1>

Article cited in:

<http://bjo.bmj.com/content/87/6/777.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Neurology](#) (1042 articles)

[Optic nerve](#) (558 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>