

none was associated with CLL or CTCL. Some pigmentary changes were described in these cases, but in a pattern quite different from that seen in our patients.

To date of approximately 6000 patients treated with denileukin diftitox the manufacturer has only nine reports of visual disturbance including these patients (data on file at Ligand Pharmaceuticals). Formal review of these reports has not revealed a clear pattern of ocular involvement. However, the similarity of the two cases reported here is striking, and raises the possibility of a novel toxic retinopathy.

Acknowledgements

We gratefully acknowledge the assistance of Ligand Pharmaceuticals in providing information from their database and reviewing the manuscript.

J B Ruddle, C A Harper

The Royal Victorian Eye and Ear Hospital, East Melbourne, Australia

D Hönemann, J F Seymour, H M Prince

Haematology Service, Peter MacCallum Cancer Centre, East Melbourne, and University of Melbourne, Australia

C A Harper

Eye Research Australia, East Melbourne, and Department of Ophthalmology, University of Melbourne, Australia

Correspondence to: Dr Jonathan B Ruddle, Royal Victorian Eye and Ear Hospital, 32 Gisborne Street, East Melbourne, Victoria 3002, Australia; jonathan.ruddle@eyeandear.org.au

doi: 10.1136/bjo.2006.091165

Accepted for publication 1 April 2006

References

- 1 **Morgan SJ, Seymour JF, Prince HM, et al.** Confirmation of the activity of the interleukin-2 fusion toxin denileukin diftitox against chemorefractory chronic lymphocytic leukemia, including cases with chromosome 17p deletions and without detectable CD25 expression. *Clin Cancer Res* 2004;**10**:3572-5.
- 2 **Olsen E, Duvic M, Frankel A, et al.** Pivotal phase III trial of two dose levels of denileukin diftitox for the treatment of cutaneous T-cell lymphoma. *J Clin Oncol* 2001;**19**:376-88.
- 3 **Thornton AM.** Signal transduction in CD4+CD25+ regulatory T cells: CD25 and IL-2. *Front Biosci* 2006;**11**:921-7.
- 4 **Takeuchi M, Keino H, Kezuka T, et al.** Immune responses to retinal self-antigens in CD25(+)CD4(+) regulatory T-cell-depleted mice. *Invest Ophthalmol Vis Sci* 2004;**45**:1879-86.
- 5 **Weleber RG, Watzke RC, Shults WT, et al.** Clinical and electrophysiologic characterization of paraneoplastic and autoimmune retinopathies associated with anti-nucleolar antibodies. *Am J Ophthalmol* 2005;**139**:780-94.
- 6 **To KW, Thirkill CE, Jakobiec FA, et al.** Lymphoma-associated retinopathy. *Ophthalmology* 2002;**109**:2149-53.
- 7 **Adamus G, Ren G, Weleber RG.** Autoantibodies against retinal proteins in paraneoplastic and autoimmune retinopathy. *BMC Ophthalmol* 2004;**4**:5.
- 8 **Sen HN, Chan CC, Caruso RC, et al.** Waldenström's macroglobulinemia-associated retinopathy. *Ophthalmology* 2004;**111**:535-9.

NOTICES

Macula of Paris

This meeting will take place on 6th September 2006 at the InterContinental Paris Le

Grand Hôtel, 2 Rue Scribe, 75009 Paris, France. For further information, please contact Gisèle Soubrane, MD, Department of Ophthalmology, University Créteil-Paris XII, Avenue de Verdun, 40, 94010 Creteil, France. Or email Anne-Sophie Caron at: as.caron@colloquium.fr

Back of the eye

The latest issue of *Community Eye Health* (No 57) assesses treatments for age related macular degeneration and other back of the eye conditions. For further information please contact: Journal of Community Eye Health, International Resource Centre, International Centre for Eye Health, Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK (tel: +44 (0)20 7612 7964; email: Anita.Shah@lshtm.ac.uk; online edition: www.jceh.co.uk). Annual subscription (4 issues) UK £28/US\$45. Free to developing country applicants

Managing human resources

The latest issue of *Community Eye Health* (No 56) assess the use of human resources in the delivery of eye care. For further information please contact: Journal of Community Eye Health, International Resource Centre, International Centre for Eye Health, Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK (tel: +44 (0)20 7612 7964; email Anita.Shah@lshtm.ac.uk, url: ; online edition: www.jceh.co.uk). Annual subscription (4 issues) UK £28/US\$45. Free to developing country applicants.

8th EUNOS Meeting – 2007

The 2007 European Neuro-ophthalmology Society meeting (EUNOS; www.eunos.org) will be taking place in Istanbul, Turkey on 26-29th May 2007. For further information please visit www.eunos2007.org or contact Pinar Aydin aydinp@eunos2007.org

Teaching courses on Retinal and Vitreous Surgery

Several teaching courses on Retinal and Vitreous Surgery have been organised throughout 2006 and 2007 around the world in association with the International Faculty. For further information on each of these courses please contact Ingrid Kressig, Univ.-Augenklinik Theodor-Kutzer-Ufer 1-3, 68164 Mannheim, Germany; email: Ingrid.kreissig@augen.ma.uni-heidelberg.de; website: <http://kressig.uni-hd.de/>.

EVER 2006

The EVER 2006 meeting will take place in Vilamoura, Portugal on 4-7th October 2006. For further information please contact the EVER Office, Kapucijnenvoer 33, 3000 Leuven, Belgium; website www.ever.be

Recruitment halted on ESCRS study on antibiotic prophylaxis of endophthalmitis following clear beneficial result

THE ESCRS has terminated recruitment for their two year study of antibiotic prophylaxis of endophthalmitis following cataract surgery. Quarterly analysis of the figures to date

by the study's statisticians at the University of Strathclyde clearly indicates a beneficial treatment effect. In January 2006 the Data Monitoring Committee recommended that the study be unmasked and found the result to be so clear that they recommended to the Study Chairman that recruitment be halted.

The study has found that the risk of contracting endophthalmitis following phacoemulsification cataract surgery is significantly reduced by an intracameral injection of cefuroxime at the end of surgery.

The ESCRS Study, a partially masked, randomized, placebo controlled, multi-national study conducted at 24 ophthalmology centres across Europe commenced recruitment in September 2003. A preliminary report on the primary results will be published in the March 2006 issue of the Journal of Cataract and Refractive Surgery. Complete follow-up data and analyses will be reported at the XXXIV Congress of the ESCRS in London in September 2006 and will subsequently be published in the Journal of Cataract and Refractive Surgery.

For further information contact Caroline Fitzpatrick European Society of Cataract and Refractive Surgeons, tel: +353 1 209 1100, caroline.fitzpatrick@escrs.org.

Prevention of Blindness Fellowship Programme

Application are invited for BCPB Fellowships to start in 2007. The aims of the Fellowships are to fund research and training in prevention of blindness for high caliber clinicians and scientists from the UK and overseas. Projects must further the goals of VISION 2020: THE RIGHT TO SIGHT, the elimination of avoidable blindness. In 2007, BCPB seeks to fund one Fellow from the UK and one Fellow from a low-income country to undertake projects that focus on Africa.

Priority will be given to applicants who:

- Demonstrate that their project is innovative and increases knowledge of the causes of blindness and/or its prevention in line with the priorities of VISION 2020
- Demonstrate the ability and ambition to pass on their skills in blindness prevention

The fellowships will be worth up to £60,000pa for 2 or 3 years. Applications must be submitted jointly by the Fellowship candidate and the supervisor at the host institution in the UK.

For full information and an application form, see www.bcpb.org or contact Jackie Webber at BCPB, 59-60 Russell Square, London WC1B 4HP or by email: info@bcpb.org. Closing date for applications is 30 September 2006.

CORRECTION

doi: 10.1136/bjo.2005.074955corr1

In the Scientific report titled, Incidence and severity of keratoconus in Asir province, Saudia Arabia (*Br J Ophthalmol* 2005;**89**: 1403-6) figure 2 was incorrect. The original legend printed for figure 2 has also changed. The authors apologise for this error. A full corrected figure is available on the BJO website at <http://www.bjophthalmol.com/supplemental>.



CORRECTION

Br J Ophthalmol 2006 90: 1071
doi: 10.1136/bjo.2005.074955corr1

Updated information and services can be found at:
<http://bjo.bmj.com/content/90/8/1071.full.html>

Email alerting service

These include:

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>