

WHENEVER
THE BENEFITS
OF MORPHINE
ARE NEEDED—

'CYCLIMORPH'

(Morphine plus Cyclizine)^{brand} INJECTION

'Cyclimorph' provides all the benefits of morphine but overcomes the distressing, even, dangerous vomiting often caused by this analgesic.

'Cyclimorph' is indicated whenever morphine is indicated—especially where nausea and vomiting would be harmful as in

myocardial infarction

pre-operative medication

post-operative pain

debilitated patients

Cyclizine, the anti-nauseant and anti-emetic in 'Cyclimorph', is one of the most successful agents for the control of morphine-induced vomiting. It acts rapidly, has a similar duration of action to morphine, and is remarkably safe.

'Cyclimorph' is subject to the D.D. regulations. It is issued in 1 ml. ampoules (boxes of 5 and 25), in two strengths, viz.:

'CYCLIMORPH' 10

Morphine tartrate	10 mg. (gr. $\frac{1}{8}$)
Cyclizine tartrate	50 mg.
	per ml.


'CYCLIMORPH' 15

Morphine tartrate	15 mg. (gr. $\frac{1}{4}$)
Cyclizine tartrate	50 mg.
	per ml.




BURROUGHS WELLCOME & CO. (The Wellcome Foundation Ltd.) LONDON

a new treatment of total hyphaema



Urokinase Leo



plasminogen activator of human origin

Recent research in the field of fibrinolytic therapy has resulted in the development of a valuable new technique for the treatment of traumatic total hyphaema. Lysis of the clot has been obtained in a matter of minutes with return of vision by irrigation of the anterior chamber with Urokinase.¹ In the initial study more than twenty cases of secondary haemorrhage in the anterior chamber have been

treated with Urokinase with very satisfactory results.² Subsequent use of Urokinase since the original communications have confirmed the practicability and success of Urokinase therapy in this condition.

Urokinase—Available in ampoules containing 5,000 Ploug units in a sterile lyophilized preparation.

1 Lancet (1963) ii, 1143

2 Trans. Ophthal. Soc. U.K. In press

Leo Laboratories Limited
Copenhagen London Dublin

