



OPEN ACCESS

Should tumbling E go out of date in amblyopia screening? Evidence from a population-based sample normative in children aged 3–4 years

Sandra Guimaraes,^{1,2,3} Tiago Fernandes,¹ Patrício Costa,^{2,3,4} Eduardo Silva^{5,6,7}¹Department of Ophthalmology, Hospital de Braga, Braga, Portugal²School of Medicine, Life and Health Sciences Research

Institute (ICVS), University of Minho, Braga, Portugal

³ICVS/3B's—PT Government Associate Laboratory, Braga, Braga, Portugal⁴Clinical Academic Center, Braga, Portugal⁵Centro Cirúrgico de Coimbra, Coimbra, Portugal⁶IBILI, Faculty of Medicine, University of Coimbra, Coimbra, Coimbra, Portugal⁷Department of Ophthalmology, CHLN, Lisbon, Portugal**Correspondence to**

Dr Sandra Guimaraes, Department of Ophthalmology, Hospital de Braga, Sete Fontes—São Victor, 4710-243 Braga, Portugal; sandraguimaraes. eye@gmail.com

Received 3 May 2017

Revised 4 August 2017

Accepted 24 August 2017

ABSTRACT**Aims** To determine a normative of tumbling E optotype and its feasibility for visual acuity (VA) assessment in children aged 3–4 years.**Methods** A cross-sectional study of 1756 children who were invited to participate in a comprehensive non-invasive eye exam. Uncorrected monocular VA with crowded tumbling E with a comprehensive ophthalmological examination were assessed. Testability rates of the whole population and VA of the healthy children for different age subgroups, gender, school type and the order of testing in which the ophthalmological examination was performed were evaluated.**Results** The overall testability rate was 95% (92% and 98% for children aged 3 and 4 years, respectively). The mean VA of the first-day assessment (first-VA) and best-VA over 2 days' assessments was 0.14 logMAR (95% CI 0.14 to 0.15) (decimal=0.72, 95% CI 0.71 to 0.73) and 0.13 logMAR (95% CI 0.13 to 0.14) (decimal=0.74, 95% CI 0.73 to 0.74). Analysis with age showed differences between groups in first-VA ($F(3,1146)=10.0$; $p<0.001$; $\eta^2=0.026$) and best-VA ($F(3,1155)=8.8$; $p<0.001$; $\eta^2=0.022$). Our normative was very highly correlated with previous reported HOTV-Amblyopia-Treatment-Study (HOTV-ATS) (first-VA, $r=0.97$; best-VA, $r=0.99$), with 0.8 to 0.7 lines consistent overestimation for HOTV-ATS as described in literature. Overall false-positive referral was 1.3%, being specially low regarding anisometropias of ≥ 2 logMAR lines (0.17%). Interocular difference ≥ 1 line VA logMAR was not associated with age ($p=0.195$).**Conclusions** This is the first normative for European Caucasian children with single crowded tumbling E in healthy eyes and the largest study comparing 3 and 4 years old testability. Testability rates are higher than found in literature with other optotypes, especially in children aged 3 years, where we found 5%–11% better testability rates.**INTRODUCTION**Although WHO and the National Academy of Sciences Committee on Vision (NASCV) have issued recommendations about the design of optotypes used in visual acuity (VA) charts that were included in international guidelines,¹ several charts used with children meet these criteria: LEA symbols, Sloan letters/numbers, tumbling E and HOTV. Although HOTV test is commonly used in the USA,^{2–4} no standard VA testing in children has been accepted worldwide.^{5,6} Directional optotypes as Landolt C and tumbling E charts are widelyused in non-spoken English countries.^{5,7} They are considered better optotypes as grating recognition and orientation is superior to letter recognition⁸ and the progression of resolution angles is similar throughout the test.⁹ Tumbling E has the advantage of performing better than Landolt C for astigmatism against-rule.¹⁰ Test-retest tumbling E studies demonstrated highly repeatable results ($r=0.97$).¹¹ Nevertheless, tumbling E has been reported as conceptually difficult for young children¹ as it requires spatial orientation skills, and recently, it has been non-recommended in young children as 'they may not yet have developed the ability to express the orientation of these optotypes'.⁴ Doubts whether up-down-right-left orientation is already mastered by children aged 3 years old (yo), or not, is doubtful.Although single optotypes have better cooperation in children if crowding is not used,^{4,12} crowded bars are more accurate^{4,13} to overcome the 'crowding phenomenon' present in amblyopia and should be used.¹⁴ In amblyopic children, the kind of crowded optotype also revealed differences depending on the optotype.¹⁵ Normative with single crowded HOTV^{2,3,16} and LEA¹² have been published. Regarding tumbling E, normative are either in charts,^{5,7,17} lines¹⁸ or single E without crowded bars,¹⁸ but to the best of our knowledge, no normative with single-crowded tumbling E is available in the optimal age screening,¹⁹ making it crucial to set a normative.Normal VA in healthy eyes should be strictly assessed when setting a normative with accurate referral criteria for vision screening. It must be ensured that all ophthalmological pathologies are excluded. Some recent normative had these important considerations into account,^{3,12,16,17} but none with single-crowded tumbling E.

The scope of this study is to determine a normative for 3–4 yo in healthy eyes with tumbling E in the single-crowded form and to determine if it is a feasible VA tool in this age group. We search for differences between 3 and 4 yo in testability, false-positive referrals and we compare it with other previous normative published. A normative regarding age will be presented.

MATERIALS AND METHODS**Participants**

Between May 2014 and June 2016, all children registered in Portuguese National Health Service with 3–4 yo, attending public and private schools in



CrossMark

To cite: Guimaraes S, Fernandes T, Costa P, et al. *Br J Ophthalmol* Published Online First: [please include Day Month Year]. doi:10.1136/bjophthalmol-2017-310691

Braga, were invited to participate in an eye exam at the Ophthalmology Department of the *Hospital de Braga*. This study was approved by the Human Research Ethics Committee of the *Hospital de Braga* and adhered to the tenets of Declaration of Helsinki. Written informed consent was obtained from the parent/guardian of child before examinations.

Procedures, ophthalmological examination and VA assessment

All children underwent a comprehensive non-invasive eye exam performed by orthoptists and paediatric ophthalmologists trained in the study's protocol. Age, sex, school source and order of testing were registered. Examination included uncorrected monocular VA with single-crowded tumbling E (VA assessment was always done by experienced orthoptists in children examination), ocular motility, cover test, Hirshberg, three Plusoptix S04 measures, Randot stereoacuity with circles, biomicroscopy and funduscopy. All astigmatisms were registered in negative cylinders for comparison proposes.

VA was recorded as the smallest optotype size which the child identified in all four directions, in order not to misdiagnose meridional amblyopia. VA scores were provided in 0.1 decimal increments, recorded in a decimal scale and converted to logMAR for analysis. After a brief training with binocular VA, testing was conducted monocularly, always starting with the right eye (OD). The eye was occluded with a non-adhesive paper occluder, explaining the child it was a 'pirate game'. The child could either use fingers or a plastic E to indicate the direction of the letter. Half of the children started the exam with VA measurement and then went to perform other exams (VA was assessed in the end in the remainder). These VA results were considered the first visual acuity assessment (first-VA) as some children were called for a second evaluation as explained ahead. For testability rates, a child was considered testable if it was possible to determine monocular first-VA in both eyes. Stereoacuity measurements were done with Randot test. Emmetropia was defined as: normal spherical equivalent (SE) based on 98.6% negative predictive value (NPV) for hyperopia adapted criteria (SE <2.0 dioptres) and 98% NPV for astigmatism (<1.5 dioptres) for Plusoptix.²⁰ Myopia and anisometropia criteria were SE <-0.5 dioptres and sphere or cylinder <1.0 difference, respectively. We applied these criteria for all three consecutive Plusoptix measurements.

If emmetropic children with no associated ophthalmological pathology had poor collaboration or a monocular first-VA without correction ≤ 0.6 or interocular difference ≥ 3 lines in decimal scale, children's VA was reassessed on a second day, this time in their parent's presence. In the second VA assessment, the same protocol was applied. We considered the best VA (best-VA) over the first and second evaluation only for comparison purposes with other recent normative published.¹⁶ Nevertheless, we used first-VA as our main normative outcome variable because it best reflects screening practice and, besides, using the best of multiple measures of VA could introduce a systematic bias, causing false conclusions.²¹

Exclusion/inclusion criteria

Only children aged between 36 and 59 months were included. After excluding 12 children that came twice to the exam, the whole population of 1735 was considered for testability assessment. For VA normative, only healthy eyes were considered and exclusion criteria are presented in [figure 1](#).

STATISTICAL ANALYSIS

Data were gathered in Microsoft Office Excel and exported to IBM Statistical Package for the Social Sciences Statistics (SPSS V.24) for statistical analysis. VA was measured in a decimal scale and converted into logMAR units for statistical analyses as recommended.²² For being more comprehensive, VA means, differences of means, SD and CIs were also expressed in decimal values right after all logMAR results (into parenthesis). The statistical procedures assumptions, such as normality, homogeneity of variances, were considered. When appropriate, Student's paired-sample t-test, independent-sample t-test, one-way analysis of variance or Welch test were used whether there was homogeneity of variances or not in Levene's test, and the corresponding post hoc testing was done, either with Bonferroni or Games-Howell, if there was statistical significance. When Levene's test indicated unequal variances, df was adjusted correspondingly. Effect size measures along with statistical significance ($p < 0.05$) were presented.

RESULTS

Participation rate of the invited children was 96%. Regarding testability, we found 89/1735 (5.1%) non-cooperative children in VA responses in at least one eye. Testability in age subgroups is presented in [table 1](#).

Testability in different age subgroups of 3.0 (36–41 months), 3.5 (42–47 months), 4.0 (48–53 months) and 4.5 (54–59 months) yo. On the right, comparison between 3 and 4 yo.

Of the 1182 children that met the criteria for sample VA normative ([figure 1](#)), median age was 47.37 ± 4.74 months, 51.0% were males, 54% attended public schools and in 43% ophthalmological examination started with VA assessment. The mean first-VA was 0.14 logMAR (95% CI 0.14 to 0.15) (0.72 (95% CI 0.71 to 0.73)) for OD and 0.14 logMAR (95% CI 0.14 to 0.15) (0.72, (95% CI 0.71 to 0.73) for OS. Normality of these measures were proven by acceptable values of kurtosis and skewness (OD: 0.75; 0.44/OS: 0.94; 0.49) and histogram distribution. As first-VA for the OD and OS were not significantly different ($t(1149) = -0.22$, $p = 0.826$, $d = 0.00646$, paired sample t-test), ($t(1150) = 0.67$, $p < 0.001$, Pearson's correlation), we used first-VA OD as the normative of monocular VA in the first observation. Differences ≥ 1 line VA logMAR was found in 116 children (9.8%).

From the 157 with at least one criteria for a second evaluation, 94 were re-examined on a second day, for VA re-assessment (participation rate 60%). Differences between the best-VA OD and OS were not significant ($t(1158) = 1.14$, $p = 0.254$, $d = 0.0335$) ($t(1159) = 0.62$, $p < 0.001$). From now on, we will no longer report OD or OS, as we will always report OD results, either regarding first-VA or best-VA, unless specified differently.

First-VA and best-VA had no differences regarding gender and the order in which the ophthalmological examination was performed, respectively, ($t(1148) = 0.428$, $p = 0.668$, $d = 0.026$) and ($t(558) = 0.126$, $p = 0.90$, $d = 0.010$) for first-VA, and ($t(1157) = 0.916$, $p = 0.36$, $d = 0.054$) and ($t(565) = 1.13$, $p = 0.261$, $d = 0.097$) for best-VA. Regarding school type, there was a difference of 0.014 logMAR (lower logMAR for public schools) found in first-VA ($t(1030.7) = -2.85$, $p = 0.005$, $d = 0.17$).

Regarding first-VA, age had a very small correlation ($r(1150) = -0.17$, $p < 0.001$) and differences found between 3 and 4 yo were 0.022 logMAR (95% CI 0.012 to 0.031) ($t(1135.2) = 4.6$, $p < 0.001$, $d = 0.27$, independent sample t-test). With best-VA, similar differences of 0.018 logMAR were




Figure 1 Inclusion/exclusion criteria for testability rates and for visual acuity (VA) normative. The interocular difference was measured in decimal lines. UCVA, uncorrected visual acuity.

observed (95% CI 0.001 to 0.027) ($t(1150.4)=4.2$, $p<0.001$, $d=0.25$, independent sample t-test). For comparison with recent normative studies, we present normative of first-VA and best-VA in four age subgroups in [table 2](#).

Analysis showed differences between groups in first-VA ($F(3,1146)=10.0$; $p<0.001$; $\eta^2=0.026$) and best-VA ($F(3,1155)=8.8$; $p<0.001$; $\eta^2=0.022$). Post hoc analysis using Bonferroni test showed no difference in first-VA between 3.0 and 3.5 yo ($p=0.83$). All other subgroups had statistically significant differences of 0.01556 logMAR (95% CI 0.0018 to 0.0293), $p=0.019$ between 3.5 and 4.0 yo, 0.02274 logMAR (95% CI 0.0011 to 0.0444), $p=0.036$ between 3.0 and 4.0 yo, 0.03906 logMAR (95% CI 0.0193 to 0.0588), $p<0.001$ between 3.5 and 4.5 yo, 0.04624 logMAR (95% CI 0.0204 to 0.0721), $p<0.001$ between 3.0 and 4.5 yo, 0.02350 logMAR (95% CI 0.0042 to 0.0428), $p=0.010$ between 4.0 and 4.5 yo.

Regarding false-positive referrals, for children aged between 36 and 47 months, with the recommended cut-off value of 0.4 logMAR (0.4), we had 1% of false-positive referral. For children

aged between 48 and 59 months, for the recommended cut-off value of 0.3 logMAR (0.5), we had 1.5% of false-positive referral. For false-positive referrals regarding anisometropias, we used a difference ≥ 2 logMAR lines between OD and OS as the cut-off value. Of the 1182 children, only two had anisometropia. We used the second evaluation result as the true diagnosis and none of those two had in fact anisometropia, giving a false-positive referral of 0.17%.

DISCUSSION

This is the first population-based normative for European Caucasian children with single-crowded tumbling E. It is also the largest study comparing 3 and 4 yo testability, such an important age range for screening purposes.

Our exclusion criteria are similar to others^{3 16 17} that were also judicious about measuring VA in only healthy eyes. Based on refractive criteria, we excluded 20.4% children, in line with Leone *et al*¹⁶ that excluded 29.9% but in a wider age range (6–72 months), where more refractive errors are expected to be found.

Although we have higher global testability rates (94.9%), our main advantage is in the 3 years' group, where our testability is considerably higher ([table 3](#)).

Although our global testability was lower than the 97.6% reported with HOTV-ATS,²³ it is important to note that higher testability reported by Kupl *et al* could have been achieved by older children, as 28% of children in that study were 5 yo, and children between 3.0 and 3.5 yo were not included. If we compare similar age groups, we find 94.3% and 98.3% testability, better than 93.3% and 96.7% at 3.5 and 4 yo respectively, found

Table 1 Testability						
Age	n		Non-cooperative		Testability (%)	
3.0	3 yo	233	929	36	75	84.5 91.9
3.5		696		39		94.3
4.0	4 yo	638	806	14	14	97.8 98.3
4.5		168		0		100.0
Total			1735		89	94.9

yo, year old.

Table 2 Normative for the first-VA and the best-VA

Age yo (months)	First-VA			Best-VA		
	n	LogMAR (95% CI)	Decimal (95% CI)	n	LogMAR (95% CI)	Decimal (95% CI)
3 yo (36–41)	118	0.15 (0.14 to 0.17)	0.70 (0.67 to 0.72)	120	0.15 (0.14 to 0.17)	0.70 (0.68 to 0.73)
3.5 yo (42–47)	453	0.15 (0.14 to 0.16)	0.71 (0.69 to 0.72)	459	0.14 (0.13 to 0.15)	0.73 (0.71 to 0.74)
4 yo (48–53)	464	0.14 (0.13 to 0.14)	0.73 (0.72 to 0.74)	465	0.13 (0.12 to 0.13)	0.74 (0.73 to 0.76)
4.5 yo (54–59)	115	0.11 (0.10 to 0.12)	0.77 (0.75 to 0.80)	115	0.11 (0.10 to 0.12)	0.78 (0.75 to 0.80)

VA of the first-VA and of the best-VA over two assessments; yo, years old; VA, visual acuity.

with HOTV-ATS.²³ Comparing our testability with HOTV-ATS in four age subgroups described in 2012,²⁴ our rates are 4.5%, 1.3%, 2.8% and 2% higher than HOTV-ATS for 3.0, 3.5, 4.0 and 4.5 yo, respectively, also showing the best benefit for 3.0 yo.

Regarding testability rates, one strong point in our study is that it reflects testability of all children before exclusion criteria, meaning that testability among children with refractive errors and/or amblyopia is also considered. If we had considered only normal eyes in our study, the overall testability rate would have been 97.3%.

We speculate that this good testability scores could be achieved because today's children stimulation with Smartphones and Tablets at very young ages leads to up-down-left-right recognition ability developed sooner and the exam could be experienced as another game. Besides, we realised that children tend to cooperate much better if they are 'in-group' with schoolteachers and colleges than when they come alone with their parents. It happens as if the child understands better his/her role of commitment to the performance of the tests, seeing it as a school extension. As a limitation, our better testability could be due to the population studied, as Braga is a modern urban city.

We used first-VA for the normative as it best reflects clinical practice. Regarding VA variation with age, there is very small

intensity of the negative correlation found. Moreover, either in first-VA or best-VA age subgroups variation analyses, the biggest difference found was between 3.0 and 4.5 yo, but differences found between all age subgroups were <0.5 logMAR lines, what, in our opinion is not clinically relevant, as the interval measurements used in logMAR VA scales are 0.1 logMAR units (1 line), making differences of <0.05 logMAR units (<0.5 lines) below the threshold scale measurement, so, not possible to detect in clinical practice and, for the same reason, would not change VA thresholds normative guidelines. Effect size of these differences analyses revealed that age only explained 2.6% and 2.2% of first-VA and best-VA, respectively. Regarding interocular differences of ≥ 1 line VA logMAR, we had only 9.8%, <26% report¹⁶ and it was not associated with age ($p=0.195$) as reported earlier.¹⁶

Comparing our best-VA normative (table 4) with the recently reported HOTV-ATS,¹⁶ we found HOTV-ATS has 0.079 and 0.072 logMAR consistently better VA than our first-VA and best-VA with very high correlations ($r=0.97$ and $r=0.99$), respectively. It is important to note that this HOTV-ATS normative is based on the best-VA of two consecutive measures in two different appointments, what is the equivalent as our best-VA. The same tendency is described in literature regarding HOTV

Table 3 Testability rates with different optotypes

Study	Optotype	n		Testability	
		3 yo	total	3 yo	total
Present study	Single-crowded tumbling E	929	1735	91.90%	94.9%
		806		98.30%	
Leone <i>et al</i> ²⁴	HOTV-ATS	363	723	86.5%	91.4%
		360		96.3%	
Pan <i>et al</i> ³	HOTV-ATS	460	1027	81%	90.5%
		567		98%	
Kvarnstrom <i>et al</i> ²⁷	HOTV chart	478	707	84.80%	87.4%
		229		92.80%	
Kvarnstrom <i>et al</i> ²⁷	LEA chart	478	707	82.80%	86.1%
		229		92.90%	
Kupl <i>et al</i> ²³	HOTV-ATS	225	858	(3–3.5 yo not included)	97.6% (5yo included)
		633		96.70%	
Becker <i>et al</i> ¹² Paediatric routine examination	Single LEA symbols	71	134	76.00%	85.00%
		63		95.00%	
Becker <i>et al</i> ¹² Ophthalmic reevaluation	Single LEA symbols	10	28	80.00%	92.80%
		18		100.00%	

For a more accurate comparison, testability rates are presented separately at 3 and 4 yo. yo, years old.

Table 4 Normative of tumbling E and HOTV

Study	Optotype	Mean logMAR best-VA (95% CI) decimal			
		3.0 yo	3.5 yo	4.0 yo	4.5 yo
Present study	Single-crowded tumbling [E]	0.15 (0.14 to 0.17) (0.70 (0.68 to 0.73))	0.14 (0.13 to 0.15) (0.73 (0.71 to 0.74))	0.13 (0.12 to 0.13) (0.74 (0.73 to 0.76))	0.11 (0.10 to 0.12) (0.78 (0.75 to 0.80))
Leone <i>et al</i> ¹⁶	HOTV-ATS	0.09 (0.07 to 0.10) (0.81 (0.79 to 0.85))	0.07 (0.05 to 0.09) (0.85 (0.81 to 0.89))	0.05 (0.03 to 0.06) (0.89 (0.87 to 0.93))	0.03 (0.02 to 0.04) (0.93 (0.91 to 0.95))

VA, visual acuity.

charts and ETDRS charts, where HOTV-ATS has approximately 1 line better VA results.²⁴

Comparing our best-VA normative with HOTV in different age subgroups of 3.0 (36–41 months), 3.5 (42–47 months), 4.0 (48–53 months) and 4.5 (54–59 months) yo. Note that single crowded optotypes and best-VA over two observations are used in both studies.

Despite the limitation of this assumption, if we consider school type as a proxy of socioeconomic status, the difference of 0.014 logMAR found in first-VA, although statistically significant, had a small effect size, and to our interpretation it should not be considered relevant.²⁵

Regarding false-positive referrals, for 3 yo we had 1% of false-positive referral, which is similar to the 0.9% found by Leone *et al*,¹⁶ using the same cut-offs, with HOTV-ATS protocol²⁶ on the electronic VA tester and better than the 5.3% they found with ETDRS or HOTV chart.¹⁶ For children 4 yo, we had 1.5% false-positive referrals, which is 1.1% higher than the 0.4% found by Leone *et al* with HOTV-ATS²⁶ and much lower than the 6% found with ETDRS or HOTV chart.¹⁶ Regarding false-positive referrals for anisometropias, we had a very low false-positive referral of 0.17%, much lower than 6% described by Pan *et al* with HOTV-ATS.³ Their normative, nevertheless represents a wider age range (30–72 months).

As limitations, we only studied Braga's population. We did not use race as a variable because being our population mainly Caucasian, we assume it reflects no influence in our results. In a 235 sample, we only reported two non-Caucasians (0.85%). The time of the exam was not assessed. The published differences of 1.12 more minutes to test a 3 yo child vs 4 yo child with LEA and HOTV charts,²⁷ arises the question as if these differences are relevant, considering the time spent for receiving the child, register the results and giving the report of the screening. Further studies are needed regarding this issue, including time to perform the exams in healthy versus unhealthy eyes.

In summary, tumbling E has the highest testability rates between 3 and 4 yo when compared with literature, specially in children 3 yo where it is considerably better. Past evidences fearing the differences in collaboration between 3 and 4 yo are no longer supported. In screenings where VA is the gold standard, tumbling E is a good instrument for VA assessment in children aged 3 and 4 years.

Acknowledgements This study has not been presented previously and had no specific support.

Contributors Research design, data analysis and interpretation: SG, PC and ES. Data acquisition and research execution: SG and TF. All authors participated in manuscript preparation.

Competing interests None declared.

Patient consent All children aged between 3 and 4 years from public and private schools in Braga, Portugal, were invited to participate in an eye exam at the Ophthalmology Department of the Hospital de Braga. All children were legal residents, registered in Portuguese National Health Service. Written informed consent was obtained from the parent or guardian of each child before examinations. The consent was not a specific consent from BMJ. The written consent was approved by

the Human Research Ethics Committee of the Hospital de Braga and adhered to the tenets of the Declaration of Helsinki.

Ethics approval Human Research Ethics Committee of the Hospital de Braga.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Panel AAoOPOS. *Preferred Practice Pattern Guidelines. Secondary Preferred Practice Pattern Guidelines*. Amblyopia, 2012.
- Drover JR, Felius J, Cheng CS, *et al*. Normative pediatric visual acuity using single surrounded HOTV optotypes on the Electronic Visual Acuity Tester following the Amblyopia Treatment Study protocol. *J Aapos* 2008;12:145–9.
- Pan Y, Tarczy-Hornoch K, Cotter SA, *et al*. Visual acuity norms in pre-school children: the Multi-Ethnic pediatric eye disease study. *Optom Vis Sci* 2009;86:607–12.
- Donahue SP, Baker CN. Committee on Practice and Ambulatory Medicine, American Academy of Pediatrics Section on Ophthalmology, American Academy of Pediatrics American Association of Certified Orthoptists American Association for Pediatric Ophthalmology and Strabismus American Academy of Ophthalmology. Procedures for the Evaluation of the Visual System by Pediatricians. *Pediatrics* 2016;137:e20153597.
- Lai YH, Wang HZ, Hsu HT. Development of visual acuity in preschool children as measured with Landolt C and Tumbling E charts. *J Aapos* 2011;15:251–5.
- Bois C, Binot MC, Jonqua F, *et al*. [Visual screening in 3- to 5-year-old preschool children: Maternal and Infant Welfare evaluation program in the Hauts-de-Seine district, France]. *J Fr Ophthalmol* 2007;30:570–6.
- Moganewari D, Thomas J, Srinivasan K, *et al*. Test Re-Test reliability and validity of different visual acuity and stereoacuity charts used in preschool children. *J Clin Diagn Res* 2015;9:NC01–5.
- NIST NfSaT. Standards for Visual Acuity doc 12930. 2006 https://www.nist.gov/sites/default/files/documents/el/isd/ks/Visual_Acuity_Standards.
- Messias A, Jorge R, Cruz AA. [Logarithmic visual acuity charts: reasons to use and how to design it]. *Arq Bras Oftalmol* 2010;73:96–100.
- Reich LN, Ekabutr M. The effects of optical defocus on the legibility of the Tumbling-E and Landolt-C. *Optom Vis Sci* 2002;79:389–93.
- Lovie-Kitchin JE. Validity and reliability of visual acuity measurements. *Ophthalmic Physiol Opt* 1988;8:363–70.
- Becker R, Hübsch S, Gräf MH, *et al*. Examination of young children with Lea symbols. *Br J Ophthalmol* 2002;86:513–6.
- Schlenker MB, Christakis TJ, Braga-Mele RM. Comparing a traditional single optotype visual acuity test with a computer-based visual acuity test for childhood amblyopia vision screening: a pilot study. *Can J Ophthalmol* 2010;45:368–74.
- Simons K. Visual acuity norms in young children. *Surv Ophthalmol* 1983;28:84–92.
- O'Boyle C, Chen SI, Little JA. Crowded letter and crowded picture logMAR acuity in children with amblyopia: a quantitative comparison. *Br J Ophthalmol* 2017;101:457–61.
- Leone JF, Mitchell P, Kifley A, *et al*. Normative visual acuity in infants and preschool-aged children in Sydney. *Acta Ophthalmol* 2014;92:e521–e529.
- Guo X, Fu M, Lu J, *et al*. Normative distribution of visual acuity in 3- to 6-year-old Chinese preschoolers: the Shenzhen kindergarten eye study. *Invest Ophthalmol Vis Sci* 2015;56:1985–92.
- Morad Y, Werker E, Nemet P. Visual acuity tests using chart, line, and single optotype in healthy and amblyopic children. *J Aapos* 1999;3:94–7.
- Solebo AL, Cumberland PM, Rahi JS. Whole-population vision screening in children aged 4-5 years to detect amblyopia. *Lancet* 2015;385:2308–19.

Clinical science

- 20 Fogel-Levin M, Doron R, Wygnanski-Jaffe T, *et al.* A comparison of plusoptix A12 measurements with cycloplegic refraction. *J Aapos* 2016;20:310–4.
- 21 Koozekanani D, Covert DJ, Weinberg DV. The use of best visual acuity over several encounters as an outcome variable: an analysis of systematic bias. *Invest Ophthalmol Vis Sci* 2010;51:3909–12.
- 22 Holladay JT. Proper method for calculating average visual acuity. *J Refract Surg* 1997;13:388–91.
- 23 Kupl MT, Dobson V, Peskin E, *et al.* Vision in Preschoolers Study G. The electronic visual acuity tester: testability in preschool children. *Optom Vis Sci* 2004;81:238–44.
- 24 Leone JF, Gole GA, Mitchell P, *et al.* Visual acuity testability and comparability in Australian preschool children: the Sydney Paediatric Eye Disease Study. *Eye* 2012;26:925–32.
- 25 Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Front Psychol* 2013;4:863.
- 26 Holmes JM, Beck RW, Repka MX, *et al.* The amblyopia treatment study visual acuity testing protocol. *Arch Ophthalmol* 2001;119:1345–53.
- 27 Kvarnström G, Jakobsson P. Is vision screening in 3-year-old children feasible? Comparison between the Lea Symbol chart and the HVOT (LM) chart. *Acta Ophthalmol Scand* 2005;83:76–80.



Should tumbling E go out of date in amblyopia screening? Evidence from a population-based sample normative in children aged 3–4 years

Sandra Guimaraes, Tiago Fernandes, Patrício Costa and Eduardo Silva

Br J Ophthalmol published online October 7, 2017

Updated information and services can be found at:

<http://bj.o.bmj.com/content/early/2017/10/07/bjophthalmol-2017-310691>

These include:

References

This article cites 25 articles, 5 of which you can access for free at: <http://bj.o.bmj.com/content/early/2017/10/07/bjophthalmol-2017-310691#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiology](#) (1103)
[Open access](#) (292)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>