most recent contribution, to my knowledge, is my paper, "The Mechanism of Aqueous Secretion in Mammalia," which appeared in the British Journal of Ophthalmology in January, 1941. Your reviewer has obviously not made himself acquainted with the demonstration presented there, for his criticisms and strictures are not only ill-judged but illuminated by prejudice and not by the light of reason.

I am, etc.,

THOMSON HENDERSON.

15, THE ROPEWALK,
NOTTINGHAM.
January 15, 1943.

MACULAR HOLE.

To the Editors of THE BRITISH JOURNAL OF OPHTHALMOLOGY.

DEAR SIRs,—May I report another case similar to that described by Lieutenant Commander Pittar?

R. A., aged 13, attended a school clinic on Nov. 4, 1942, complaining of a black spot in his right vision. Vision was 6/12 6/5, the eyes emmetropic. There was oedema of the right macula. Thinking it might be an early central choroiditis I treated him with atropine and rest. The oedema gradually subsided, being replaced by an appearance of abnormal redness. About this time the boy's mother suggested that the trouble might be due to his having looked at the sun without protection.

On Dec. 30, there was a definite macular "hole" with clean cut edges, though vision was 6/18. This retention of fair vision is surprising, but one occasionally meets it in cases of old central choroiditis in which the ophthalmoscopic appearance suggests a total loss of central vision. No doubt it will deteriorate further.

Yours truly,

H. DOUGLAS SMART.

18, WIMPOLE STREET, W.1

SYMPATHETIC OPHTHALMITIS

To the Editors of THE BRITISH JOURNAL OF OPHTHALMOLOGY.

DEAR SIRs,—On p. 572 of your Journal for December, 1942, a letter from Mr. T. Harrison Butler is published in which is described a case supposed to be one of sympathetic ophthalmitis. The interval of time between the date of excision of the exciting eye and that of the first signs of inflammation in the sympathising eye was