A WASP STING

BY

S. W. K. NORRIS

R.A.M.C.

THE following case is sufficiently unusual to be worthy of a report.

On October 26, 1943, Lance-Cpl. S. reported complaining of epiphora, pain and photophobia in the right eye of nearly seven weeks duration. He stated that he had been treated for "kerato-conjunctivitis" in three different hospitals.

The eye was red, watering and difficult to examine because of the blepharospasm evoked by the light. Vertical striae, faintly staining with fluorescein, were present over the upper third of the cornea. A subtarsal foreign body was looked for; all that could be seen, however, was a small papule with a brownish centre in the middle of the tarsal conjunctiva. The skin of the upper lid was normal.

An attempt was made to remove the brown spot in spite of the patient's protestations. Apparently five efforts had already been made to do this, by three different ophthalmologists elsewhere, without success. Again, nothing could be found. Next morning when the capillary ooze had stopped a fine hair was seen and removed from the centre of the papule. "That's the sting" cried the patient, "nobody would believe me."

It then transpired that on September 6, 1943, he had been stung on the right eyelid by a wasp, he had refrained from mentioning it on this occasion as he had been discouraged by the past reception of his story.

The "hair" was sent to Mr. B. Jobling, Entomologist of the Wellcome Bureau of Scientific Research, who kindly reported the specimen to be "the distal part of the lancet of the sting of a wasp." He also made a sketch which showed that the tip of the lancet is armed with projecting notches. These undoubtedly got caught in the tarsal plate and the wasp had been unable to withdraw it.

The next day the patient was symptom free with 6/6 vision in each eye.

I wish to thank Major E. M. G. Galton, R.A.M.C. for his advice and the Command Specialist, S.E.C. Ophthalmic Centre, for his permission to publish the case.