THE BRITISH JOURNAL OF OPHTHALMOLOGY

JUNE, 1947

COMMUNICATIONS

BINOCULAR VISION * †

BY

DOROTHY CAMPBELL

COVENTRY

ALTHOUGH binocular vision has not so far been made the subject of a Middlemore Lecture—certain allied disorders of vision have been discussed by previous lecturers. It is interesting to find that Mr. Fulford Eales—in 1920—deplored the failure of the Education Authorities to deal adequately with cases of visual deficiency, and particularly with those suffering from squint and amblyopia—while Mr. Haycraft, from his experience in examining recruits in 1943, found the incidence of amblyopia so high, that he too deplored the fact that so many apparently healthy eyes should grow up practically blind. Moreover, he regarded it as “a slur on the care which we oculists devote to our young helpless patients.” The interval between the two lectures was that of a complete generation—and it seems to be in keeping with the

* The Middlemore Lecture delivered at the Birmingham and Midland Eye Hospital, October 25, 1946.

† Received for publication, February 3, 1947.

progressive spirit of Richard Middlemore that we should meet this challenge by a consideration of the physiological factors which maintain binocular vision, and an inquiry as to how we can employ this knowledge—to improve the treatment of cases of abnormal binocular vision.

In presenting my thoughts to you I shall undoubtedly repeat certain familiar facts, but I hope to present them to you in fresh perspective, and to include in them an account of recent observations. The one-eyed man is perhaps not now regarded as an oddity, but he is certainly debarred from many interesting activities, while perfect binocular vision is required for so many occupations in the skilled industrial world of the immediate future, that its lack may determine the fate of a man. The high standards of binocular vision required by the Royal Air Force in this last war led to a mass-enrolment of orthoptic trainers, and the work which they accomplished under the guidance of Air Commodore Livingston in restoring binocular vision to pilots—has I think removed the last barriers which existed between ophthalmic surgeons and orthoptists as to the usefulness of orthoptic training. Indeed, it repeats the challenge to ophthalmic surgeons henceforth to make their proper contribution towards the art of orthoptic training—in order that it shall not run the danger of becoming merely a technical exercise.

The common disorders of binocular vision are well known to you—viz. primary amblyopia—and the allied condition of suppression; strabismus—both paralytic and concomitant—and heterophoria.

Paralytic squint, from its sudden onset and its embarrassing clinical symptoms of double vision and false projection, has been recognised by physicians for many years and has been correctly attributed to a pathological disorder of the cranial nerves or nuclei supplying one or other of the extra-ocular muscles. Concomitant strabismus on the other hand was first regarded as a naturally occurring malposition of one eye—and Dieffenbach in 1840 was the first to attempt its surgical correction. Nevertheless, as long ago as 1743 Buffin had suggested that it was due to inequality of vision between the two eyes—while over a century later Priestley Smith ascribed it to a disturbance of those cortical nervous centres which govern binocular vision.

Up to the beginning of this century—the treatment of squint consisted in a correction of the errors of refraction and cosmetic operation. Claud Worth (1902) was the first to inquire into the significance of the amblyopia which is so often present in a squinting eye—and to treat it by occlusion of the good eye—followed by simple binocular fusion exercises on the amblyoscope.

He believed that a fusion centre existed in the brain which was essential to binocular vision, and was absent or lacking in function in cases of concomitant strabismus. The realisation that a faulty state of binocular vision, *viz.*, heterophoria, can exist without an obvious malposition of the eyes—came later still.

Now if, in the light of our present knowledge, we consider the factors which contribute to the maintenance of perfect binocular vision in man, we find that they are (1) The overlapping of visual fields (2) A strongly developed sense of fusion—and (3) The dominance of macular over peripheral vision.

1. *The Overlap in Visual fields.*—Let us first consider the origin of the overlap in visual fields. Most of us are sufficiently familiar with the phylogenetic development of animals, to know that the eyes were first placed laterally—and then assumed a frontal position. We must also bear in mind the fact that in order of development, conjugate movements of the eyes preceded convergence.

What is known as the nasal half of the retina must originally have been occupied with the receipt of visual stimuli arising *behind* the animal—and with the assumption of the frontal position of the eye—this half of the retina is still occupied with stimuli arising on the outer side of the body. In the first case the reaction of the animal as a whole will be to move away from the object behind it, and in the second case—the animal will be stimulated to make divergent movements of the eyes.

The temporal halves of the retinae, on the other hand, are concerned with the reception of stimuli arising *in front of* the animal. The eyes are mutually attracted instead of being distracted—and there is every incentive for the eyes to converge in an attempt to locate the object more precisely. When the visual axes of the two eyes meet—there is a considerable overlap of the visual fields—which must increase the dominance of the temporal halves of the retinae and strengthen visual attention in the binocular area. A re-arrangement of the optic nerve fibres occurs first in the cats, when instead of there being a total decussation of fibres—the semidecussation with which we are familiar is instituted. It is noteworthy that this anatomical linkage appears to subserve conjugate vision; while the non-decussation of fibres from the temporal halves of the retinae favours binocular vision.

Another interesting anatomical fact is that the power of *convergence* only occurs in animals which possess maculae—*i.e.*, the higher monkeys and man. It is associated with a great development of the brain. Not only are there visual sensory areas in the occipital cortex, but a prefrontal cortical area which is associated

with the acquisition of a wide range of conjugate movement and with convergence. In fact, Kennard (1940) found that destruction of one entire frontal lobe in monkeys had the same effect as destruction of one occipital area, in that the animal did not appear to see objects in a contralateral field of vision. In other words though the sensory stimulus was undoubtedly being received by the animals, appropriate mental reaction was stopped. The *localisation* of an image in a visual field is not then purely a sensory phenomenon but is a sensory-motor response. Holmes has reported two cases in which impairment of the parietal lobe in a patient caused defective spatial orientation—although the visual fields were unimpaired and visual acuity was fair. Riddoch (1935) has pointed out that visual disorientation can occur when the association pathways from the occipital cortex to other parts of the brain are interrupted, the nodal point appearing to be in the region of the supramarginal and angular gyri in the parietal lobe.

This new anatomical evidence is against the theory held by so many ophthalmologists that spatial localisation arises purely from the position of the stimuli which fall on the retinae, or from any exactly recorded pattern of them in the visual sensory cortex.

It is probably the correct conclusion to say that the overlap of visual fields lays the foundation of binocular vision, by concentrating visual attention, so that an object can be located correctly, and that the added powers of convergence and accommodation—serve to make the location precise. The mere overlap of fields does not guarantee the fusion of two images into one. For this a fusion sense appears to be necessary—and is intimately bound up with the sense of projection.

Fusion Sense.—Paralytic squint is the most dramatic instance of a breakdown of fusion. It arises from a motor defect in the extra-ocular muscles, which disturbs the normal position of the visual axes. An attempt has been made to explain the accompanying false projection, also on the basis of a failure of *motor* innervation. It was said to be due to the fact that "When an ocular muscle is weak the increased innervation of the opposing muscle gives an impression of a greater movement of the eye than has really taken place; thus the mind receives the idea that the object is placed farther to one side than it really is, and in an attempt to touch it, the finger may go beyond it." Even so eminent a physiologist as Gowers stated that "We judge of the relation of external objects to each other by the relation of their images on the retina; but we judge of their relation to our own body by the position of the eyeball as indicated to us by *the innervation we give to the ocular muscles.*"

While such an argument might conceivably be true for paralytic

squint, in which defective orientation is partly overcome by secondary deviation of the other eye, and by alteration of head-posture, it is contrary to the facts to interpret all abnormalities of binocular vision as a failure of motor innervation. In concomitant squint the visual axes retain the same relationship to one another over a wide range of conjugate movement—and a *sensory* disturbance occurs in the form of amblyopia or suppression in the majority of cases. Worth explained the onset of suppression as being due to the absence of a fusion centre in the brain—but there is no physiological or anatomical evidence for the existence of such a centre. There is obviously a *sensory-motor* balance in binocular vision—and Franklin (1942) has also emphasised the idea that the functions of fusion, projection and stereopsis depend on this balance—rather than on visual sensations which arise from retinal stimuli alone.

He points out that before the development of the eyes the movements of the body were guided by the impressions received by the lateral sense organs, and that in course of time the eyes took on the same function as the sense organs. We have further evidence of this in the fact that no bird will fly in a fog, because it loses its directional sense, and that a blindfolded pigeon, when loosed from an aeroplane, fails to fly, and drops like a stone.

In animals with laterally placed eyes in which the binocular fields do not overlap, *e.g.*, herbivores and fishes, the animal shows that it can see an object ahead by moving up to it and seizing it. In the carnivores, which have frontally placed eyes—there is a definite correlation between vision and the use of the forelimbs, in the catching and killing of fast-moving prey. The mode of life adopted by the primates, of swinging by hand from branch to branch—requires an exact sense of distance—and projection of an object in space, and we find the anatomical response—of specialisation of the brain, convergence, accommodation and macular vision to which I have already referred.

Thus, in animals, in the fundamental association of body action with visual stimuli, each forelimb is correlated with vision in the *contra*-lateral field. In man, the intimate association of body movement with visual stimuli is well shown in the lack of general co-ordination which often accompanies paralytic strabismus, and is also seen in certain cases of concomitant strabismus when the patient attempts to rely on the amblyopic eye alone.

Franklin has emphasised the fact that fusion and projection in binocular vision are based on *bitemporal* retinal representation—*i.e.*, on the overlapping of nasal fields, but that they are anticipatory in character—and comparable to potential movements.

Let us now consider the third factor which influences binocular vision—*viz.*, *macular dominance*.

Birds with laterally placed eyes, whose visual axes are set at an angle of 120° or more, have a fovea, so that it would appear that the primary function of the fovea cannot be one of fusion. Nocturnal birds of prey, with frontally placed eyes, have two foveae in each eye.

The macular fibres in man are distributed over an area which extends for approximately 20° in a horizontal direction on either

FIG. 1.

The course of the visual fibres of the retina.

side of the macula and embraces the blind spot on the nasal side (Fig. 1). Vertically the area extends for about 3° above and below the macula. Visual acuity within this central area rises from 6/60 at the periphery to 6/6 at the fovea (Fig. 2).

On account of the superiority which this visual acuity and pattern vision has conferred on man—we are apt to consider that it dominates over the more primitive visual function of fusion. It is important to remember in this respect, that bitemporal retinal representation persists throughout the central area of the retina—and the overlap of the visual fields must be as shown in Figs. 3 and 4. One would expect to find, as is indeed the case, that fusion is very strong in the paramacular area, and it decreases peripherally.

FIG. 2.

The regional variation of visual acuity (after Wertheim).

Overlap of visual fields in binocular fixation (central area).

FIG. 3.

The overlap of visual fields in man.

□ Binocular vision. ■ Monocular vision.

n.r. = nasal limit for "central field," right eye. n.l. = nasal limit for "central field," left eye.

The maintenance of perfect binocular vision must depend on a bilateral harmony within the central area, of the two mechanisms of pattern vision and fusion. This probably occurs only in a minority of individuals, since the possibilities of the breakdown of so fine an adjustment are so many. It has in fact been said (Bielchowsky, 1937) that only about 20 per cent. of individuals with good binocular vision are really orthophoric, and recent observers have found evidence of a continual competition between foveal vision and the fusional processes occurring in the area immediately outside the fovea.

In certain pathological conditions there is a surprising maintenance of binocular vision; *e.g.*, subjects suffering from retinitis pigmentosa, in which the function of the peripheral retina is entirely lacking, can still maintain binocular vision when only a small central field remains and their visual acuity is very low. Conversely subjects with bilateral macular degeneration or disease also may retain some degree of fusion. A study of binocular vision in conditions of partial loss of visual field, would, I think, elucidate the functional balance between the peripheral and central areas of the retina, still further.

In a series of tests on another aspect of this problem, which have been made recently at this hospital, we have found that good binocular vision persists in the dark adapted eye—even when the illumination of the test object is so low as to favour rod, rather than cone, vision. The subjective angle tends to become convergent; there is a general tendency to giddiness, and colours are

suppressed. Some degree of stereopsis persists even when visual acuity is greatly diminished,

The *physiological* abnormalities of binocular vision (see Table 1) may be tabulated as shown. We find all degrees of variation, from a complete competition between the eyes, as in alternating strabismus, down to total suppression of central vision in one eye. Thus at each end of the scale—vision becomes uniocular rather than binocular. In the intermediate grades some attempt

TABLE I
Physiological Abnormalities of Binocular Vision

1. Alternating strabismus	...	Uniocular vision
2. Congenital nystagmus	...	Usually some degree of binocular vision
3. Heterophoria Low degrees of concomitant strabismus False associated fixation	} ...	Abnormal binocular vision due to rivalry within the "central" area of the retina
4. Large degrees of concomitant strabismus (a) with, and (b) without suppression	} ...	Abnormal function in both "central" and peripheral retina
5. Advanced suppression with loss of central fixation	} ...	Peripheral function only in affected eye; binocular vision poor or absent

at binocular vision is made. In by far the majority of cases this is accomplished by the partial suppression of pattern vision—which has been abundantly proved to be a mental process of ignoring the image from one eye. In others, although visual acuity is full, foveal fixation gives way in order that extra-foveal fusion can be maintained; or competition is avoided by establishment of a false macula—in the condition we recognise as false associated fixation.

When the central area of one eye is very defective—abnormalities of orientation occur. The onset or non-occurrence of strabismus is, I believe, determined at its onset by the initial level of visual acuity in the eye affected. By referring again to the curve for visual acuity (Fig. 2) we are reminded that the central area includes the blind spot, so that "the margin of escape" is only 10° on the nasal side of the macula—and any failure in this region will favour the development of a convergent strabismus.

You will notice that I have so far avoided making any reference to the subject of retinal correspondence. It has long been considered that stereopsis was due to the stimulation of corresponding points on the retina, and to an exact reinterpretation of such stimuli in the visual sensory cortex. I have already shown that this is not true in the case of the primitive fusional mechanisms—and recent work has shown that it is too precise a theory to be applied even to foveal fixation.

McAllister (1905), Dodge (1907) and Clark (1936) have shown that in normal binocular vision an area of $1-1.5^\circ$ from the fovea

FIG. 5.

Diagram showing the doubling of a centrally fixated object (1° square) by displacement of peripheral images (print), A, superimposed, B, displaced (Burian, 1939).

can be used for fixation—and with a disparity of $2-4^\circ$ horizontally and 2° vertically in either direction an object will still be reported as single. Burian (1939) made a further discovery that peripheral retinal stimuli, if displaced, by as small an interval as 0.25° , will break the fusion of images situated on corresponding areas of the macular region, and no matter how hard the observer tries he is unable to maintain central fusion in the presence of displaced peripheral stimuli (Fig. 5). Burian made these tests within an area of 12° round about the macula. Stimulation of smaller areas of $\frac{1}{2}-1^\circ$ at a distance of 12° from the fovea caused noticeable fusional movements of the eyes. Burian concluded from his observations that there must be a large number of individuals who will never be able to fuse central and paracentral images simultaneously; some in fact maintain a rapid small alternation.

He later (1941) examined 75 cases of concomitant strabismus with less than 12° strabismus and full visual acuity. He found that the eyes performed small fusional movements in order to maintain

binocular vision, and that patients who had no foveal suppression, showed a small degree of suppression in the paramacular area. Paramacular fusion could only be maintained at the expense of foveal fixation being out of alignment. The exceptions were (a) alternation, due to suppression of all the images of each retina

FIG. 6.

Peripheral fusion (of superimposed line pattern) and foveal test lines on a level. Images seen as above, *i.e.*, unchanged by (a) Patient with normal binocular vision. (b) Patient with strabismus with anomalous correspondence adapted to the angle of squint (Burian, 1941).

in turn; (b) The presence of a false macula, which supported all the attributes of binocular vision except stereopsis (Fig. 6). His methods of experiment were both accurate and ingenious.

A comparison of the average number of eye-movements made per second, while the eyes "fixed" isolated points and points in context. (Clark), 1936.

(By kind permission of Professor Clark, *Amer. Inst. Psychol.*, 1939)

	Horizontal component		Vertical component	
	R. E.	L. E.	R. E.	L. E.
Isolated point A	7.55	7.01	4.11	6.20
Point A in context ... (<i>i.e.</i> viewed in a picture)	6.88	7.59	3.88	6.39
Isolated point B	6.98	6.66	4.28	5.53
Point B in context	7.08	6.55	3.83	5.44

FIG. 7.

The improvement of the camera has made it possible to make objective records of the movements of the eyes—rather than having to rely entirely on subjective phenomena.

Judd (1907) was the first to photograph the movements of the eyes in convergence and divergence. He found that as they viewed two fixed points in space, the eyes of some subjects made conjugate movements prior to convergence, while others moved one eye faster than the other. In short there was no exactitude of movement in fixation. The same observation was made by Clark (1936) who from an analysis of over 3,000 photographic records found that the quality and quantity of the movements was the same for stereoscopic vision, as in the fixation of an isolated point (Fig. 7). The eyes made an average of 8 movements per second, but one eye moved more often than the other (Fig. 8). Clark

Average number of movements per second made by the eyes
fixing points in a Stereogram (Clark).

(By kind permission of Professor Clark, *Amer. Inst. Psychol.*, 1939)

Fixating					Fixating				
Near Point			Far Point		Near Point			Far Point	
Subject	R.	L.	R.	L.	Subject	R.	L.	R.	L.
1	5.9	9.9	10.2	13.8	9	5.3	8.1	3.3	7.0
2	5.1	5.8	6.0	6.2	10	4.1	7.0	11.9	9.2
3	11.8	13.7	15.9	14.1	11	9.0	6.8	8.2	7.2
4	7.7	9.2	9.1	9.9	12	5.8	2.4	3.1	3.8
5	7.1	9.8	8.9	12.0	13	8.3	6.9	8.8	10.2
6	9.8	7.2	8.7	10.9	14	11.1	9.9	10.0	9.1
7	9.0	11.7	5.9	5.8	15	5.2	7.1	5.1	5.2
8	7.9	8.3	6.2	10.8	16	—	—	8.8	11.8

FIG. 8.

concluded that clear visual perception can only occur at the movement of fixation—and in the interval visual acuity must be diminished. Obviously then, visual perception must constantly be both binocular and monocular, and shift from one eye to another. He suggests that the highest grade of binocular vision must therefore depend on the organisation and unification of minimal changes—rather than on the stimulation of specific geometrical or corresponding points on the retina. He believes that stereopsis must be made up of a rapid comparison of uniocular impressions, and in this he is supported by Litinski (1939) who found that in 72.5 per cent. of normal subjects which he examined, the uniocular perception of solidity was equal to the binocular perception. Verhoeff too supports the theory that binocular vision occurs by uniocular replacement—rather than by simultaneous vision of the two eyes. He believes too, that while one eye is occupied with the observations of one portion of space, the other eye is suppressed in this region only—so that the two eyes divide between each other the entire field of vision in an ever-changing mosaic of small uniocular part-fields. Verhoeff's conclusions were based on subjective phenomena—but the objective experiments of Burian and Clark go far to substantiate them. In an analysis of amblyopia which I made earlier this year (1946) I arrived at the same conclusion, *viz.*, that binocular vision depends on some form of physiological replacement. It is certainly the only theory which explains all the abnormalities of binocular vision, particularly those which arise from functional disorders of the central area of the retina. Thus Hering's theory which states that in binocular vision the image is projected along the visual axis of a hypothetical cyclopic eye—midway between the two eyes—appears obsolete. The elaboration of this theory to explain stereopsis as due to the simultaneous stimulation of corresponding and disparate points of the retina also appears to me to be impossible. I was surprised to find that the latest definition of stereoscopic vision which has been given to orthoptic students is "the ability to fuse images falling on *slightly* disparate points of the retina." It is difficult to reconcile this with any theory of binocular vision.

In conclusion, I shall attempt to apply these newer concepts of binocular vision to the practice of orthoptics.

The young child resembles the animal in its development, and its early years are spent in learning to correlate its body reactions with visual stimuli.

I cannot resist quoting from Richard Middlemore's "Treatise on diseases of the eye," certain passages in which he lays stress on the very early onset of strabismus (1835).

In early life, children are sometimes placed very injudiciously in their cradle, or they are, perhaps, very generally carried in one and the same position, so that the light gains more frequent and complete access to one organ than to the other, in consequence of which, that eye which is most familiarized with the influence of the luminous rays, is more readily stimulated by them, and the other is rendered, in a corresponding degree, the more insensible to them. Again, sometimes the infant's cap is ornamented with a long and partly coloured border, so that it readily attracts the child's attention, and he is continually turning the eyes towards it, as he lies in the cradle. In the same way a small spot, naevus, or tumour upon the nose, may produce squinting by disposing the child to look at it very intently, and he is inclined to make great and repeated efforts to turn the eyes towards the object of attention, because it is not distinctly seen. Of these two causes of infantile strabismus, the former will cause strabismus divergens, and the latter strabismus convergens.

7.—Imitation is certainly a cause of strabismus; parents will often tell their medical attendant that their child did not squint when he left home for school, but, on his return, the strabismus was complete; and the child will probably mention that many of his school-fellows also squinted, because they were frequently mocking or imitating each other. No cause of strabismus is more satisfactorily made out than this.

8.—Sometimes one of the eyes happens to be inflamed in early life, and it is carefully bound up, until its visual powers become impaired from prolonged repose from its natural actions and the long absence of its natural stimulus; such an eye will be liable to be affected with strabismus, unless, which is very improbable, its retina should again recover its sensibility. In other instances both eyes may, perhaps, be slightly inflamed, and the surgeon may tell the parents of the child to protect them from the influence of bright light, and they will bind them up, as they conceive, very carefully, in order to fulfil what they presume to be the Doctor's wish the better, and in this way the patient will be totally deprived of his sight during the continuance of the ophthalmia, at least, according to the parent's intention; but children do not like this deprivation, and therefore so change the situation of the bandage, as to see either upwards or downwards, and, in this way, acquire a habit of squinting. Many of the worst cases of strabismus of both eyes, which it has fallen to my lot to notice, have been so caused.

It is certainly a pity to lose the early years as a time for training. The sense of fusion can only be properly developed if the two eyes are used equally. Visual acuity should be brought to a maximum by ordinary glasses at as early an age as possible—and occlusion should not be left, as it so often is in the hospital child, until the age of 5 (Fig. 9). It can be employed much earlier—or perhaps a better method is the use of atropine drops in the good eye, in order to bring about fixation in the squinting eye. I have never known this method to cause false associated fixation to develop—and it has the advantage that it can be started in infancy—and that the child continues to exercise some degree of binocular vision—whereas occlusion cut out any stimulus to fusion. There is a good deal of evidence that the amblyopic eye supplies a spatial background to the good eye, and this must be the explanation as to why quite a number of adult amblyopes have straight eyes and a certain degree of fusion.

The next stage of fusion training is not usually started until about the age of 6, because the child cannot understand the synoptophore and has not sufficient mental application. I think more use might be made of the toy games in the interim.

The development of visual acuity in the young child, following treatment. (Campbell, 1946)

TREATMENT UNDER 5 YEARS					TREATMENT OVER 5 YEARS	
Group A			Group B		Group C	
Vision	Before	After	Before	After	Before	After
	per cent.	per. cent.	per cent.	per cent.	per cent.	per cent.
<6/60 ...	15	0	27	0	10	5.2
6/60-6/24	28	2.4	38	2.2	43	5.2
6/18-6/9	57	9.6	35	20.0	47	12.0
6/6 ...	40	88.0	0	77.8	0	77.6
No. of cases	47	Treated by atropine drops followed by occlusion	45	Treated by occlusion only	58	Treated by occlusion only

FIG. 9.

Again fusion training is delayed until vision is nearly equal in both eyes, but Swan and Laughlin (1944) have recently advocated it, when the vision in one eye is still much below standard. They state that provided that the target is large enough to correspond with the degree of vision, or to avoid the central scotoma—peripheral fusion can be established and appears to assist in the development of central vision.

Cases of heterophoria excluding large degrees of height, seem to improve adequately on duction exercises. I have found that quite a number, *e.g.*, 42 out of 78 cases, show a small degree of suppression—which is frequently alternating and appears only in the fixing eye. This suppression is no obstacle to their cure—because of the strong fusion sense which is always present.

False associated fixation presents one of the most difficult problems of training. In a group of 74 cases I found only one cure. The data showed that they all had, or regained, equal vision in both eyes—and showed a macular dominance, but with very little power of fusion. The subjective angle in every case except one

was under 10° —*i.e.*, it corresponded to the retinal area between the disc and the macula, while in one case only the angle was 19° —*i.e.*, immediately to the nasal side of the disc.

It has been suggested that in these cases operation should precede orthoptic training, but what is needed appears to be some form of training to develop peripheral fusion and by the use of large images, rather than to attempt to change macular vision from "false" to "true," a process which takes an exceedingly long time and has such strikingly poor results. Events might prove that the resumption of peripheral fusion would lessen the disparity in central fixation.

Finally, although the standard set by the British Orthoptic Council for a cure is quite commendable, I think it is far too high for every day purposes.

Costenbader (1945) in an analysis of the causes of failure in the treatment of squint, has suggested 4 grades of cure :—

- (1) British Standard (1939). Straight eyes, normal vergences—good fusion and stereopsis.
- (2) Good fusion and stereopsis, but varying from fusion to diplopia with Worth's test, and inconstant ability to bar read.
- (3) Fair fusion—fair vision and a small degree of residual strabismus.
- (4) Cosmetic cure—no fusion—a small degree of residual strabismus.

You will notice that fusion with convergence is only achieved in the highest degree of binocular vision. Stutterheim (1940) has advocated a new form of training based solely on convergence exercises. He starts, as it were, at the other end of the scale—and states that amblyopia may persist for a considerable time *after* the squint has been cured. Although he pours scorn on the recognised methods of orthoptic training, his book is worth reading, by way of contrast.

I have not attempted to analyse the surgical treatment of binocular abnormalities, but I think you will agree that good surgical results depend more on the moment chosen for operation, rather than on the exact adjustment of the angle of squint. I must end my discourse with an apology for the inadequate analysis of the subject of binocular vision, which I have been able to make. Perhaps some Middlemore lecturer of the future will be able to give us more advanced information. The present jig-saw of facts will then be transformed into a complete picture.

REFERENCES

- BIELCHOWSKY, A. (1937).—The etiology of squint. *Amer. Jl. Ophthal.*, Vol. XX, p. 478.
- BURIAN, H. M. (1939).—Fusional movements. *Arch. Ophthal.*, Vol. XXI, p. 486.
- (1941).—Fusional movements in permanent strabismus. A study of the role of the central and peripheral regions in the act of binocular vision in squint. *Amer. Arch. Ophthal.*, Vol. XXVI, p. 626.
- CAMPBELL, D. A. (1946).—Primary amblyopia. *Trans. Ophthal. Soc. U.K.*
- CLARK, B. (1936).—An eye movement study of stereoscopic vision. *Amer. Jl. Psychol.*, Vol. XLVIII, p. 82.
- COSTENBADER, F. D. (1945).—Causes of failure in the treatment of squint. *Amer. Jl. Ophthal.*, Vol. XXVIII, p. 1123.
- DODGE, R. (1907).—An experimental study of visual fixation. *Psychol. Monogr.*, 1.
- EALLES, FULFORD W. H. (1920).—Some aspects of convergent strabismus. Middlemore Lecture.
- FRANKLIN, I. (1942).—Fusion projection and stereopsis. *Amer. Jl. Ophthal.*, Vol. XXV, p. 1316.
- HAYCRAFT, G. H. (1943).—Amblyopia ex anopsia (with special reference to recruitment for the armed forces). Middlemore Lecture.
- HERING (1863).—See DUKE-ELDER, Text-book of Ophthalmology, Vol. I, p. 1062.
- JUDD, C. H. (1907).—Photographic records of convergence and divergence. *Psychol. Monogr.*, pp. 6, 370.
- KENNARD, M. A. (1940).—Alteration in response to visual stimuli following lesions of frontal lobe in monkeys. *Arch. Neurol. and Psychiat.*, Vol. XLI, p. 1153.
- LITINSKI, G. A. (1939).
- MCALLISTER (1905).—The fixation of points in the visual field. *Psychol. Monogr.*, pp. 7, 17.
- RIDDOCH, G. (1935).—Visual disorientation in homonymous half fields. *Brain*, Vol. LVIII, p. 376.
- STUTTERHEIM, N. A. (1940).—Squint and convergence. H. K. Lewis & Co., Ltd., London.
- SWAN, K. C. and LAUGHLIN, E. (1944).—Binocular orthoptic training for amblyopic patients. *Amer. Arch. Ophthal.*, Vol. XXXII, p. 302.
- WORTH, C. (1902).—Squint, its causes, pathology and treatment. Sixth edition, 1929.
- VERHOEFF, F. H. (1935).—A new theory of binocular vision. *Arch. Ophthal.*, Vol. XIV, p. 419.
- MIDDLEMORE, R. (1835).—A treatise on the diseases of the eye.

THE REGENERATION OF WOUNDS OF EXTERNAL MEMBRANE OF THE EYE IN THE LIGHT OF NEW PATHOLOGICO-ANATOMICAL RESULTS*†

BY

Professor E. TH. LEVKOIEVA

MOSCOW

PART I

Introduction

WHILE paying due consideration to the questions raised by ophthalmic surgery during the last war, we have to point out that insufficient attention was paid to the conditions at the moment of

* From the pathological-histological Department of the Central Institute of Ophthalmology, named after Helmholtz Director, Professor Kolen; in charge of pathological-histological department, Professor Levkoieva. Translated by Dr. N. Pines. † Received for publication, November, 1946.