CORRESPONDENCE

CLINICAL GLAUCOMA

To the Editorial Committee of the BRITISH JOURNAL OF OPHTHALMOLOGY

SIRS,—In my recently published book "Clinical Glaucoma" [reviewed on p. 335], in the section dealing with Applanation Tonometry, I have stated that the pressure should be read off at the diastolic point of the pulsation of the green half-circles. Professor Goldmann has kindly written to me to tell me that this is not what he advises. He always takes the mean value between the systolic and diastolic. I would like to bring this important point before readers, because discrepancies might otherwise arise and I would also like to apologise to Professor Goldmann for misrepresenting his instructions regarding the use of his superb instrument. He tells me that the English version of the instructions for use of the applanation tonometer provided by the manufacturers is being rewritten to make the point clear.

Yours faithfully,
REDMOND J. H. SMITH

46 WIMPOLE STREET,
LONDON, W.1.
April, 1965.

BOOK REVIEWS

Stereoscopic Manual of the Ocular Fundus. By F. C. BLODI, L. ALLEN, and A. E. BRALEY. 1964. Pp. 132, 206 illus. Henry Kimpton, London. (£12 3s. 6d.)

If any ophthalmologist wearies of ophthalmology he should acquire this book; to him, to his family, and to their guests after dinner it could be a never-ending source of joy. In recent years there has been a spate of photographic atlases of the ocular fundus, many of them excellent, a break-through rendered possible by finer cameras, faster colour-film, and better lighting; but this is the first stereoscopic atlas in colour. Complete with its fifteen reels of seven pairs of stereo-photographs arranged in two neatly packed rows on the inside of the front cover and a collapsible stereo-viewer (with full instructions) tucked in an envelope fixed on to the back cover, the delighted owner of the book can study beautiful and dramatic pictures of the normal fundus, its congenital anomalies and age changes, and the pathological changes in the more common diseases. At the same time a short text tells the story of each individual case illustrated, and excellent histological illustrations are provided to explain the underlying pathology. The authors, the ophthalmic department of Iowa College, and the publishers—everyone in fact—are to be congratulated.

Contribution to Ocular Pathological Anatomy. (Travaux d'anatomie pathologique oculaire.) By F. HERVOUËT. 1964. Pp. 172, 263 figs, bibl. Masson, Paris. (52 F.)

The third of a series of monographs on ocular pathology written by Hervouët from the clinic at Nantes fully maintains the excellence of its predecessors. There is an interesting description of the effect of alpha-chymotrypsin on human eyes on enucleation: fragmentation of the zonule occurs rapidly, as effectively after 30 seconds as after 40 hours, so that the enzyme need not be retained within the eye longer than one minute, when the anterior chamber should be washed out. Other interesting subjects are epithelial ingrowth into the anterior chamber after cataract surgery; Fuchs's combined corneal dystrophy; intra-ocular myiasis; periphlebitis of the retina and retinal detachment—in which the studies of the pigmentary epithelium and the vitreous are especially interesting. Congenital lesions are represented by two complete studies on colobomatous eyes and two cases of microcoria. The histological illustrations are particularly good.