blocking the filtration angle. But this form of treatment is manifestly too dangerous for other than experimental work; Lagrange thinks that the escape can be sufficiently limited by encouraging the formation of a band of fibrous tissue on the outside of the sclerotic, round about the region of the filtration angle. If the escape is limited, the tension will rise. Certain cases have been observed by Kummel, Guarino, and Lagrange himself in which glaucoma has followed on extensive scarring of the conjunctiva.

Guardino is quoted as expressing the opinion that the high frequency of glaucoma in Egypt is due to the cicatricial state of the limbal conjunctiva in old trachoma.

Lagrange adopts a measure which he calls colmatage; it consists in burning the sclerotic deeply, through the conjunctiva, in a number of points—20 or more—arranged in three rows round one half of the limbus; later, if necessary, the other half is treated similarly. This is followed by a series of subconjunctival injections.

He produces the notes of a number of cases in which detachment of the retina was cured by these means.

As a means of treatment of this disastrous condition, anything that will allow us to raise the intraocular tension deliberately, and to a definite extent, holds out large hopes.

The book as a whole is interesting and stimulating; there is much that is controversial in it—some of this has been indicated here—but no one can doubt the sincerity of the author or deny the great debt which the world owes him for his operation of limbal sclerectomy as a means of combating chronic glaucoma.

The book loses from the absence of an index.

HAROLD GRIMSDALE.

CORRESPONDENCE

VENOUS PULSATION

To the Editor of THE BRITISH JOURNAL OF OPHTHALMOLOGY

SIR,—I am unaware whether it is on record that in the majority of cases of physiological retinal venous pulsation—as far as I have gathered from observation on about a dozen cases—the pulsation ceases when the eye is turned to the extreme lateral limit in either direction. The direct view of the disc under such circumstances is

the more easily secured with a self-luminous ophthalmoscope, for the obvious reason that the observer is thereby independent of picking up the light at a difficult angle.

Yours truly,
BASIL GRAVES.

February 9, 1922.

COLLOSOL ARGENTUM (FOR THE MEDICAL PROFESSION ONLY)

To the Editor of THE BRITISH JOURNAL OF OPHTHALMOLOGY

SIR,—It was with great surprise that I read in the letter of Mr. M. S. Mayou in the April number of the journal, that he found the use of collosol argentum causes pain and irritation.

I have continued to use it ever since I wrote my article on the treatment of ulcer of the cornea and interstitial keratitis in some 40 to 50 cases daily.

As regards the pain, I have had only one complaint and that was from a neurotic old lady with ulcer of the cornea. Subsequently she had several relapses and made no complaint.

It is obvious that a considerable proportion of the above cases would be children, therefore exceedingly apt to resent anything of the kind.

The formula I use is 12 minims of sol. of scopolamine 1 in 800 to two drachms of collosol argentum dropped into the eye five times a day and the eye kept closed for five minutes afterwards.

If this is not sufficient to keep the pupil dilated a separate supply of the mydriatic is given.

As regards the irritation mentioned, I have never met any. In ophthalmia neonatorum it is used as supplied. It must be somewhat difficult to determine as to whether or not it causes pain in an infant, especially when combined with other treatment as recommended by me.

Possibly the method of its use may account for this great discrepancy.

For some time during the war and after, I found it to vary in colour from light sherry colour to almost black; recently it has been more uniform. I have never been able to get a satisfactory explanation as to why it should have been so.

I am, Sir, Yours faithfully,
A. LEGGE ROE.

HULL, April 3, 1922.