Anterior chamber depth in eyes with pseudoexfoliation

R. S. BARTHOLOMEW

From the University Department of Ophthalmology, Edinburgh

SUMMARY The depth of the anterior chamber was measured in 34 eyes with pseudoexfoliation but without glaucoma and in 334 normal controls. No significant difference was found.

Angle-closure glaucoma is likely to develop in eyes with shallow anterior chambers and narrow angles.¹ Primary angle-closure glaucoma is thought to be rare and to occur by chance association in eyes with pseudoexfoliation.²⁻⁶ Glaucoma with pseudoexfoliation may, however, present with symptoms suggesting angle closure,⁷ have a high incidence of narrow angles⁸—perhaps caused by phospholipid iodide—or be associated with pupil block⁹ or iridocapsular block.^{10 11}

The anterior chamber depth is similar in eyes with primary open-angle glaucoma and glaucoma associated with pseudoexfoliation^{12 13} and in eyes with and without pseudoexfoliation¹⁴—in an admittedly small sample.

This paper presents the results of comparisons between the anterior chamber depths of non-glaucomatous eyes with and without pseudoexfoliation.

Material and methods

The anterior chamber depth of 334 normal eyes and 34 eyes with pseudoexfoliation was measured on a randomly chosen sample during an epidemiological survey of the Negroes of Pondoland, South Africa (this survey is fully reported by Bartholomew¹⁵). Measurements were made with the Goldmann pachymeter attached to the Haag-Streit slit-lamp on right eyes only and excluded those with glaucoma. The corneal thickness and the combined depth of cornea and anterior chamber were measured. The anterior chamber depth was then obtained by subtracting the corneal thickness from the combined measurement. The mean of 3 readings was recorded.

Correspondence to Dr R. S. Bartholomew, Department of Ophthalmology, University of Edinburgh, The Eye Pavilion, Chalmers Street, Edinburgh EH3 9HA.

Table 1 *Anterior chamber depths in 334 normal eyes without pseudoexfoliation by age and sex*

Age group	Males			Females			Significance levels	
	No.	Mean (mm)	SD	No.	Mean (mm)	SD	t	2P
40-49	44	2.74	0.29	44	2.61	0.24	2.17	<0.05
50-59	48	2.73	0.23	40	2.49	0.28	4.00	<0.001
60-69	39	2.61	0.30	36	2.52	0.29	1.32	>0.05
70+	44	2.60	0.32	39	2.51	0.32	1.31	>0.05

Table 2 *Anterior chamber depths in 158 normal eyes and 34 eyes with pseudoexfoliation*

	Pseudoexfoliation			Normal			Significance levels	
	No.	Mean (mm)	SD	No.	Mean (mm)	SD	t	2P
Males	17	2.58	0.36	83	2.60	0.30	0.21	>0.8
Females	17	2.46	0.32	75	2.51	0.30	0.59	>0.5

Results

The anterior chamber depths of the control sample are shown in Table 1.

There was a decrease in anterior chamber depth with increasing age for men ($t=2.15$, $P<0.05$) and for women ($t=1.6$, $P>0.05$). The chamber depth was greater in men than women for those between 40 and 59 years of age but was not significantly different in those over 60 years of age.

The anterior chamber depths in eyes with pseudoexfoliation were all measured in persons over 60 years of age and are shown separately for men and women in Table 2. The measurements are compared with those from controls over 60 years of age.

The observed differences were very small and not statistically significant.

Discussion

There was no significant difference in the depth of the anterior chamber of eyes with or without pseudoexfoliation. This confirmed the findings of Forsius *et al.*¹⁴ It is unlikely that eyes with pseudoexfoliation have the anatomical predisposition of a shallow anterior chamber, which is common in primary angle-closure glaucoma.

If angle closure does develop in association with pseudoexfoliation, some other mechanism must be responsible, as suggested by Herbst,⁹ Bartholomew,¹⁰ and Dark.¹¹

This work was supported by the University of the Witwatersrand and the South African National Council for the Blind.

References

- ¹Törnquist R. Shallow anterior chamber in acute glaucoma. *Acta Ophthalmol* 1953; Suppl. 39.
- ²Gradle WS, Sugar HS. Glaucoma capsulare. *Am J Ophthalmol* 1947; **30**: 12–9.
- ³Ross R. A case of acute congestive glaucoma with exfoliation of the lens capsule. *Acta Ophthalmol (Kbh)* 1949; **27**: 475–82.
- ⁴Audibert J. (1957). La pseudo-exfoliation capsulaire sénile. Etude anatomoclinique et pathogénique. Thèse, Lyon.
- ⁵Tarkannen A. Pseudoexfoliation of the lens capsule. A Clinical study of 418 patients. *Acta Ophthalmol (Kbh)* 1962; Suppl. 71.
- ⁶Lowe RF. Primary angle closure glaucoma with capsular exfoliation of the lens. *Br J Ophthalmol* 1964; **48**: 492–4.
- ⁷Gillies WE, West RH. Pseudoexfoliation of the lens capsule and glaucoma. *Aust J Ophthalmol* 1977; **5**: 18–20.
- ⁸Layden WE, Shaffer RN. Exfoliation syndrome. *Am J Ophthalmol* 1974; **78**: 835–41.
- ⁹Herbst RW. Angle closure glaucoma in a patient with pseudoexfoliation of the lens capsule. *Ann Ophthalmol* 1976; **8**: 853–6.
- ¹⁰Bartholomew RS. The effect of cataract extraction on the intra-ocular pressure of normal eyes and eyes with pseudoexfoliation. *Trans Ophthalmol Soc UK*, in press.
- ¹¹Dark AJ. Cataract extraction complicated by capsular glaucoma. *Br J Ophthalmol* 1979; **63**: 465–8.
- ¹²Hørvén I. Exfoliation syndrome. Incidence and prognosis of glaucoma capsulare in Massachusetts. *Arch Ophthalmol* 1966; **76**: 505–11.
- ¹³Ehlers N, Hansen FK, Aasved H. Biometric correlations of corneal thickness. *Acta Ophthalmol (Kbh)* 1975; **53**: 652–9.
- ¹⁴Forsius H, Sveinsson K, Als E, Luukka H. Pseudoexfoliation of the lens capsule and depth of anterior chamber in Northern Iceland. *Acta Ophthalmol (Kbh)* 1974; **52**: 421–8.
- ¹⁵Bartholomew RS. Glaucoma in a South African black population: a population study on the Pondo tribe of South Africa. *S Afr Arch Ophthalmol* 1976; **3**: 135–50.