Postural change of IOP in normal persons and in patients with primary wide open-angle glaucoma and low-tension glaucoma

SHIGEO TSUKAHARA AND TAKAYA SASAKI

From the Departments of Ophthalmology, Yamanashi Medical College, Yamanashi, and the Suwa Red Cross Hospital, Nagano, Japan

SUMMARY The values of intraocular pressure (IOP) measurements by an Alcon pneumatic tonometer and a Goldmann applanation tonometer were found to be close, with a correlation coefficient of 0.92. Measured by the pneumatonograph the IOP after 30 minutes in the supine position was highest in normal persons, in patients with primary wide open-angle glaucoma and low-tension glaucoma. The greatest difference in IOP between subjects sitting and supine was observed in patients with low-tension glaucoma.

Many authors have discussed the change of IOP measured by various tonometers in the sitting and supine positions.¹⁻⁶ However, methodologically^{7,8} Schiøtz's tonometer presents problems in measuring IOP⁹ in subjects sitting down, and there is also the problem of scleral rigidity.¹⁰ The reported data showed that there was an average rise in IOP of about 1 to 2 mmHg in subjects lying down.¹¹⁻¹⁵ The best method is to employ an Alcon pneumatic tonometer for measuring IOP in subjects either sitting or lying down.¹⁶⁻¹⁸ It is very important in the treatment of glaucoma patients to know the postural change of IOP.

The purpose of this paper is to report the postural change of IOP in normal persons and in patients with primary wide open-angle glaucoma and low-tension glaucoma by measuring it with an Alcon pneumatic tonometer.

Materials and methods

The materials consisted of 20 normal persons (average age 67.6, SD 13.1 years), 20 patients with primary wide open-angle glaucoma (average age 60.8, SD 20.6 years), and five patients with low-tension glaucoma (average age 63.3, SD 16.8 years).

An Alcon pneumatic tonometer was used for both the sitting and the supine positions. A Goldmann

tonometer was used only for the sitting position. In 20 normal eyes the relationship between the IOP measurement with the Goldmann tonometer and the Alcon pneumatonograph (PTG) was examined in the sitting position. In all procedures one drop of 0.4% oxybuprocaine was instilled into both eyes. The IOP was measured in the right eye in the same room between 10 and 12 o'clock in the morning. No pillow was allowed. Glaucoma patients had not received treatment for 3 days prior to the test.

The order of the measurements is shown in Fig. 1. First, Goldmann applanation readings were taken with the subjects seated. Immediately after that IOP

Fig. 1 Order of measurements.

Correspondence to Professor S. Tsukahara, Department of Ophthalmology, Yamanashi Medical College, Tamaho, Yamanashi, Japan 409-38.

Fig. 2 Comparative IOP measurements using Goldmann applanation and Alcon pneumatic tonometer with patients seated. Number of eyes=45. Correlation coefficient=0.91. Regression equation $y=0.12+0.93x$. Goldmann applanation tonometry - PTG=0.91 (SD 1.53) mmHg.

was measured with PTG in the same position. The patients then lay flat on the bed without a pillow about 1 m away from the tonometer and were measured again with PTG. They were measured again with PTG after 30 minutes. Finally, the subjects sat in the chair in front of the slit-lamp and were measured again for IOP with PTG and the Goldmann tonometer.

The Alcon pneumatonometer was calibrated each day before use. The value of measurements with PTG was judged by the mean level of the recorded pulse wave. All measurements were made by the same examiner.

Results

Fig. 2 shows the correlation between Goldmann applanation and the PTG tonometer on subjects in the sitting position. The mean difference in IOP between the Goldmann applanation tonometer and the PTG tonometer was 2.8, SD 0.83 mmHg. The former tended to be higher than the latter. Both were in close agreement with a correlation coefficient of 0.92. In the controls the mean applanation IOP recorded in the sitting position with the PTG tonometer was 13.0, SD 1.86 mmHg, and 14.0, SD 2.14 mmHg with the Goldmann applanation tonometer (Table 1).

The IOP measured after 30 minutes in the supine position by PTG tonometry was highest in normal persons, lower in the patients with primary wide

open-angle glaucoma and with low-tension glaucoma (Tables 1, 2, 3).

The mean difference in IOP between the sitting and supine positions was 5.6, SD 1.71 mmHg in the normal persons, 6.5, SD 2.66 mmHg in the patients with primary wide open-angle glaucoma, and 8.6, SD 3.5 mmHg in the patients with low-tension glaucoma (Fig. 3). It should be pointed out that the most significant difference was observed in the patients with low-tension glaucoma.

Discussion

Many authors have reported on the correlation of IOP between the Goldmann and the Alcon PTG.¹⁸⁻²² Some of them used the Goldmann tonometer with subjects in the sitting position and the Schiötz tonometer for the supine position. In these experiments the Goldmann tonometer proved of more value than the PTG. Moreover, some of the workers observed that the rise in pressure when patients become supine is higher in glaucomatous eyes.^{11 12 16} We have confirmed this, and the IOP by the Alcon PTG was 1.5 mmHg greater than by the

Table 1 Normal eyes (20 eyes, average ages 67.6, SD 13.1 yr)

	IOP mmHg ($\pm$ SD)
Goldmann applanation tonometry (sitting)	14.0 $\pm$ 2.14
Pneumatograph (sitting)	13.0 $\pm$ 1.86
Pneumatograph (supine)	16.9 $\pm$ 2.63
Pneumatograph (30 min supine)	18.9 $\pm$ 2.65
Pneumatograph (sitting)	13.1 $\pm$ 2.52
Goldmann applanation tonometry (sitting)	13.8 $\pm$ 2.52

Table 2 Primary wide open angle glaucoma eyes (20 eyes, average ages 60.8, SD 20.6 yr)

	IOP mmHg ($\pm$ SD)
Goldmann applanation tonometry (sitting)	18.3 $\pm$ 3.42
Pneumatograph (sitting)	17.1 $\pm$ 4.00
Pneumatograph (supine)	20.3 $\pm$ 4.15
Pneumatograph (30 min supine)	23.6 $\pm$ 5.20
Pneumatograph (sitting)	18.6 $\pm$ 4.72
Goldmann applanation tonometry (sitting)	19.1 $\pm$ 4.44

Table 3 Low-tension glaucoma eyes (5 eyes, average ages 63.3, SD 16.8 yr)

	IOP mmHg ($\pm$ SD)
Goldmann applanation tonometry (sitting)	13.0 $\pm$ 1.67
Pneumatograph (sitting)	13.8 $\pm$ 2.40
Pneumatograph (supine)	19.8 $\pm$ 1.94
Pneumatograph (30 min supine)	22.4 $\pm$ 1.62
Pneumatograph (sitting)	15.0 $\pm$ 0.00
Goldmann applanation tonometry (sitting)	13.4 $\pm$ 1.50

Fig. 3 Histogram of pressure differential between patients sitting and lying down.

Goldmann tonometer in the same position. This difference is thought to be due to the correction of gain in the PTG tonometer.

At present it is generally accepted that measurements of IOP in the sitting and supine positions are best taken with the Alcon PTG. However, some problems still remain. Krieglstein *et al.*²² found a non-linear relationship between IOP increase and body position by measuring IOP in different body positions from 60° semiupright to 30° with the head down. The IOP postural response in individual subjects was roughly correlated with ophthalmic arterial pressure and with the episcleral venous pressure postural response. It is concluded that the IOP postural response depends on arterial and venous vascular changes when the subject moves from an erect to a horizontal position. The IOP measured in the supine position will be influenced by the use of a pillow and its thickness though pillows were not used in this study. Various reasons have been given to explain the rise of IOP in the recumbent position.

Weber and Price²³ found an average rise in the IOP of 6 mmHg in 177 eyes of subjects in the sitting and supine positions by measuring with PTG. They claimed that this increase in the diastolic ophthalmic arterial pressure would be transmitted to the intraocular blood vessels, causing them to dilate. They also indicated the possibility that it might be of importance in the treatment and early detection of glaucoma to demonstrate a larger postural IOP response in glaucomatous eyes. Obtaining a greater postural rise of IOP in a given eye might be another diagnostic factor in glaucoma.^{16 22 24}

The effect of placing the head below the body should be investigated in the future. A change of episcleral venous pressure would be expected in carotid cavernous fistula and mediastinal tumour, and in these diseases the postural change of IOP

should be examined. Finally, it has been suggested that damage to the optic nerve in low-tension glaucoma might occur when the patient is asleep in the supine position.

We have confirmed the fact that IOP measured with the body in different positions was greater in glaucomatous eyes than in normal eyes, and in patients with low-tension glaucoma the postural difference in IOP particularly was remarkable.

References

- 1 Linnér E, Rickenbach C, Werner H. Comparative measurements of the pressure in the aqueous veins and the conjunctival veins using different methods. *Acta Ophthalmol (Kbh)* 1950; **28**: 469-78.
- 2 Goldmann H, Schmidt Th. Über Applanations-tonometrie. *Ophthalmologica* 1957; **134**: 221-8.
- 3 Weekers R, Laverne G, Gougner L. The origin of functional deficiencies in the hypertensive eye. *Doc Ophthalmol* 1959; **13**: 251-60.
- 4 Drance SM. The coefficient of scleral rigidity in normal and glaucomatous eyes. *Arch Ophthalmol* 1960; **63**: 668-70.
- 5 Leydecker W. *Glaucoma: ein handbuch*. Berlin: Springer, 1960: 286.
- 6 Leith AB. Episcleral venous pressure in tonography. *Br J Ophthalmol* 1963; **47**: 271-8.
- 7 Langham ME, McCarthy E. A rapid pneumatic applanation tonometer: comparative findings and evaluation. *Arch Ophthalmol* 1968; **79**: 389-99.
- 8 Moses RA. Repeated applanation tonometry. *Ophthalmologica* 1961; **142**: 663-8.
- 9 Friedenwald JS. Tonometer calibration: an attempt to remove discrepancies found in the 1954 calibration scale for Schötz tonometers. *Trans Am Acad Ophthalmol Otolaryngol* 1957; **61**: 108-14.
- 10 Roberts W, Rogers W. Postural effects on pressure and ocular rigidity measurements. *Am J Ophthalmol* 1969; **57**: 111.
- 11 Anderson DR, Grant WM. The influence of position on IOP. *Invest Ophthalmol Visual Sci* 1973; **12**: 204-12.
- 12 Galin MA, McIlvor JW, Magruder GB. Influence of position on IOP. *Am J Ophthalmol* 1963; **55**: 720-3.
- 13 Tarkkanen A, Leikola J. Postural variations of the IOP as measured with the Mackay Marg tonometer. *Acta Ophthalmol* 1967; **45**: 569-75.

- 14 Tsamprlakakis J, Casey TA, Howell W, Edridge E. Dependence of IOP on induced hypotension and posture during surgical anaesthesia. *Trans Ophthalmol Soc UK* 1980; **100**: 521–6.
- 15 Vaz F, Gupta AK. The influence of posture on intraocular pressure. *Indian J Ophthalmol* 1981; **29**: 1–3.
- 16 Hetland-Eriksen J. On tonometry 5. The pressure of glaucomatous eyes measured in the sitting and the lying positions by means of the Goldmann applanation tonometer. *Acta Ophthalmol (Kbh)* 1966; **44**: 515–21.
- 17 Hetland-Eriksen J. On tonometry 6. Comparative tonometry with Goldmann applanation tonometer and the Schötz tonometer in the lying position. *Acta Ophthalmol (Kbh)* 1966; **44**: 522–38.
- 18 Jain JR, Marmion VJ. A clinical evaluation of the applanation pneumatonograph. *Br J Ophthalmol* 1976; **60**: 107–10.
- 19 Walker RE, Litovitz TL. An experimental and theoretical study of the pneumatic tonometer. *Exp Eye Res* 1972; **13**: 14–23.
- 20 Walker RE, Litovity TL. Pneumatic applanation tonometer studies. II. Rabbit cornea data. *Exp Eye Res* 1972; **13**: 187–93.
- 21 Walker RE, Langham ME. Pneumatic applanation tonometer studies. III. Analysis of floating tip sensor. *Exp Eye Res* 1975; **20**: 167–72.
- 22 Kriegelstein GK, Waller WK, Leydecker W. The vascular basis of the positional influence on the IOP. *Albrecht von Graefes Arch Klin Ophthalmol* 1978; **206**: 99–106.
- 23 Weber AK, Price K. Pressure differential of IOP measured between supine and sitting position. *Ann Ophthalmol* 1981; **13**: 323–6.
- 24 Wuthrich UW. Postural change and intraocular pressure in glaucomatous eyes. *Br J Ophthalmol* 1976; **60**: 111–14.