Correspondence

Are their spectacles really necessary?

SIR, This editorial,¹ which is overtly critical of opticians, unjustifiably so in my view, has been brought to my attention. Although I am writing in my private capacity as an experienced ophthalmic optician, I am also the Scottish representative of the British College of Ophthalmic Opticians and chairman of the National Optical Consultative Committee.

The main defect in your editorial lies in the assumption that '80–90% of the prescriptions' for the 516 wearers of supposedly unnecessary spectacles out of 13 800 school children aged 10 years or less 'have been issued by non-medically qualified persons.' You give no source for that figure of 80–90% but it presumably applies to all age groups in the population.

But the 'School Eye Service' is staffed almost entirely by medically qualified ophthalmic practitioners who very probably account for a much higher proportion than 20 to 10% of refractions in the age group 10 years or under.

I have no access to figures in the Bristol area where Dr Stewart-Brown's survey was done,² but I personally have ascertained information about all spectacle dispensing in the Lothian, Scotland, area in February 1985. That month was chosen because it was probably an average month and because after 1 April 1985 these figures will not be available. Of the 85 children under the age of 10 years who had spectacles dispensed 37 were seen by an ophthalmic optician and 48 by a medically qualified ophthalmologist. I suspect the same would be true of the Bristol area, in which case a high proportion of the unnecessary spectacles, if indeed they were unnecessary, were prescribed by ophthalmologists.

Both Dr Stewart-Brown's paper, and indeed your own editorial, admit that hypermetropia requiring spectacles can well exist in the presence of normal visual acuity, yet these cases may constitute a significant proportion of those who received unnecessary spectacles.

Your editorial directs attention to a possible tendency for spectacles to be prescribed for headaches. The belief is widespread among ophthalmologists as well as general medical practitioners that even minor refractive errors are a cause of headaches which can be cured by spectacles.

Dr Stewart-Brown admits (p 879) that a proportion of children who appear to have normal visual acuity may have been inadequately tested by a series of medical officers with variable training (i.e., non-specialists, presumably).

You are critical of a 'non-medically qualified person' converting 'what begins as a simple task, a refraction,' into a 'pseudo-medical event.' In rebuttal I would claim that the main contributor to the early diagnosis of glaucoma as well as cataract and other diseases (mainly ophthalmic but some general) in this country is the ophthalmic optician.

30 Elder Street, Edinburgh

JACK BROWN

References

- 1 Editorial: Are their spectacles really necessary? *Br J Ophthalmol* 1985; 69: 873.

- 2 Stewart-Brown S. Spectacle prescribing among 10-year-old children. *Br J Ophthalmol* 1985; 69: 874–80.

**The figure for '80–90% of the prescriptions' being issued by opticians was an estimate based on information issued by the Federation of Optical Corporate Bodies, 22 Nottingham Place, London W1, 'Optics at a Glance' 1984. This document states that 86.7% in 1983 were carried out by opticians and 67% of these tests resulted in a prescription for spectacles. Mr Brown appears to think that Dr Stewart-Brown's survey refers only to the Bristol area. However, if he consults the original paper again he will see that the children studied were all those born in the United Kingdom (excluding Northern Ireland) in the one week 7–11 April 1970.—Editor, *BJO*.

Transient monocular obscuration—amaurosis fugax?

SIR, I was very much interested in the paper 'Transient monocular obscuration—? amaurosis fugax: a case report' by Kosmorsky, Rosenfeld, and Burdel.¹ This was the case of a 73-year-old hypertensive man five years after right cataract extraction with iris fixation lens implant intermittently treated with pilocarpine to stabilise the lens. He presented repeated visual blurring the cause of which was found to be bleeding from abnormal superior angle vessels.

In 1972 we published the case of a 58-year-old hypertensive woman² who presented spontaneous hyphaema after topical administration of phenylephrine 10% for routine funduscopy followed by repeated pilocarpine 2% in order to speed the recontraction of the pupils. A blood clot was suspended at 12 o'clock of the pupillary border, but we could not observe any abnormal vessel in the area. We believed that perhaps a previously present abnormal vessel was pressed upon by the fibres of the pupillary sphincter in rapid contraction caused by pilocarpine and then ruptured, causing the bleeding; however, no evidence of the ruptured vessel could be found afterwards. Fechner³ described a case of spontaneous hyphaema with a ruptured microaneurysm at the pupillary margin and quoted a similar case described in 1932 by Tyson which was accompanied by vascular abnormality in one occipital lobe. Our patient,² who presented in 1976, has not had a recurrence of spontaneous hyphaema since then.

I am wondering if on the background of fragile new formed vessels near the iridectomy in the patient described by Kosmorsky *et al.* the intermittent administration of pilocarpine did not also play a role (perhaps by iris 'gymnastics' or by some pharmacological properties) in the appearance of hyphaema.

The differential diagnosis from carotid insufficiency was suggested by us in 1975 in a case of a 47-year-old woman⁴ with anomalous iris vessels in the right eye, right orbit cavernous haemangioma, and a seven-year period of repeated attacks of intermittent blurring of vision due to spontaneous hyphaema. After the excision of the orbital haemangioma in 1973 there have been no more attacks of spontaneous hyphaema up to the present.

Owing to the rarity of cases of spontaneous hyphaema I