there were children with optic atrophy, macular degeneration, retinoblastoma, and bilateral retinoblastoma who would not be expected to derive much benefit from spectacles; there were also ESN(S) children and severely emotionally disturbed children, for whom there may have been valid non-optical reasons for withholding spectacles. A detailed analysis of reasons for lack of spectacle prescription among all children in this group had not been undertaken when the paper was published, and I felt therefore it would be inappropriate to make a comment about this group. Some of these data have now been analysed and will be submitted for publication in a further paper.

The second group of children whom Drs Charman and Jennings comment upon are those with isolated near vision defects; only a small proportion of all the children in this group had been prescribed spectacles. I was not surprised to find a low prescription rate in this group, since screening for defective near vision is not common in schools,¹ and such children would therefore rarely be identified. This group was, however, the only one among children with defective vision in which an associated educational disadvantage that might be attributable to their untreated hypermetropia was detectable.² It is therefore a group which is worthy of comment and, indeed, further investigation.

Hypermetropia is very common in primary school children. It may be that these children would all benefit from spectacles, but it is important, before embarking upon such treatment policy, that a controlled trial be undertaken to establish exactly how spectacles could be of benefit. Such a study is currently being planned in the Department of Child Health Bristol and the Department of Ophthalmic Optics UWIST Cardiff.

S L STEWART-BROWN

Department of Child Health,
Bristol University

References

- 1 Stewart-Brown SL, Brewer R. The significance of minor defects of visual acuity in school children: implications for screening and treatment. *Trans Ophthalmol Soc UK* in press.
- 2 Stewart-Brown S, Haslum M, Butler NR. Educational attainment of children with visual defects. *Dev Med Child Neurol* 1985; 27: 498-503.

Screening for visual defects in preschool children

SIR, As part of the general ophthalmic service in Ayrshire and Arran Health Board primary orthoptic screening has been offered to all children at the age of 3½ since January 1978. A preliminary report of this has been published.¹ The methods used are often similar to those described by Ingram *et al.*,² but are always non-invasive and differ from those used by Ingram *et al.* in other ways. For example, the Sheridan Gardiner test (seven letters) is used for testing visual acuity, and the Frisby test is used for testing stereo acuity. Both those tests can give very accurate results and are eminently suitable for mass screening. Virtually no defects are missed and the false positive rate is approximately 1.5%. Unlike Ingram *et al.*, we consider an acceptable visual acuity to have to be at least 6/6.

We are now in the process of assessing the first five years of screening for both clinical effectiveness and cost effectiveness.

M CAMERON

Heathfield Hospital,
Heathfield Road,
Ayr KA8 9DZ

References

- 1 Cameron JH, Cameron M. Visual screening of preschool children. *Br Med J* 1978; ii: 1693-4.
- 2 Ingram RM, Walker C, Wilson JM, Arnold PE, Dally S. Prediction of amblyopia and squint by means of refraction at age 1 year. *Br J Ophthalmol* 1986; 70: 12-5.

SIR, I did know about the Camerons' paper in 1978, but overlooked reference to it, for which I apologise. I will await with interest publication of both their results of treating defective vision identified at 3½ and their methods of assessing 'cost effectiveness.' I still believe that a nationwide introduction of vision screening at 3½ should depend primarily on somebody being able to demonstrate that defective vision is more effectively treated if identified at 3½ than if it is left untreated until age 5+ (as at present).

I agree that the cost of any vision screening programme should also be considered, because we can no longer introduce a good idea just because it might be a 'good idea.'

R M INGRAM

Kettering and District
General Hospital,
Rothwell Road,
Kettering NN16 8UZ

Atracurium and intraocular pressure

SIR, Murphy *et al.*'s paper on intraocular pressure (IOP) has just come to hand.¹ Their claim that the effect of atracurium had 'not hitherto been documented' is incorrect, as my coworkers and I studied and published² the effects of atracurium on IOP in 1984. It was nevertheless pleasing to see the results of our study independently corroborated.

R J MAHARAJ

5th Floor,
Durdac Centre,
Smith Street,
Durban 4001,
South Africa

References

- 1 Murphy DF, Eustace P, Unwin A, Magner JB. Atracurium and intraocular pressure. *Br J Ophthalmol* 1985; 69: 673-5.
- 2 Maharaj RJ, Humphrey D, Kaplan N, *et al.* Effects of atracurium on intraocular pressure. *Br J Anaesth* 1984; 56: 459-63.

SIR, At the time of submission of our manuscript for publication there were no published reports on the effects of atracurium on intraocular pressure (IOP). Since that time two studies have been published apart from ours, that of Dr Maharaj *et al.*¹ and one by Tattersall, Manus, and Jackson.²

All three studies confirm the acceptability of atracurium for ophthalmic surgery.

In terms of purity of pharmacodynamic study, Dr Maharaj and colleagues' work is the most commendable. However, in the light of our study we feel that a more important fact should not be overlooked. The patient for whom changes in intraocular pressure are of most importance is the patient who has suffered a perforating eye injury. In this patient any change in IOP may cause further extrusion of intraocular contents and may critically jeopardise the eye. Our study, and that of Tattersall, Manus, and Jackson showed that, while atracurium is an acceptable drug for use in ophthalmic surgery, the considerable rise in IOP which results from laryngoscopy and tracheal intubation is of much greater significance. Until such time as we have an anaesthetic technique which is both rapid and unaccompanied by autonomic nervous overactivity, the perforated eye will remain a danger. Until this time, minor

differences such as those demonstrated between atracurium, pancuronium, and fazadinium pale into insignificance.

DERMOT MURPHY

Sir Charles Gairdner Hospital,
The Queen Elizabeth II Medical Centre,
Verdun Street,
Nedlands,
Western Australia 6009

References

- 1 Maharaj RJ, Humphrey D, Kaplan N, *et al.* Effects of atracurium on intraocular pressure. *Br J Anaesth* 1984; **56**: 459–63.
- 2 Tattersall MP, Manus NJ, Jackson DM. The effect of atracurium or fazadinium on intraocular pressure. A comparative study during induction of general anaesthesia. *Anaesthesia* 1985; **40**: 805–7.

Obituary

A G Palin, BM, BCL, FRCS, FRCSEd

Mr A G Palin, formerly consultant ophthalmologist at the Bristol Eye Hospital, died on 26 January 1986 after a long illness patiently borne.

Anthony Palin graduated from Oxford and St Thomas's Hospital Medical School in 1934. He was appointed consultant ophthalmologist at Bristol in 1937 and served in the Royal Air Force during the war. On his return to Bristol until his retirement in 1972 he played a major part in establishing the Eye Hospital as one of the best units outside London. For many of these years he was senior surgeon in the days when this position carried the onerous duties of

being chairman, head of the university department, and representative on many committees. Tony had no difficulty in carrying out these functions with aplomb, at the same time maintaining his reputation as a first class clinician. One mark of this distinction was the conferring of an honorary fellowship by the Royal College of Surgeons of England in 1970. There are many younger ophthalmologists, both in the UK and in Australia, who will remember his warm friendship and wise teaching.

Tony is survived by his wife Bobby, and two sons. NLD