Pink hypopyon: a sign of *Serratia marcescens* endophthalmitis

Selwa A F Al Hazzaa, Khalid F Tabbara, J Allan Gammon

Abstract

Endogenous bacterial endophthalmitis in infants is uncommon. We recently examined and treated an infant who presented with pink hypopyon which followed a *Serratia marcescens* septicaemia. Culture of the aspirate from the anterior chamber showed no red blood cells, and grew *Serratia marcescens*, which was also isolated from the tip of the child's umbilical artery catheter. The presence of a pink hypopyon in the absence of hyphaema may suggest the diagnosis of *Serratia marcescens* endophthalmitis.

(*Br J Ophthalmol* 1992; 76: 764-765)

Serratia marcescens, a Gram negative bacillus once believed to be a harmless saprophyte may cause infection in humans. This bacterium can cause meningitis, endocarditis, otitis media, sepsis, and ocular infections such as conjunctivitis and keratitis.¹⁻³ Endogenous bacterial endophthalmitis is uncommon but its occurrence in newborns has been reported.² We report a newborn infant who developed a pink hypopyon caused by *Serratia marcescens*.

Case report

After an uneventful pregnancy, a healthy 18-year-old primigravida had an uncomplicated labour and delivered a 2.65 kg full term infant. The neonate required resuscitation after delivery. Apgar scores were 1 at 5 minutes, 3 at 6 minutes, and 5 at 10 minutes. During the first few hours of life, the baby developed convulsions and was treated with 60% oxygen and phenobarbitone. Three days later the infant developed clinical signs of septicaemia and disseminated intravascular coagulopathy. *Serratia marcescens* was isolated from blood cultures and from the infant's ear, nose, and throat. *S. marcescens* was also cultured from the tip of the child's umbilical artery catheter.

Ophthalmic examination disclosed a normal right eye. The left cornea was hazy centrally. A fibrinous exudate was seen in the anterior chamber occluding the pupil and precluding visualisation of the fundus. A pink hypopyon occupied one third of the anterior chamber. Ultrasound evaluation of the posterior segment revealed dense, dispersed vitreous opacities and retinohoroidal thickening consistent with endophthalmitis. No intraocular calcification was noted.

An anterior chamber paracentesis was performed. Gram stain of the aqueous specimen revealed Gram negative bacilli and polymorphonuclear cells. Haematoxylin and eosin stain of the fibrinous exudate showed polymorpho-

nuclear leucocytes but not red blood cells. *Serratia marcescens* was isolated from the aqueous. Vitreous was aspirated for microscopy and culture. A 25 gauge needle on a tuberculin syringe was inserted through the pars plana at the 9 o'clock position and was advanced less than 1 cm to reach the mid vitreous. Liquefied vitreous (0.3 ml) was aspirated, and through the same site gentamicin 100 µg and cephazolin 100 µg were injected intravitreally. Culture of vitreous aspirate revealed no growth. The view of the fundus was not possible because of the hazy media.

Antibiotic sensitivity studies showed the organism to be susceptible to amikacin, chloramphenicol, carbenicillin, gentamicin, minocycline, and trimethoprim-sulphamethoxazole, and resistant to ampicillin, cephalosporins, erythromycin, polymyxin B, and tetracycline. Radiological examination of the extremities confirmed the presence of periosteal reaction suggestive of osteomyelitis. The patient responded to parenteral gentamicin therapy. The left eye showed progressive healing of the inflammatory reaction. Three months later the left eye started showing evidence of phthisis.

Discussion

Serratia marcescens was first described by Bizio in 1823.⁴ Subsequently, the bacterium became known as *Bacterium prodigiosum* because it produces a red pigment (prodigiosin). Its red pigment has been a useful marker in the study of bacteraemia after dental extractions⁵ and bacteriuria after urinary tract procedures.⁶ *S. marcescens* endophthalmitis has rarely been reported. A review of 72 cases of metastatic endophthalmitis in the past 10 years disclosed only three cases of *S. marcescens*, all in the adult age group.⁷⁻⁹

Endophthalmitis in infants has received little attention. Three cases of bacterial endophthalmitis reported in the neonatal age group were reported to have been caused by *Streptococcus pneumoniae*, *Streptococcus algalactiae*, and group B *Streptococcus*.¹⁰⁻¹²

Serratia marcescens ocular infections have been described as purulent conjunctivitis in six premature infants.¹³ In 1971 Bigger *et al* reported two cases of severe purulent endophthalmitis owing to *S. marcescens* in adults.¹⁴ We assume that our patient had metastatic endophthalmitis following septicaemia acquired by umbilical artery catheter infection.

Because of the potential severity of ocular involvement by this organism, a characteristic presentation of pink hypopyon may suggest *S. marcescens* and should prompt the ophthalmologist to look for laboratory confirmation

Department of
Ophthalmology, College
of Medicine, King Saud
University and the King
Khaled Eye Specialist
Hospital, Riyadh, Saudi
Arabia
S A F Al Hazzaa
K F Tabbara
J A Gammon

Correspondence to:
Dr K F Tabbara, Department
of Ophthalmology, College of
Medicine, King Saud
University, PO Box 55307,
Riyadh 11534, Saudi Arabia.

Accepted for publication
22 July 1992

and institute immediate management with appropriate antibiotics such as gentamicin.

Endogenous metastatic endophthalmitis in infants is rare. Vitrectomy and intraocular antibiotic injections may be indicated in such patients but are technically demanding if the view is obstructed, and may carry considerable risks.

- 1 Gammon JA, Schwab I, Joseph P. Gentamicin-resistant *Serratia marcescens* endophthalmitis. *Arch Ophthalmol* 1980; 98: 1221-3.
- 2 Lazarcheck GW, Boyle GL, Schwartz AL, Leopold IH. *Serratia marcescens*, an ocular pathogen. *Arch Ophthalmol* 1971; 85: 599-603.
- 3 Davis JT, Foltz E, Blackmore WS. *Serratia marcescens*: a pathogen of increasing clinical importance. *JAMA* 1970; 213: 2190-2.
- 4 Gaughran ER. From superstition to science. The history of a bacterium. *Trans NY Acad Sci* 1969; 31: 3-24.
- 5 McEntegart MG, Porterfield JS. Bacteremia following dental extractions. *Lancet* 1949; ii: 596-8.
- 6 Kass EH, Schneiderman LJ. Entry of bacteria into the urinary tracts of patients with inlaying catheters. *N Engl J Med* 1957; 256: 556-7.
- 7 Greenwald MJ, Wohl LG, Sell CH. Metastatic bacterial endophthalmitis: a contemporary reappraisal. *Surv Ophthalmol* 1986; 31: 81-101.
- 8 Malnou F. Metastatic septique endoculaire per *Serratia*. *Bull Soc Ophthalmol Fr* 1981; 81: 803-6.
- 9 Radda TM. Metastatic *Serratia marcescens* endophthalmitis. *Ophthalmologica* 1982; 185: 65-8.
- 10 Greene GR, Carroll WL, Morozumi PA. Endophthalmitis associated with group-B streptococcal meningitis in an infant. *Am J Dis Child* 1979; 133: 752.
- 11 Weintraub MI, Otto RN. Pneumococcal meningitis and endophthalmitis in a newborn. *JAMA* 1972; 219: 1763-4.
- 12 Berger B. Endophthalmitis complicating neonatal group B streptococcal septicemia. *Am J Ophthalmol* 1981; 92: 681-4.
- 13 Stenderup A, Faergeman O, Ingerslev M. *Serratia marcescens* infections in premature infants. *Acta Pathol Microbiol Scand* 1966; 68: 157-60.
- 14 Bigger JF, Meltzer G, Mandell A, Burde RM. *Serratia marcescens* endophthalmitis. *Am J Ophthalmol* 1971; 72: 1102-5.