- 9 Young R, Price J, Gorham N, Cowart M. Selective abnormality of the cone b-wave in a patient with retinal degeneration. *Doc Ophthalmol* 1985; 60: 211–8.
- 10 Wakabayashi K. Electrodiagnosis of primary macular dystrophies. *J Jusen Med Soc* 1986; 96: 399–439.
- 11 Karpe G. The basis of clinical electroretinography. *Acta Ophthalmol* 1945; 24 (suppl): 1–118.
- 12 Schubert G, Bornschein H. Beitrag zur Analyse des menschlichen Elektoretinogramms. *Ophthalmologica* 1952; 123: 396–413.
- 13 Miyake Y, Yagasaki K, Horiguchi M, Kawase Y, Kanda T. Congenital stationary night blindness with negative electroretinogram. A new classification. *Arch Ophthalmol* 1986; 104: 1013–20.
- 14 Miyake Y. Incomplete type congenital stationary night blindness. In: Heckenlively JR, Arden GB, eds. *Principles and practice of clinical electrophysiology of vision*. St Louis: Mosby, 1991: 721–5.
- 15 Weleber RG, Pillers DAM, Powell BR, Hanna CE, Magen RE, Buist NRM. Aland Island eye disease (Forsius-Eriksson syndrome) associated with contiguous deletion syndrome at Xp21. Similarity to incomplete congenital stationary night blindness. *Arch Ophthalmol* 1989; 107: 1170–9.
- 16 Young RSL. Low-frequency component of the photopic ERG in patients with x-linked congenital stationary night blindness. *Clin Vis Sci* 1991; 6: 309–15.
- 17 Hirose T, Wolf E, Hara A. Electrophysiological and psychophysical studies in congenital retinosis of x-linked recessive inheritance. *Doc Ophthalmol Proc Ser* 1977; 13: 173–84.
- 18 Kellner U, Brümmer S, Foerster MH, Wessing A. X-linked congenital retinosis. *Graefes Arch Clin Exp Ophthalmol* 1990; 228: 432–7.
- 19 Rhode N, Täumer R, Pernice D. Vorschlag eines verbesserten klinischen EOG-Testes. *Ber Dtsch Ophthalmol Ges* 1977; 74: 747–50.
- 20 Foerster MH, Kellner U, Wessing A. Cone dystrophy and supernormal dark-adapted b-waves in the electroretinogram. *Graefes Arch Clin Exp Ophthalmol* 1990; 228: 116–9.
- 21 Marmor MF, Arden GB, Nilsson SEG, Zrenner E. Standard for clinical electroretinography. *Arch Ophthalmol* 1989; 107: 816–9.
- 22 Granit R. *Sensory mechanisms of the retina*. London: Oxford University Press, 1947.
- 23 Bacon P, Spalton DJ, Smith SE. Blindness from quinine toxicity. *Br J Ophthalmol* 1988; 72: 219–24.
- 24 Krill AE. *Hereditary retinal and choroidal diseases*. Vol II. Hagerstown: Harper & Row, 1977: 403–4.
- 25 Smith EL, Harwerth RS, Crawford MLJ, Duncan GC. Contribution of the retinal ON channels to scotopic and photopic spectral sensitivity. *Vis Neurosci* 1989; 3: 225–39.
- 26 Condon GP, Brownstein S, Wang NS, Kearns AF, Ewing CC. Congenital hereditary (juvenile x-linked) retinosis. Histopathologic and ultrastructural findings in three eyes. *Arch Ophthalmol* 1986; 104: 576–83.
- 27 Birch DG. Flicker electroretinography. In: Heckenlively JR, Arden GB, eds. *Principles and practice of clinical electrophysiology of vision*. St Louis: Mosby, 1991: 348–51.

History of ophthalmology

On private practice

Those who sometimes feel overwhelmed by the financial and logistic complications of private practice, may safely direct their ire at Hammurabi, who seems to have started the whole business in 2500 BC by publishing the following table of fees:

- 'Setting of man's broken bone' – 5 Shekels
- 'Setting freeman's bone' – 3 Shekels
- 'Setting slave's bone' – 2 Shekels
- 'Treating ox or ass' – 1/6 Shekel

This should actually cheer up one's accountant, as today's fees aren't tailored to patients' income (whether the NHS soon will be is a moot point). And we don't treat oxen – in fact are legally prevented from doing so. Whether we ever treat asses is a matter of opinion.

Private practice has been a lucrative business for centuries. Pope Clement V had to forbid medical practice by ecclesiastical persons, as it was so financially rewarding that all the monks were leaving the monasteries to practise full time. There weren't many bishops left, either.

A little later, many Tudor doctors were able to purchase country estates with their earnings. But this situation becomes less desirable when the reason for such high fees is known – they had to reside in the patient's house 'until all suppuration had healed.' One such experience with a 'heart-sink' patient could undoubtedly break a practitioner.

During many periods of history, payment was only made for successful operations. Oph-

thalmologists in Babylon received no shekels at all (and occasionally had their hands hacked off) if the patients' sight did not return. This must have imbued the postoperative ward rounds with a certain tension. One wonders whether they had insurance for 'loss of income owing to disablement.'

Collection of the fee has long been fraught with difficulty. A Tudor surgeon advised that 'It is surer to be paid while the patient is not entirely well, as the outstretched hand may be drawn back after cure is complete.' This advice is hard to follow nowadays. For one thing, the ophthalmology private patient would be unable to write the cheque until the dressings came off.

The Tudors did have a system of legal redress, though. In 1537 Dr Andrew Boorde petitioned for the sum of £53, 'in money, cloth and two horses' which was owed to him. As the average annual wage was £2, Boorde could have hung up his instrument case and retired had he got it. Fortunately for medicine – as his later writings were excellent – he did not.

FIONA ROMAN

- Gordon BL. *Medieval and renaissance medicine*. London: Peter Owen, 1959: 759–61.
- Graham H. *Surgeons all*. London: Rich and Cowan, 1939: 33–8.
- Riesman D. *Medicine in the middle ages*. New York: Paul B Hoeber Inc, 1935: 17–27.
- Sigerist HE. *A history of medicine*. New York: Oxford University Press, 1951: 246–52.