- 1 Algawi K, Goggin M, O'Keefe M. Refractive outcome following diode laser versus cryotherapy for eyes with retinopathy of prematurity. *Br J Ophthalmol* 1994;78:612-4.
- 2 Ling CS, Fleck BW, Wright E, Anderson C, Laing I. Diode laser treatment for retinopathy of prematurity: structural and functional outcome. *Br J Ophthalmol* 1995;79:637-41.
- 3 Scharf I, Zonis S, Zeltzer M. Refraction in premature babies. *Metab Ophthalmol* 1978;2:395-6.
- 4 Fledelius HC. Pre-term delivery and the growth of the eye. An oculometric study of eye size around term-time. *Acta Ophthalmol* 1992;204(Suppl):10-5.
- 5 Fledelius HC. Myopia of prematurity—changes during adolescence. A longitudinal study including ultrasound. *Doc Ophthalmol Proc Series* 1981;29:217-23.
- 6 Fledelius HC. Changes in refraction and eye size during adolescence. With special reference to the influence of low birth weight. *Doc Ophthalmol Proc Series* 1981;28:63-9.
- 7 Fledelius HC. Pre-term delivery and subsequent ocular development. A 7-10 year follow-up of children screened 1982-84 for ROP. 3) Refraction. Myopia of prematurity. *Acta Ophthalmol Scand* 1996;74:297-300.
- 8 Nissenkorn I, Yassur Y, Mashkowski D, Sherf I, Ben Sira I. Myopia in premature babies with and without retinopathy of prematurity. *Br J Ophthalmol* 1983;67:170-3.
- 9 Quinn GE, Dobson V, Repka MX, Reynolds J, Kivlin J, Davis B, et al. Development of myopia in infants with birth weights less than 1251 grams. *Ophthalmology* 1992;99:329-40.
- 10 Gallo JE, Fagerholm P. Low-grade myopia in children with regressed retinopathy of prematurity. *Acta Ophthalmol* 1993;71:519-23.
- 11 Robinson R, O'Keefe M. Follow up study on premature infants with and without retinopathy of prematurity. *Br J Ophthalmol* 1993;77:91-4.
- 12 Fledelius HC. Pre-term delivery and subsequent ocular development. A 7-10 year follow-up of children screened 1982-84 for ROP. 4) Oculometric—and other metric considerations. *Acta Ophthalmol Scand* 1996;74:301-5.
- 13 Laws D, Shaw DE, Robinson J, Jones HS, Ng YK, Fielder AR. Retinopathy of prematurity: a prospective study. Review at six months. *Eye* 1992;6:477-83.
- 14 Cryotherapy for Retinopathy of Prematurity Cooperative Group. The natural outcome of premature birth and retinopathy. Status at one year. *Arch Ophthalmol* 1994;112:903-12.
- 15 Seiberth V, Knorz MC, Trinkmann R. Refractive errors after cryotherapy in retinopathy of prematurity. *Ophthalmologica* 1990;201:5-8.
- 16 Cryotherapy for Retinopathy of Prematurity Cooperative Group. Multi-center trial of cryotherapy for retinopathy of prematurity. 3½ year outcome—structure and function. *Arch Ophthalmol* 1993;111:339-44.
- 17 Pohlant F. Hypothesis: myopia of prematurity is caused by postnatal bone mineral deficiency. *Eur J Pediatr* 1994;153:234-6.
- 18 Gerding H, Busse H. Myopia of prematurity (MOP) is definitely not a consequence of skull deformation. *Eur J Pediatr* 1995;154:245-6.
- 19 Fielder AR, Levene MI, Russell-Eggitt IM, Weale RA. Temperature—a factor in ocular development? *Dev Med Child Neurol* 1986;28:279-84.
- 20 Stone RA, Lin T, Desai D, Capehart C. Photoperiod, early post-natal eye growth, and visual deprivation. *Vis Res* 1995;35:1195-202.
- 21 Fielder AR, Foreman N, Moseley MJ, Robinson J. Prematurity and visual development In: Simons K, ed. *Early visual development, normal and abnormal*. New York: Oxford University Press, 1993:485-504.
- 22 Billson FA, Fitzgerald BA, Provis JM. Visual deprivation in infancy and childhood: clinical aspects. *Aust NZ J Ophthalmol* 1985;13:279-86.
- 23 Von Noorden GK, Khodadoust A. Retinal hemorrhage in newborns and organic amblyopia. *Arch Ophthalmol* 1973;89:91-3.
- 24 Miller-Meeks MJ, Bennett SR, Keech RV, Blodi CF. Myopia induced by vitreous hemorrhage. *Am J Ophthalmol* 1990;109:199-203.
- 25 Lue C-L, Hansen RM, Reisner DS, Findl O, Petersen RA, Fulton AB. The course of myopia in children with mild retinopathy of prematurity. *Vis Res* 1995;35:1329-35.
- 26 Hittner HM, Rhodes LM, McPherson AR. Anterior segment abnormalities in cicatricial retinopathy of prematurity. *Ophthalmology* 1979;86:803-16.
- 27 Gordon RA, Donzis PB. Myopia associated with retinopathy of prematurity. *Ophthalmology* 1986;93:1593-8.
- 28 Kelly SP, Fielder AR. Microcornea associated with retinopathy of prematurity. *Br J Ophthalmol* 1987;71:201-3.
- 29 Laws DE, Haslett R, Ashby D, O'Brien C, Clark D. Axial length biometry in infants with retinopathy of prematurity. *Eye* 1994;8:427-30.
- 30 Nissenkorn I, Kremer I, Ben Sira I, Cohen S, Garner A. A clinicopathological case of retinopathy of prematurity (ROP) treated by peripheral cryopexy. *Br J Ophthalmol* 1984;68:36-41.
- 31 Benner JD, Huang M, Morse LS, Hjelmeland LM, Landers MB. Comparison of photocoagulation with the argon, krypton, and diode laser indirect ophthalmoscopes in rabbit eyes. *Ophthalmology* 1992;99:1554-63.
- 32 McNamara JA. Laser treatment for retinopathy of prematurity. *Curr Opin Ophthalmol* 1993;4:76-80.

Cataract and driving

Onset of cataract and its surgery have both practical and legal consequence for driving. In the UK the need to satisfy the number plate test is absolute in law (a standard number plate of 79.4 mm in height at 20.5 metres in good daylight). Empirically found to be equivalent to 6/10 Snellen,¹ the test is consistent with the European Union (EU) Directive (now law),² which requires a minimum acuity of 0.6 metric (= 6/10 Snellen), as well as a minimum field of vision of 120°.

The paper in this issue of the *BJO* by Mönestam and Wachtmeister (p 16), states a legal requirement for Sweden as 0.5 (6/12 Snellen) at the time of the study, though Sweden is now subject to the EU directive (and 0.6 acuity). This variation does not detract from the value of their paper which investigates driver patients' largely subjective estimation of their visual function while driving, as well as their actual visual acuities both before and after modern cataract surgery. It is to some extent a measure of their driver patients' needs from cataract surgery as well as their expectations. The latter are sometimes unrealistically high in UK experience. Their results show a commendably high standard of visual improvement for driving. In the light of driving safety the authors show the real value of pseudophakic cataract surgery, as well as the value of surgery for the second eye. They look specifically at the problems of estimating distance, which is of some importance in driving and is relevant to other research—for example, into 'estimated time to collision', and other psychophysical factors.³

Interestingly, their paper records 23% of patients before surgery who continued driving while below the national standard, a not unknown clinical finding in the UK. They

find little correlation between visual acuity and difficulty in driving before surgery; perhaps a reflection of the weak statistical correlation of the multifactorial visual factors to driving accidents.^{4,5}

The authors reasonably rely on much very subjective data, bearing in mind the subjective nature of visual perception itself. Perception is at least as important as vision in driving. New studies and methods to test drivers' visual perception are beginning to emerge—for instance, 'usable field of view' and driving simulation.^{6,7}

There are, of course, other factors to be addressed in the context of cataract surgery and driving—that is, the role of contrast, and also discomfort and disability glare in both bright daylight and from oncoming headlights.^{8,9} Glare in some of these circumstances can produce both optical flare (and disability glare) as well as diffraction effects from headlamps and street lamps, both before and after surgery. Brightness acuity and glare tests have been used to assess both discomfort and visual function for driving in glare conditions.^{10,11}

Personal investigation has shown glare and diffraction effects can be quite serious in early cortical and, especially, cuneiform cataracts. These indicate the value of the number plate test as a practical visual acuity and square wave contrast sensitivity test in the real highway environment.⁸ The same investigations have shown that diffraction lines occur with bright sources such as headlamps, street lighting, and surface water reflections. These are also significant after extracapsular surgery where there are capsule traction lines present. The angle of the diffracted flare line can be directly related to the traction line axis. Capsule pearls can produce a more diffuse flare and disability

glare.¹² These phenomena can reduce driver patients' subjective satisfaction after pseudophakic surgery. Even after effective YAG laser capsulotomy diffractive phenomena can still occur with pupil dilatation in dark conditions, especially where the capsulotomy has been made small for quite valid reasons. 'In the bag' implantation and continuous curvilinear capsulotomy within the normal pupil margin to some extent increase the possibility of these diffractive phenomena.

There are other factors which may reduce patient satisfaction with their cataract surgery, in particular altered visual space as a result of changed refractive error and, rarely, anisometropia. The latter is less common since the use of A scan biometry has become almost universal, though errors still occur. Equally, changed or added astigmatism, fairly common in extracapsular surgery, can alter perception of visual space and driving coordination. Newer small incision techniques, such as phacoemulsification, barely alter or even reduce astigmatism, and have gone a long way to overcome these sources of dissatisfaction.¹³ These postoperative alterations make it prudent that there should be a period of adaptation before driving after cataract surgery, even where the legal standards can be met. This period might be as short as a week with current surgical methods.

There are still some patients driving with aphakia (especially in the developing countries) though the problems of magnification, astigmatism, spherical and chromatic aberration, and the roving ring scotoma of non-aspheric aphakic glasses are true handicaps for driving. Some of these drivers barely reach or fail the visual field standard² and there is a natural reluctance of many ophthalmologists to jeopardise their patients' right to drive by too rigorous investigation.

The paper from Sweden shows that a few driver patients decide not to resume driving after surgery, but most improve greatly from modern extracapsular cataract extraction or phacoemulsification cataract surgery and appreciate the benefit of retaining that important civil liberty, the right to drive.

GREG MUNTON

Maidstone, Kent ME14 1HG

- 1 Drasdo M, Haggarty CM. *A comparison of British number plates and Snellen vision tests for car driving. Research report RF676.* Crowthorne, Berks: Transport and Road Research Laboratory, 1977.
- 2 Commission of the European Communities. Second Directive on Driving. 1991;439:EC.
- 3 Berthelon C, Mestre D. Visual cues for the detection of impending collision at crossroads. *Vision in Vehicles* 1993;IV:101-8.
- 4 Burg A, Hills BL. *Analysis of California drivers vision data. Report 768.* Crowthorne, Berks: Transport and Road Research Laboratory, 1977.
- 5 Davison PA. Inter relationships between British drivers visual disabilities, age and road accident histories. *Ophthalm Physiol Optics* 1985;5:195-204.
- 6 Dunne MC. Defective vision and road accidents. Pilot Study. Birmingham: University of Aston, 1995:18-19, 86, 94.
- 7 Ball K, Owsley C, Roenker DL, Bruni JR. Visual attention problems as a predictor of vehicle crashes in older drivers. *Invest Ophthalmol Vis Sci* 1993;34:31-40.
- 8 Neuman AC, McCary SR, Seedle TO, Sanders DR, Raanan MG. The relationship between indoor and outdoor Snellen visual acuity in cataract patients. *J Cataract Implant Surg* 1988;14:35-9.
- 9 Jaffe N. Glare and contrast: indications for cataract surgery. *J Cataract Implant Surg* 1986;12:372-5.
- 10 Hirsc RP, Nadler MP, Miller D. Glare measurements as a predictor of outdoor vision among cataract patients. *Ann Ophthalmol* 1984;16:965-8.
- 11 Holladay JT, Prager TC, Trujillo J, Ruiz RS. Brightness acuity test and outdoor visual acuity in cataract patients. *J Cataract Implant Surg* 1987;13:67-9.
- 12 Behrent S, Trier HD, Altenhar A, Hildenbran G. The effect of posterior chamber lenses or YAG laser capsulotomy on the sensitivity to glare and twilight vision in comparison to phakic control groups. *Klin Monatsbl Augenheilkd* 1988;193:249-56.
- 13 Pfeleger T, Skorpik C, Menapace R, Schotz U, Weghaupt H, Zehetmayer M. Long term course of induced astigmatism after clear corneal incision cataract surgery. *J Cataract Implant Surg* 1996;22:72-7.