Effects of scleral buckling and encircling procedures on human optic nerve head and retinochoroidal circulation

Miyuki Nagahara, Yasuhiro Tamaki, Makoto Araie, Shuichiro Eguchi

Abstract

Aims—To study the effects of segmental scleral buckling and encircling procedures on tissue circulation in the human optic nerve head (ONH) and choroid and retina.

Methods—Using the laser speckle method, the normalised blur (NB) value, a quantitative index of tissue blood velocity, was measured every 0.125 seconds and averaged over three pulses in the optic nerve head (NB_{ONH}) and choroid and retina (NB_{ch-ret}) in 10 patients with unilateral rhegmatogenous retinal detachment (mean age 52 (SD 17)). NB_{ONH} , NB_{ch-ret} , and intraocular pressure (IOP) in both eyes, and blood pressure (BP) were measured before, and 1, 4, and 12 weeks after the scleral buckling and encircling procedure. **Results**— NB_{ch-ret} on the buckled side was significantly reduced after surgery and smaller than that in the unoperated contralateral eye throughout the study period (ANOVA, $p < 0.0001$). NB_{ch-ret} on the unbuckled side, in the foveal area, NB_{ONH} , IOP, and BP showed no significant change.

Conclusions—It was indicated that the segmental scleral buckling procedure with encircling elements decreased tissue blood velocity in the choroid and retina on the buckled side but caused no significant change on tissue circulation in other areas of the fundus or ONH.

(Br J Ophthalmol 2000;84:31–36)

Scleral buckling procedures are the treatment of choice for the majority of uncomplicated rhegmatogenous retinal detachments. Postoperative complications related to alterations in blood supply, such as anterior and posterior segment ischaemia which usually severely influence visual recovery, are rare but have been reported by several authors.^{1–4} It may be possible that subclinical ischaemia after retinal detachment surgery exists in many more cases than reported.^{5–11} Thus, the effects of scleral buckling procedures on ocular circulation, if they exist, have important clinical implications.

Studies in which the laser Doppler technique was used indicated that the pulsatile characteristics of the retinal arterial blood flow measured in the treated eyes were altered,⁵ and that arterial flow rates in the major temporal retinal arteries were, on average, 50% lower in the treated eyes than in the unoperated contralateral eyes; removal of scleral buckling

elements in two patients produced increases of 73% and 44% in arterial blood flow rates.⁶ One study using colour Doppler imaging indicated that scleral buckling procedures with encircling elements decreased average blood flow velocity in the central retinal artery by 35% on the first operative day and by 50% 1 week postoperatively but did not affect that in the ophthalmic artery,⁷ whereas another study reported that the same surgery reduced the average blood flow velocity in the ophthalmic artery by 31% 2 days after surgery.⁸ In patients with rhegmatogenous retinal detachment, the ocular pulse amplitude in the surgically treated eyes was reportedly lower by 53% compared with fellow eyes 6 months after scleral buckling.¹¹ These findings suggest that scleral buckling decreases not only retinal but also choroidal and/or optic nerve head (ONH) circulation because the ophthalmic artery is the origin of the arterial branches that supply blood to the eye. To date, however, there have been no reports on the effect of the scleral buckling procedure on tissue circulation in different parts of the human fundus—that is, in the ONH, choroid in the foveal area, and choroid and retina on the buckled and unbuckled side.

We recently constructed an apparatus for non-contact, two dimensional estimation of ocular fundus tissue circulation, utilising the near infrared diode laser and the laser speckle phenomenon. The normalised blur (NB) value, a quantitative index of the tissue blood velocity, is measured non-invasively in a portion of the ONH, choroid, and retina with reasonable reproducibility.^{12–15} Using this laser speckle method, we studied the effects of the scleral buckling procedure with encircling elements on tissue circulation in the ONH, the choroid in the foveal area, and the choroid and retina on the buckled and unbuckled side in patients with rhegmatogenous retinal detachment before and after primary scleral buckling.

Materials and methods

SUBJECTS

From February 1996 to July 1996, 12 patients (mean age 52 years; range 21–7; four males and eight females) with unilateral, localised, quadrant rhegmatogenous retinal detachment repaired by uncomplicated scleral buckling and encircling procedures were consecutively enrolled. The entry criteria for the study were retinal detachment localised to the periphery of the fundus and not involving the macula, no other known retinal vascular disease, visual acuity correctable to at least 0.4, adequate

Department of
Ophthalmology,
University of Tokyo
School of Medicine
M Nagahara
M Araie
S Eguchi

Eye Clinic, Tokyo
Metropolitan Geriatric
Hospital
Y Tamaki

Correspondence to:
Makoto Araie, MD,
Department of
Ophthalmology, University
of Tokyo School of Medicine,
7-3-1, Hongo, Bunkyo-ku,
Tokyo 113, Japan

Accepted for publication
11 August 1999

pupillary dilatation, and clear media. Eyes which would need exophts crossing the mid-line were not included in the study. The protocol of the study was approved by the ethics review committee of the University of Tokyo School of Medicine and written informed consent was obtained from each participant after full explanation of the procedure. No patient had any known haematological disorder. The primary surgeon (SE) was notified of the patient's enrolment into the study after the surgery.

Surgery was performed under local anaesthesia for all 12 patients by one of the authors (SE). Retinal breaks were identified in all patients and were treated with transscleral cryotherapy. A Mira No 240 silicone encircling element (Mira, Inc, Waltham, MA, USA), in conjunction with a silicone tyre exopht, was used in all patients. The encircling element was secured to the sclera 13 mm posterior to the limbus with 5-0 Surgiron mattress sutures placed in the four quadrants between the muscles and fastened with a Mira No 270 sleeve with moderate tightness. The extent of the exopht was from 1.5 to 2.5 quadrants and was consistent with the size and location of retinal breaks and co-existing lattice degeneration. Subretinal fluid was drained externally in every patient. Extraocular muscle disinsertion was not required for any patient. There were no intraoperative complications. At the end of the procedure, indirect ophthalmoscopy revealed retinal reattachment along with the absence of central retinal artery pulsations in all patients. In none of the patients was the eye believed to be excessively firm at the end of the operation. The height of the scleral buckle also was roughly judged at this point by the angle of bending the retinal vessel onto the segmental buckle by the primary surgeon (SE); about 30° to 60° designated as medium and 60° to 90° as high. (designated as low, medium, or high). The type and location of each exopht was drawn in the chart after the procedure.

INSTRUMENT

A fundus camera was equipped with a diode laser (wavelength 808 nm) and image sensor (100 × 100 pixels). The fundus where the laser beam was focused was observed by means of an infrared CCD camera. The scattered laser light was imaged on an image sensor corresponding to a field of 1.06 × 1.06 mm (45 degree visual angle of the fundus camera) in the human fundus, on which a speckle pattern appeared. The difference between the average of the speckle intensity (I_{mean}) and the speckle intensity for successive scanings of the image speckles at the pixels on the sensor plane was calculated, and the ratio of I_{mean} to this difference was defined as NB.¹²⁻¹⁴ NB is nearly equivalent to the reciprocal of the speckle contrast defined by Fercher and Briers.¹⁶⁻¹⁷ The NB was calculated using a logic board every 0.125 seconds successively for a maximum of 7 seconds and divided into 50 colour coded levels, which were displayed as colour graphics on a colour monitor showing the two dimensional variation of the NB level over the field of

interest.¹⁸ The average NB level (NB_{av}) in any rectangular field of interest on a displayed colour map can be calculated and the change of NB_{av} over 7 seconds can be monitored at the same time. The maximum exposure of the retina with the present apparatus is below the permissible variable limits of American National Standard Institute (ANSI).¹⁸

NB MEASUREMENT IN HUMAN ONH AND CHOROID RETINA

The pupil was dilated with one drop of Mydrin M (0.4% tropicamide, Santen Pharmaceutical, Osaka, Japan) 30 minutes before measurements were taken. During the measurement, the subject was asked to watch a target light; an electrocardiogram was monitored simultaneously. The image speckles from the measurement field located in the temporal site of the ONH, corresponding to a field of 0.72 × 0.72 mm (30 degree visual angle of the fundus camera), were recorded. The average NB across the largest rectangular field in the measurement field free of visible surface vessels further averaged over three cardiac pulses when fixation was satisfactory, was calculated to obtain mean NB_{av} from the ONH (NB_{ONH}). The size of this rectangular field varied among subjects to avoid visible surface vessels, ranging from 0.15 × 0.22 mm (20 × 30 pixels) to 0.37 × 0.48 mm (50 × 65 pixels).

Image speckles from a field corresponding to a field of 1.06 × 1.06 mm (45 degree visual angle of the fundus camera) were recorded and the average NB across the largest rectangular field free of visible surface retinal vessels was averaged as described above to obtain the mean NB_{av} from the choroid and retina ($NB_{\text{ch-ret}}$). Although large choroidal vessels can be seen on the display, their contours were not as discrete as those of surface retinal vessels and the density of choroidal vessels are greater than surface visible retinal vessels. Therefore, we did not avoid visible choroidal vessels, while we did avoid visible retinal vessels. The size of this rectangular field varied among subjects to avoid surface visible retinal vessels, ranging from 0.48 × 0.53 mm (45 × 50 pixels) to 0.64 × 0.80 mm (60 × 75 pixels). Image speckles from the macula, corresponding to a field of 1.06 × 1.06 mm (45 degree visual angle of the fundus camera), were recorded and the average NB across the foveal area free of retinal vessels was averaged as above to obtain the mean NB_{av} from the foveal area (NB_{fovea}). The size of this square field was 0.32 × 0.32 mm (30 × 30 pixels).

The movement of the subject's eye in any direction during the measurement period was checked by the method previously described.¹⁴ It was also checked by inspecting the colour map and the time course plot of NB_{av} taken every 0.125 seconds; when there was no eye movement during measurement, visible surface vessels did not change position on the colour map and the time course plot of NB_{av} exhibited periodic fluctuations synchronised with cardiac pulse.¹⁸ On the other hand, when eye movement occurred, visible surface vessels in the colour map changed position according

Figure 1 Measurement field of normalised blur (NB) in human choroid and retina and optic nerve head (ONH). Image speckles from a field on the buckled side (A) and unbuckled side (B) with no discrete retinal vessels visible in the operated eye and those on the mirror image sides of the unoperated contralateral eye were recorded to measure NB in the choroid and retina. Image speckles from fovea and a field in temporal ONH, free of visible surface vessels, were recorded to measure NB in the fovea (C) and that in ONH tissue (D).

to the eye movement. The coefficients of reproducibility for 5 minute and 24 hour interval measurements in normal humans were 11.7% and 13.0% for NB_{ONH} and 8.7% and 9.7% for NB_{ch-ret} respectively.

Image speckles from all subjects were digitally stored as colour maps. NB_w was calculated by a blinded investigator (MN) for each subject in the same position and size of the rectangular field free of visible surface vessels by referring to a polaroid fundus picture and colour map.

STUDY PROTOCOL

The both eyes of a patient underwent identical assessment at each time point of measurement. The baseline measurements were carried out between 1100 and 1300 one day before surgery. After the pupil was dilated by one drop of Mydrin M, the NB_{ONH} and NB_{fovea} in both eyes were measured. The NB_{ch-ret} was measured in the eye with the retinal detachment in an area where the retina was not detached, outside the vascular arcade, and close to the area where the buckle was going to be located (the buckled side), and an area 180° from the buckled side (unbuckled side) (Fig 1). Thereafter, the NB_{ch-ret} on the mirror image side of the unoperated contralateral eye in relation to the buckled or unbuckled side of the operated eye was measured. Polaroid fundus photographs were

taken to record the sites of measurement. The IOP in both eyes and brachial arterial blood pressure were also measured. It took 5–10 minutes to complete all procedures in one eye.

During the prospective follow up period of 12 weeks, patients were examined 1, 4, and 12 weeks after the surgery and at each visit, brachial arterial blood pressure, and the IOP, NB_{ONH} , NB_{fovea} , and NB_{ch-ret} in both eyes were measured as described above. These NB measurements were performed in the same area as measured before surgery, referring to a fundus photograph and colour map of NB obtained before surgery to determine the preoperative locations. All measurements were performed between 1100 and 1300. The ocular perfusion pressure (OPP) was also calculated by:

$$OPP = \frac{2}{3}BP_m - IOP \tag{1}$$

where BP_m was mean brachial arterial blood pressure. All patients were monitored for evidence of anterior segment ischaemia or other postoperative complications.

DATA ANALYSIS

Statistical significance was determined with Student's *t* test for paired variates. Analysis of variance (ANOVA) for the data obtained by sequential measurements was also used.

Results

Two of the 12 patients, who could not visit the hospital on the scheduled postoperative day, were excluded from the study and the following analysis was performed in 10 patients (Table 1). In both the operated and unoperated contralateral eyes, there was no significant change in IOP from the baseline and no bilateral differences at any of the time points (paired *t* test, *p* > 0.1). There was no significant change from the baseline at any time point in mean blood pressure or OPP (paired *t* test, *p* > 0.1; Table 2).

There was no significant difference in the baseline NB_{ch-ret} , NB_{fovea} , and NB_{ONH} at any of the corresponding sites between the operated and the unoperated contralateral eyes (Table 3). There was a significant approximately 40% decrease from the baseline in the NB_{ch-ret} on the buckled side of the operated eye (Table 4), while no significant change was observed in NB_{ch-ret} on the mirror image side of the unoperated contralateral eye. Further, no significant change was observed in the NB_{ONH} and NB_{fovea} in both eyes, and NB_{ch-ret} on the unbuckled side of the operated eye and that on the mirror

Table 1 Patient characteristics

Patient	Age (years)/sex	Eye	Visual acuity				Explant type*
			Pre	1 week	4 week	12 week	
1 AY	62/M	R	0.8	0.5	1.0	0.8	240 287 (M)
2 SH	21/M	R	1.2	0.4	0.6	1.5	240 287 (H)
3 CY	75/M	R	0.7	0.5	0.8	0.8	240 287 (M)
4 TS	68/M	R	1.2	1.0	1.2	1.2	240 287 (H)
5 MO	46/M	R	1.5	1.0	1.0	1.5	240 220 (H)
6 ST	53/M	R	0.4	0.5	0.9	1.2	240 220 (H)
7 ME	23/F	R	0.8	0.8	0.8	1.0	240 287 (H)
8 WY	63/F	L	1.5	0.7	1.0	1.0	240 219 (M)
9 SK	41/M	R	1.5	0.9	2.0	2.0	240 220 (H)
10 OT	49/M	L	1.2	1.2	1.5	1.5	240 287 (H)

*Mira silicone element number (buckle height: H=high, M=medium).

Table 2 Measurement of variables in patients with retinal detachment

	Before	1 week	4 weeks	12 weeks
Intraocular pressure (mm Hg) (operated eye)	13.4 (4.7)	13.8 (3.8)	13.8 (3.8)	13.1 (3.5)
Intraocular pressure (mm Hg) (unoperated contralateral eye)	12.9 (3.8)	14.1 (4.2)	13.8 (3.6)	12.6 (2.9)
Mean blood pressure (mm Hg)†	96 (18)	90 (15)	88 (15)	88 (15)
Ocular perfusion pressure (mm Hg)‡ (operated eye)	51 (15)	46 (9)	45 (13)	46 (9)
Ocular perfusion pressure (mm Hg)‡ (unoperated contralateral eye)	51 (12)	46 (12)	45 (9)	46 (9)

Figures are mean (SD) in 10 patients. †Diastolic blood pressure + 1/3 (systolic blood pressure—diastolic blood pressure), ‡2/3 (mean blood pressure)—intraocular pressure. There were no significant intergroup differences observed in any of the variables. There was no significant difference observed between the initial measurement and subsequent measurements in any of the variables.

Table 3 Baseline value in the NB_{ch-ret}, NB_{fovea} and NB_{ONH} in patients

	Operated eye	Unoperated contralateral eye
NB _{ch-ret} (buckled side)*	25.1 (8.8)	23.9 (6.5)
NB _{ch-ret} (unbuckled side)*	22.0 (6.6)	22.1 (3.6)
NB _{fovea}	28.2 (5.8)	27.5 (6.5)
NB _{ONH}	23.6 (4.4)	25.2 (7.6)

Figures are mean (SD) in 10 patients. NB_{ch-ret}, NB_{fovea} and NB_{ONH} represent mean tissue blood velocity in the largest rectangular field in the measurement field free of visible surface vessels in choroid and retina, foveal area, and optic nerve head, respectively.

*Mirror image side of the buckled (unbuckled) side in the unoperated contralateral eye.

No significant bilateral difference is observed in any of variables (paired *t* test, *p*>0.1).

Table 4 Change in NB_{ch-ret}, NB_{fovea} and NB_{ONH}

	Before	1 week	4 weeks	12 weeks
Operated eye				
Δ NB _{ch-ret} (buckled side)	0	-7.86 (13.1)*	-8.95 (10.6)*	-12.8 (8.70)*
Δ NB _{ch-ret} (unbuckled side)	0	0.53 (7.47)	-1.66 (5.81)	-1.66 (8.05)
Δ NB _{fovea}	0	-0.95 (4.33)	-2.73 (5.74)	-2.47 (5.89)
Δ NB _{ONH}	0	-0.14 (3.15)	-0.10 (3.24)	-1.54 (3.12)
Unoperated contralateral eye				
Δ NB _{ch-ret} (buckled side)†	0	1.06 (6.41)	-0.03 (5.88)	1.87 (4.75)
Δ NB _{ch-ret} (unbuckled side)†	0	0.19 (3.66)	-0.24 (4.62)	-0.40 (1.72)
Δ NB _{fovea}	0	0.93 (4.21)	-0.13 (3.10)	0.53 (4.59)
Δ NB _{ONH}	0	-0.46 (2.92)	0.50 (3.45)	-0.71 (3.15)

Figures are mean (SD) in 10 patients. NB_{ch-ret}, NB_{fovea} and NB_{ONH} represent mean tissue blood velocity in the largest rectangular field in the measurement field free of visible surface vessels in choroid and retina, foveal area, and optic nerve head, respectively. ΔNB_{ch-ret}, ΔNB_{fovea} and ΔNB_{ONH} indicate difference in NB_{ch-ret}, NB_{fovea} and NB_{ONH} between before and each time point of measurement, respectively.

Bilateral difference is significant for ΔNB_{ch-ret} in buckled side (*p*<0.0001, ANOVA), but not for in unbuckled side, ΔNB_{fovea} and ΔNB_{ONH}.

*Significantly smaller than zero (*p*=0.0003–0.040).

†Mirror image side of the buckled (unbuckled) side in the unoperated contralateral eye.

image side of the unoperated contralateral eye (Table 4). There were significant bilateral differences in the change in the NB_{ch-ret} from the baseline (ΔNB_{ch-ret}) on the buckled side (ANOVA of repeated measurements, *p*<0.0001), but not the unbuckled side (Table 4). The bilateral difference was not significant for the change in both NB_{fovea} and NB_{ONH} from the baseline (ΔNB_{fovea} and ΔNB_{ONH}, respectively; Table 4).

Discussion

METHOD

NB is primarily a quantitative index of tissue blood velocity.^{12–14} In rabbits, when the ocular blood flow rate was artificially changed by increasing the IOP^{12–14} or by intravenous injection of a calcium antagonist,¹⁵ averaged NB levels over the sampling field, NB_{av}, had a good correlation with the blood flow rate determined using the microsphere method in the retina or choroid.^{12–14–15} These findings suggest that the NB also correlates with the blood flow rate in the choroid or retina, at least under certain conditions. According to Koelle *et al*,¹⁹ the penetration depth of near infrared laser (wavelength 811 nm) in the cat optic nerve exceeds 1 mm. When ONH blood flow rate in rabbits was artificially changed by a small amount of

endothelin-1 or by inhalation of 10% carbon dioxide, NB_{av} obtained from ONH tissue strongly correlated with the blood flow rate in the ONH determined using the hydrogen gas clearance method where a hydrogen electrode was inserted into the ONH tissue to a depth of approximately 0.7 mm.¹³ Although blood flow rate measured by the hydrogen gas clearance method does not directly reflect the movement of red blood cells as the NB_{av} measured by the laser speckle method, these findings suggest that the NB correlates with the blood flow rate also in the ONH, at least under certain conditions.

Unlike rabbit eyes, in which retinal vasculature exists only in the medullary wing area, the NB obtained from the fundus in human eyes is affected by both retinal and choroidal circulation.²⁰ In the case of Dutch rabbits, no significant difference was observed between the NB_{av} obtained using the diode laser from the medullary field where no discrete vessels were observed and the NB_{av} from the extramedullary field, within an IOP range between 10 and 80 mm Hg.¹² This finding suggests that the contribution of retinal microcirculation to the NB obtained from the fundus using the diode laser is very minor in rabbits. Measurement of NB_{av} obtained from the posterior fundus of monkeys using the diode laser before and after obstruction of a retinal artery (experimental branch retinal artery occlusion) suggested that the contribution of retinal microcirculation to NB_{av} obtained with the diode laser was about 20% or less in monkeys.²¹ These results suggest that the contribution of choroidal circulation to the NB_{av} obtained in the present study from the human fundus with no discrete retinal vessels visible is probably much greater than the contribution of retinal microcirculation. The NB obtained from the foveal area is supposed to be affected only by choroidal circulation because the area is free of retinal capillaries. It must be noted that the laser speckle method does not give an absolute value for the tissue blood velocity, but a relative value, because NB is also influenced by the reflectivity of the laser beam in the tissue. Therefore, the NB_{av} obtained from the ONH, foveal area, or mid-peripheral fundus should not be directly compared and may not be suited for interindividual comparison. On the other hand, this method yields good reproducibility in the same tissue of an individual,¹⁸ and its change relative to the baseline correlates well with the relative change of blood flow determined using the microsphere method^{12–14–15–22} or hydrogen gas clearance method.¹³ These characteristics and the

simplicity of the measurements makes this method very useful in following the time course of change in an individual tissue.

EFFECT OF SCLERAL BUCKLING PROCEDURE

The present study indicated that segmental scleral buckling procedures with encircling elements significantly decreased NB_{ch-ret} on the buckled side. Using bidirectional laser Doppler in conjunction with retinal photography, Ogasawara *et al.*⁶ reported that in the patients who had undergone uncomplicated scleral buckling procedures the blood flow rates through the major temporal retinal arteries were on average 50% lower in the surgically treated eyes than in the unoperated contralateral eyes. If we assume that the contribution of retinal microcirculation to the NB_{ch-ret} obtained in the present study is approximately 20% according to the result reported by Isono,²¹ it seems difficult to explain a 40% decrease in the NB_{ch-ret} on the buckled side after scleral buckling only by its effect on retinal circulation. A decrease in choroidal tissue blood velocity after the scleral buckling procedure seems very likely. This finding is consistent with previous reports using the microsphere²³ or hydrogen gas clearance method²⁴ in rabbits that choroid and retina blood flow buckled with encircling elements was significantly decreased²³ or that choroidal blood flow was reduced at the posterior and peripheral sides of the encircling band.²⁴ Ohkubo²⁵ found signs of reduced retinal perfusion in regions of reattached retina that were indented by scleral buckling procedures. Compressional mechanisms have been cited as the cause of reduced choroidal blood flow following scleral buckling procedures.^{9, 23, 24} Thus, the reduced blood velocity in the choroid was probably attributed to direct obstruction of the choroidal venous drainage due to scleral buckling and/or local effects of cryotherapy.

The important finding in the present study is that the NB_{ch-ret} on the unbuckled side, the NB_{fovea} , and NB_{ONH} showed no significant change in spite of the significant decrease in NB_{ch-ret} on the buckled side. This result may not be consistent with those of previous reports suggesting substantial decrease in the posterior segment circulation on the whole at various intervals after buckling procedure⁵⁻¹¹—that is, 50% decrease in the flow rates in the major temporal retinal arteries,⁶ 35%–50% decrease in flow velocities in the central retinal artery,⁷ 31% decrease in the flow velocity in the ophthalmic artery,⁸ and 53% decrease in the ocular pulse.¹¹ These previous reports, however, did not study the circulatory change in different parts of the fundus. Findings of the present study suggested that the segmental scleral buckling procedure with encircling elements does not cause clinically significant circulatory changes elsewhere on the buckled side, including the ONH and the foveal area where postoperative vision might be affected. The discrepancy between the present result and previous reports might be partly due to that the reduction of blood flow rate depends

upon the degree of constriction of the encircling band, and/or the extent of the exoplane as reported by Mano²⁴ or Diddie *et al.*²³ In the present study, the encircling element was fastened with a moderate tightness and the extent of the exoplane was 1.5 or 2.5 quadrants. If the encircling element was fastened more tightly and/or the exoplane was more than three quadrants, it may be possible that not only the NB_{ch-ret} on the unbuckled side, but also NB_{fovea} and/or NB_{ONH} are also affected significantly. In all of the present subjects, retinal detachment was successfully treated without any complications and it seems very unlikely that buckling and/or fastening of the encircling element were unreasonably mild. As discussed above, the present NB_{ch-ret} is thought to mainly reflect choroidal circulation, and partly retinal circulation. Therefore, the present result does not exclude the possibility that the retinal blood flow was slightly decreased also on the unbuckled side, as Ogasawara *et al.*⁶ reported, while the NB_{ch-ret} showed no significant change. Whether the NB_{fovea} or NB_{ONH} was also unaltered after segmental scleral buckling and encircling procedure in patients with pre-existing compromise of ocular tissue circulation awaits future study.

In summary, the present study demonstrated in otherwise healthy human eyes that the segmental scleral buckling procedure with encircling elements fastened with a moderate tightness causes a decrease in the blood velocity only in the choroid and/or retina on the buckled side, but little change in other areas of the fundus, including the ONH, foveal area, and fundus on the unbuckled side.

This work was supported in part by grant in aid for developmental scientific research (B) No 01870074 from the Ministry of Education, Science, Sports, and Culture of Japan.

- 1 Boniuk M, Zimmerman LE. Necrosis of uvea, sclera, and retina following operations for retinal detachment. *Arch Ophthalmol* 1961;66:318–26.
- 2 Foulds WS, Reid H, Chrisholm IA. Factors influencing visual recovery after retinal detachment surgery. *Mod Probl Ophthalmol* 1974;12:49–57.
- 3 Jarret WH, Brockhurst RJ. Unexplained blindness and optic atrophy following retinal detachment surgery. *Arch Ophthalmol* 1965;73:782–91.
- 4 Robertson DM. Anterior segment ischemia after segmental episcleral buckling and cryopexy. *Am J Ophthalmol* 1975;79:871–4.
- 5 Yoshida A, Fekke GT, Green GT, *et al.* Retinal circulatory changes after scleral buckling procedures. *Am J Ophthalmol* 1983;95:182–8.
- 6 Ogasawara H, Fekke GT, Yoshida A, *et al.* Retinal blood flow alterations associated with scleral buckling and encircling procedures. *Br J Ophthalmol* 1992;76:275–9.
- 7 Regillo CD, Sergott RC, Brown GC. Successful scleral buckling procedures decrease central retinal artery blood flow velocity. *Ophthalmology* 1993;100:1044–9.
- 8 Santos L, Capeans C, Gonzalez F, *et al.* Ocular blood flow velocity reduction after buckling surgery. *Graefes Arch Clin Exp Ophthalmol* 1994;32:666–9.
- 9 Dobbie JG. Circulatory changes in the eye associated with retinal detachment and its repair. *Trans Am Ophthalmol Soc* 1980;78:503–66.
- 10 Johnson MW, Han DP, Hoffman KE. The effect of scleral buckling on ocular rigidity. *Ophthalmology* 1990;97:190–5.
- 11 Yoshida A, Hirokawa H, Ishiko S, *et al.* Ocular circulatory changes following scleral buckling procedures. *Br J Ophthalmol* 1992;76:529–31.
- 12 Tamaki Y, Araie M, Kawamoto E, *et al.* Non-contact two-dimensional measurement of microcirculation in choroid and optic nerve head using laser speckle phenomenon. *Exp Eye Res* 1995;60:373–84.
- 13 Sugiyama T, Utsumi T, Azuma I, *et al.* Measurement of optic nerve head circulation: comparison of laser speckle and hydrogen clearance methods. *Jpn J Ophthalmol* 1996;40:339–43.

- 14 Tamaki Y, Araie M, Kawamoto E, et al. Noncontact, two-dimensional measurement of retinal microcirculation using laser speckle phenomenon. *Invest Ophthalmol Vis Sci* 1994;35:3825-34.
- 15 Tamaki Y, Araie M, Tomita K, et al. Time change of nicardipine effect on choroidal circulation in rabbit eyes. *Curr Eye Res* 1996;15:543-8.
- 16 Fercher AF, Briers JD. Flow visualization by means of single-exposure speckle photography. *Opt Commun* 1981;37:326-30.
- 17 Briers JD, Fercher AF. Retinal blood-flow visualization by means of laser speckle photography. *Invest Ophthalmol Vis Sci* 1982;22:255-9.
- 18 Tamaki Y, Araie M, Tomita K, et al. Real-time measurements of tissue circulation in optic nerve head or choroid in human eyes using laser speckle phenomenon. *Jpn J Ophthalmol* 1997;4:49-54.
- 19 Koelle JS, Riva CE, Petrig BL, et al. Depth of tissue sampling in the optic nerve head using laser Doppler flowmetry. *Lasers Med Sci* 1993;8:49-54.
- 20 Isono H, Kimura Y, Aoyagi K. Analysis of choroidal blood flow by laser speckle flowgraphy. *Nippon Ganka Gakkai Zasshi (J Jpn Ophthalmol Soc)* 1997;101:684-91.
- 21 Isono H. *Invest Ophthalmol Vis Sci* 1997;38(Suppl):1050.
- 22 Tomidokoro A, Araie M, Tamaki Y, et al. In vivo measurement of iridial circulation using laser speckle phenomenon. *Invest Ophthalmol Vis Sci* 1998;39:364-71.
- 23 Diddie KR, Ernest JT. Uveal blood flow after 360° constriction in the rabbit. *Arch Ophthalmol* 1980;78:503-66.
- 24 Mano T. Choroidal circulation after encircling procedure I. Immediate effect. *Folia Ophthalmol Jpn* 1982;33:460-7.
- 25 Ohkubo H. Fluorescein angiographic findings in the detached and reattached retina. *Jpn J Ophthalmol* 1988;32:423-8.