Matrix metalloproteinases and their natural inhibitors in fibrovascular membranes of proliferative diabetic retinopathy

Joel Salzmann, G Astrid Limb, Peng T Khaw, Zdenek J Gregor, Lynne Webster, Anthony H Chignell, David G Charteris

Abstract

Aim—To examine epiretinal membranes of proliferative diabetic retinopathy (PDR) for the presence of selective matrix metalloproteinases (MMPs) and their natural inhibitors (TIMPs), in order to determine whether neovascularisation and fibrosis, characteristic of this complication of diabetes mellitus, are associated with specific anomalies of MMP or TIMP expression.

Methods—The presence of selected MMPs and TIMPs was investigated in 24 fibrovascular epiretinal membranes of PDR, and the findings compared with that observed in 21 avascular epiretinal membranes of proliferative vitreoretinopathy (PVR) and five normal retinas. Specimens were examined for deposition of interstitial collagenase (MMP-1), stromelysin-1 (MMP-3), gelatinase A (MMP-2), gelatinase B (MMP-9), and three tissue inhibitors of metalloproteinases (TIMP-1, TIMP-2, and TIMP-3).

Results—The results showed that unlike normal retina, which constitutively expresses MMP-1 and TIMP-2, a large proportion of PDR membranes (> 62%) stained for MMP-1, MMP-2, MMP-3, MMP-9, TIMP-1, TIMP-2, and TIMP-3. There were no differences in the expression of these molecules when compared with PVR membranes. A characteristic staining for MMP-9 was observed within the perivascular matrix of PDR membranes, and there was a significant increase in TIMP-2 expression by PDR membranes ($p = 0.036$) when compared with PVR membranes.

Conclusions—The findings that MMPs involved in degradation of fibrovascular tissue matrix, as well as TIMP-1 and TIMP-2, are found in a large proportion of PDR membranes, and that their expression does not differ from that of PVR membranes, suggest the existence of common pathways of extracellular matrix degradation in pathological processes leading to retinal neovascularisation and fibrosis.

(Br J Ophthalmol 2000;84:1091–1096)

Proliferative diabetic retinopathy (PDR) is a common complication of diabetes mellitus characterised by preretinal neovascularisation and development of epiretinal fibrovascular tissue.¹ Early features of diabetic retinopathy include selective loss of intramural pericytes from retinal capillaries which leads to damage of the inner blood-retinal barrier.¹ This process is followed by neovascularisation, involving the production of angiogenic factors,^{2,3} as well as synthesis of extracellular matrix (ECM) necessary for anchorage of migrating endothelium and other cells such as retinal pigment epithelium (RPE), glial cells, and fibroblasts.⁴ Degradation of ECM proteins is exerted by matrix metalloproteinases (MMPs),⁵ a family of zinc binding, calcium dependent enzymes, whose activity is in turn regulated by natural inhibitors known as tissue inhibitors of metalloproteinases (TIMPs).⁶ MMPs have an important role in connective tissue remodelling and in the degradation of basement membrane and surrounding extracellular matrix during processes of angiogenesis,^{7,8} while TIMPs are multifunctional proteins with both cell growth promoting and inhibitory activities.⁹ A balance between MMP and TIMP activities may preserve tissue integrity, while excessive MMP or TIMP production may lead to excessive degradation or increased accumulation of extracellular matrix.^{5,6} It is therefore possible that an imbalance between MMPs and TIMPs may lead to the neovascularisation that characterises the development of retinal abnormalities in insulin dependent diabetes mellitus (IDDM), and to the fibrocellular proliferation observed in neovascular and avascular retinopathies.

Increased levels of MMP-9, a type IV collagenase implicated in tumour angiogenesis,¹⁰ are found in vitreous from eyes with PDR¹¹ and PVR,^{12,13} and retinal membranes of proliferative vitreoretinopathy (PVR) stain for MMPs 1, 2, 3, and 9, as well as TIMP-1 and TIMP-2.¹⁴ This contrasts with the expression of MMP-1 and TIMP-2 but not other MMPs by normal retina.¹⁴ Although PVR and PDR have different aetiologies and clinical characteristics, retinal membranes from both conditions share the features of fibroplasia, excessive

Institute of
Ophthalmology and
Moorfields Eye
Hospital, London, UK
J Salzmann
G A Limb
P T Khaw
Z J Gregor
D G Charteris

St Thomas's Hospital,
London
L Webster
A H Chignell

Correspondence to:
Dr G A Limb, Department
of Pathology, Institute of
Ophthalmology, 11–43 Bath
Street, London EC1V 9EL,
UK
g.limb@ucl.ac.uk

Accepted for publication
23 May 2000

matrix protein deposition, and cellular infiltration.¹⁵ They differ in that PDR membranes are highly vascular, owing to the angiogenic activity that takes place within the diabetic retina,¹⁶ while PVR membranes are relatively avascular¹⁷ and are not regarded as a complication of systemic disease. At present it is not known whether retinal tissues from eyes with PDR express MMPs or TIMPs associated with neovascularisation, or if there are differences in the expression of these molecules between epiretinal membranes of PDR and PVR.

The aim of this study was to examine PDR membranes for the presence of MMPs and TIMPs potentially responsible for neovascularisation and fibrosis, and to compare their expression with that observed in avascular retinal membranes from non-diabetic patients and normal retina. On this basis we compared PDR and PVR membranes for the presence of interstitial collagenase (MMP-1), stromelysin-1 (MMP-3), gelatinase A (MMP-2), gelatinase B (MMP-9), and their natural inhibitors TIMP-1, TIMP-2, and TIMP-3.

Experimental tissue and methods

Epiretinal membranes were obtained from 24 eyes undergoing pars plana vitrectomy for treatment of diabetic tractional retinal detachment, and from 21 eyes with rhegmatogenous retinal detachment complicated by PVR. Normal retina was obtained from five cornea donor eyes 12–24 hours post mortem. Four membranes derived from patients with non-insulin dependent diabetes mellitus (NIDDM). The study was performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. Membranes were collected in 4% paraformaldehyde in phosphate buffered saline (PBS) and fixed for 1–2 hours. After immersion for 1 hour in 15% sucrose in PBS, tissue was embedded in OCT compound (BDH, UK)

and stored in liquid nitrogen until use. Serial cryostat sections (7 µm thick) were placed on slides coated with poly-L-lysine (0.01% w/v: Sigma, UK), fixed in 4% paraformaldehyde for 5 minutes followed by 15% sucrose for 10 minutes, and processed for immunohistochemistry.

IMMUNOHISTOCHEMICAL DETECTION OF MMPs AND TIMPs

The presence of MMPs and TIMPs was determined by immunohistochemical staining using the alkaline phosphatase anti-alkaline phosphatase detection system, performed according to our published methods.^{14, 18} Tissue sections were washed in TRIS buffered saline pH 7.4 (0.05M TRIS in 0.15NaCl TBS) and incubated for 3 hours with a panel of commercially available mouse monoclonal antibodies to selected human MMPs and TIMPs (Oncogene research products: Calbiochem-Novabiochem International, Nottingham, UK). All antibodies were diluted in 0.5% blocking reagent (Boehringer Mannheim) in TBS to a final concentration of 10 µg/ml. These included anti-MMP-1 (clone 41-1E5, detecting active and latent forms), anti-MMP-2 (clone 42-5D11, detecting latent form), anti-MMP-3 (clone 55-2A4, detecting active and latent form), anti-MMP-9 (clone 6-6B, detecting active and latent forms), anti-TIMP-1 (clone 7-6C1), anti TIMP-2 (clone T2-101), and anti-TIMP-3 (clone 136-13H4). These antibodies are known to be specific for the above molecules as previously confirmed by zymography¹⁹ and to identify epitopes that are preserved by fixation with PFA.¹⁴ Mouse total IgG antibody (Sigma, UK) was used as a negative control in the assay. Non-specific binding of the antibodies to constitutive proteins of retinal membranes was excluded by the fact that not all membranes stained with each of the antibodies used in the study.

Table 1 Presence of selected matrix metalloproteinases (MMPs) and their natural inhibitors in PDR membranes

Membrane No	Type of diabetes	MMP-1	MMP-2	MMP-3	MMP-9	TIMP-1	TIMP-2	TIMP-3	Total mouse IgG
2	IDDM	+++	–	–	–	–	+	ND	–
3	IDDM	–	–	–	+	+	–	ND	–
4	IDDM	+++	–	–	++	+	+	++	–
5	IDDM	–	–	+	–	–	–	ND	–
6	IDDM	+	++	++	++	+	–	+	–
7	IDDM	–	–	–	–	–	–	ND	–
8	IDDM	+++	–	++	+++	+	+	–	–
9	IDDM	–	–	–	–	–	–	ND	–
10	IDDM	–	+	++	++	+	+	ND	–
11	IDDM	+	+	++	++	++	+	+	–
12 ^a	IDDM	+++	+++	+++	+++	++	++	ND	–
13 ^b	IDDM	–	++	–	+++	–	–	–	–
15	IDDM	+++	+++	+	+	++	+	ND	–
16	IDDM	++	++	–	–	–	+	ND	–
17	IDDM	++	+	+++	+	++	+	++	–
19	IDDM	+++	+++	+++	+++	++	++	++	–
21	IDDM	+++	+	+	–	–	+	–	–
22	IDDM	++	++	–	+++	++	+	–	–
23 ^c	IDDM	–	+	–	+	+	–	ND	–
24 ^d	IDDM	–	–	+	–	+	–	ND	–
1	NIDDM	+++	++	++	++	+	–	++	–
14	NIDDM	+++	+++	+++	–	+++	+	ND	–
18	NIDDM	+++	+++	++	+++	++	+	++	–
20	NIDDM	++	++	+++	+++	++	+	+	–

[a–b] Same patient; left and right eye respectively; [c–d] same patient, right and left eye respectively. Intensity of staining: +++ = strong, ++ = moderate, + = mild, – = negative, ND = not done.

Figure 1 Photomicrographs of PDR membranes showing deposition of selective MMPs and TIMPs. The pink precipitate indicates the presence of these molecules within the membranes. Specimens were counterstained with Mayer's haematoxylin. (A) PDR membrane (No 15) showing strong staining (+++) for MMP-1. (B) PDR membrane (No 12) showing intense cellular and extracellular staining (+++) for MMP-2. (C) PDR membrane (No 1) exhibiting staining for MMP-9 within the extracellular and perivascular matrix. (D) Section of PDR membrane (No 17) showing detail staining for MMP-9 in the perivascular matrix. (E) PDR membrane exhibiting moderate staining (++) for TIMP-1 (No 11). (F) Normal retina showing negative staining for MMP-9.

After incubation with the primary antibody, specimens were washed for 2 minutes in TBS, incubated for 30 minutes with 50 µl rabbit anti-mouse antibody (Dako, UK), and further washed for 2 minutes in TBS. Sections were then incubated with 50 µl monoclonal mouse alkaline phosphatase anti-alkaline phosphatase antibodies (Dako) for 30 minutes and washed for 2 minutes. The last two steps were repeated for 10 minutes each to enhance staining intensity, followed by incubation for 30 minutes with developing substrate (1 mg/ml fast red in 0.1M TRIS buffer pH 8.2 with 1.2 mM naph-

thol AS-MX phosphate, 0.25 dimethylformamide, and 1 mM levamisole). Specimens were finally washed in distilled water, counterstained with Mayer's haematoxylin, and mounted with glycerol mounting medium (Dako).

Membranes were examined under light microscopy for positive staining identified by a red precipitate. Intensity of staining was graded by two independent observers into four categories: negative (-), mild (+), moderate (++), and strong (+++), according to the strength and distribution of the red precipitate.

Table 2 Comparison between retinal membranes of PDR and PVR and normal retina in the expression of MMPs and TIMPs

	MMP-1 (active + latent)	MMP-2 (latent)	MMP-3 (active + latent)	MMP-9 (active + latent)	TIMP-1	TIMP-2*	TIMP-3
PDR	66%	66%	62%	66%	71%	62%	66%
PVR	86%	81%	52%	48%	43%	33%	75%
Normal retina	100% ^a	0%	0%	0%	0%	100% ^b	0%

The figures represent the percentage of samples staining for each antibody in the different groups.

* $p = 0.036$ v PVR.

^aStaining of the inner and outer nuclear cell layers and plexiform layers.

^bStaining of the inner and outer nuclear cell layers only.

STATISTICAL ANALYSIS

The significance of difference between corresponding groups of observations was evaluated by the non-parametric Fisher's exact test. Acceptable significance was recorded when p values were < 0.05 .

Results

EXPRESSION OF MMPs AND TIMPs IN PDR MEMBRANES

Table 1 shows the distribution of MMPs and TIMPs within the extracellular matrix of epiretinal membranes of PDR. The majority of membranes (22/24, 92%) stained for at least one of the MMPs or TIMPs examined, while two of the membranes did not show deposition of any of the molecules investigated. None of the specimens stained with all antibodies to MMPs and TIMPs used in this study. Of the 24 membranes analysed, 16 (66%) stained for interstitial collagenase (MMP-1) (Fig 1A), 16 (66%) for gelatinase A (MMP-2) (Fig 1B), and 15 (62%) for stromelysin-1 (MMP-3). Expression of gelatinase B (MMP-9) was observed in 16 membranes (66%), some of which showed staining of the perivascular extracellular matrix (Figs 1C and D). Presence of TIMP-1 (Fig 1D) and TIMP-2 was observed in 17 (71%) and 15 (62%) membranes respectively, and the intensity of staining for TIMP-1 was in general greater than that for TIMP-2 (Table 1). TIMP-3 was observed in 8/12 (66%) membranes and none of the membranes investigated stained with total mouse IgG (Table 1).

Of the 16 specimens stained for MMP-1, 13 were also positive for MMP-2, 12 for MMP-3, 12 for MMP-9, 13 for TIMP-1, and 14 for TIMP-2 (Table 1). All membranes staining for TIMP-3 (eight) were also positive for TIMP-1 and MMP-1, and six were positive for TIMP-2. Of the 15 membranes staining for MMP-3, 10 also stained for both MMP-2 and MMP-9, and 12 stained for TIMP-1 (Table 1). Thirteen of the 16 membranes positive for MMP-9 also stained for MMP-2, 15 for TIMP-1, and 11 for TIMP-2. Twelve of the 17 specimens positive for TIMP-1 also stained for TIMP-2 (Table 1), but overall, the intensity of staining for TIMP-1 was higher than that for TIMP-2 (Table 1). The strongest staining for any of the molecules investigated was observed with antibodies to MMP-1, and no differences in the staining for MMPs and TIMPs were evident when comparing membranes from patients with IDDM and NIDDM. Four of PDR membranes that derived from two patients who underwent pars plana vitrectomy in their left and right eyes on different occasions did not

show the same pattern of staining for the MMPs and TIMPs investigated (Table 1).

COMPARISON BETWEEN EPIRETINAL MEMBRANES OF PDR AND PVR AND NORMAL RETINA IN THE EXPRESSION OF MMPs AND TIMPs

There was no significant difference between the proportion of PDR membranes staining for MMP-3 (15/24) and MMP-9 (16/24) when compared with PVR membranes (11/21 and 10/21, respectively; $p > 0.5$) (Table 2). The proportion of PDR membranes staining for TIMP-1 (17/24) was slightly higher than that of PVR membranes staining for this molecule (9/21), but this difference was not significant ($p = 0.075$). However, the percentage of PDR membranes staining for TIMP-2 (15/24) was significantly higher when compared with that of PVR membranes (7/21; $p = 0.036$). A similar proportion of TIMP-3 positive membranes was observed in both PDR (8/12) and PVR (15/20).

As previously shown,¹⁴ normal retina did not express MMP-9 (Fig 1F, Table 2), but stained for MMP-1 in the inner and outer nuclear and plexiform layers, as well as for TIMP-2 in both the inner and outer nuclear layers (Table 2).

Discussion

This study shows that a large proportion of PDR membranes (92%) exhibited deposition of MMPs potentially responsible for degrading matrix constituents of blood vessels and epiretinal membranes. In addition, a large percentage of PDR membranes (75%) also expressed TIMPs, which by inhibiting matrix degradation by MMPs may promote its accumulation with resultant fibrosis and new vessel formation. The proportion of membranes staining for the MMPs investigated was similar in both PDR and PVR, suggesting that the same molecular mechanisms of extracellular matrix degradation and deposition may operate in vascular and avascular proliferative processes affecting the retina.

Since MMP-1 is constitutively expressed in normal retina,¹⁴ its presence in retinal membranes of both PVR and PDR could be anticipated. It is possible that this MMP may have a physiological role within the retina, such as aiding degradation of the shed tips of photoreceptors, and this merits further investigation. Membrane expression of MMP-2, MMP-3, and MMP-9, which are not observed in normal retina may indicate an active involvement of these molecules in the neovascularisation observed in PDR and in the fibrocellular proliferation observed in both PDR and PVR. Early vascular changes in diabetic retinopathy

include the loss of pericytes¹ and it has been postulated that this phenomenon may be facilitated by MMP degradation of basement membrane and surrounding matrix.^{7,8} In this context, our observation that matrix surrounding the vessels of PDR membranes stains for MMP-9 (Fig 1C, D), strongly suggests that this molecule, which degrades denatured collagens and type IV basement membrane collagen, may contribute to retinal neovascularisation in PDR. That MMP-9 may be actively involved in diabetic retinal neovascularisation is supported by observations that high vitreous levels of this molecule may be found in eyes with PDR,^{11,12} and that microvascular endothelial permeability is increased in the presence of this MMP.²⁰ This is further supported by the demonstration that levels of expression of MMP-9 correlate with tumour neovascularisation in vivo,¹⁰ and that increased plasma levels of this molecule precede the development of renal microvascular complications in patients with IDDM.²¹ This is of special interest, as renal microangiopathy is often associated with proliferative retinopathy in diabetic individuals.²²

This study showed a slight but not statistically significant increase in the percentage of PDR membranes staining for TIMP-1 (70%) when compared with PVR membranes (42%; $p = 0.06$). In contrast, there was a significant increase in the proportion of PDR membranes staining for TIMP-2 (62%) when compared with PVR membranes (33%; $p = 0.036$). These observations suggest that differences in the kinetic mechanisms of TIMP production and activity within the retina may determine the development of different patterns of matrix accumulation, and therefore whether new vessels are formed during a pathological process. Although TIMP-1 and TIMP-2 inhibit angiogenesis and exhibit growth factor-like activity,⁹ it is difficult to speculate on their specific role in the process of neovascularisation in the diabetic retina. It may be possible that these molecules are produced to control MMP activity, but if excessive production of MMPs occurs, or insufficient levels of TIMPs are released, they may not exert an adequate anti-angiogenic activity.

It is unlikely that the MMPs and TIMPs observed in PDR membranes derived from the microvascular compartment, as there were no differences with that observed in membranes from eyes with PVR, in which no vitreous haemorrhage was present and in which neovascularisation does not occur. However, it is possible that these molecules may be locally produced by cells of the retinal microenvironment, such as RPE and glial cells, which are known to produce various MMPs and TIMPs.²³⁻²⁵

The mechanisms that trigger release of MMPs and TIMPs during retinal neovascularisation are not known, but it is conceivable that cytokines produced upon endothelial cell activation by ischaemia may induce local release of these molecules. It is suggested by observations that cytokines such as tumour necrosis factor α (TNF α) and interleukin-1

(IL-1), which are found in vitreous²⁶ and membranes¹⁸ from eyes with PDR, induce production of MMPs and TIMPs by vascular endothelial cells, fibroblasts, and retinal pigment epithelial cells.^{24,25,27,28} This is of special relevance to the pathogenesis of PDR, as there is an association between susceptibility to retinopathy and TNF α polymorphism in individuals with NIDDM,²⁹ and subjects with IDDM expressing the TNF2 allele produce high levels of TNF α .³⁰

The observations that MMPs and TIMPs implicated in neovascularisation and fibrosis are found within PDR and PVR membranes, indicate that although both conditions have different aetiologies and clinical characteristics, these molecules may play an important part in the pathogenesis of both conditions. Further studies may elucidate whether deposition of MMP-9 in the perivascular matrix is unique to PDR, or whether it is a common feature of all microvascular complications of diabetes mellitus. From the present findings it is not clear whether there are selective defects in the balance between MMPs and TIMPs which either promote fibrosis and neovascularisation in PDR, or fibrosis in PVR. However, investigation of the kinetics of production of selective MMPs and their control by specific TIMPs may aid in the design of new therapeutic approaches to treat and prevent PDR and other microvascular complications of diabetes mellitus.

Supported by the Henry Smith Charity.

- Garner A. Histopathology of diabetic retinopathy in man. *Eye* 1993;7:250-3.
- Forrester JV, Shafiee A, Schröder S, *et al.* The role of growth factors in proliferative diabetic retinopathy. *Eye* 1993;7:276-87.
- Adamis AP, Miller JW, Bernal M-T, *et al.* Increased vascular endothelial growth factor levels in the vitreous of eyes with proliferative diabetic retinopathy. *Am J Ophthalmol* 1994;118:445-50.
- Hiscott P, Waller HA, Grierson I, *et al.* The extracellular matrix of reparative tissue in the vitreous: fibronectin production in proliferative diabetic retinopathy membranes. *Eye* 1993;7:288-92.
- Matrisian LM. The matrix-degrading metalloproteinases. *Bio Assays* 1992;14:455-63.
- Dollery CM, McEwan JR, Henney AM. Matrix metalloproteinases and cardiovascular disease. *Circ Res* 1995;77:863-8.
- Taylor CM, Thompson JM, Weiss JB. Matrix integrity and the control of angiogenesis. *Int J Radiat Biol* 1991;60:61-4.
- Furness PN. Basement membrane synthesis and degradation. *J Pathol* 1997;183:1-3.
- Gomez DE, Alonso DF, Yoshiji H, *et al.* Tissue inhibitors of metalloproteinases: structure, regulation and biological functions. *Eur J Cell Biol* 1997;74:111-22.
- Kurizaki T, Toi M, Tominaga T. Relationship between matrix metalloproteinase expression and tumor angiogenesis in human breast carcinoma. *Oncol Rep* 1998;60:673-7.
- Brown D, Hamdi H, Bahri S, *et al.* Characterization of an endogenous metalloproteinase in human vitreous. *Curr Eye Res* 1994;13:639-47.
- De La Paz MA, Itoh Y, Toth CA, *et al.* Matrix metalloproteinases and their inhibitors in human vitreous. *Invest Ophthalmol Vis Sci* 1998;39:1256-60.
- Kon CH, Occlleston NL, Charteris DG, *et al.* A prospective study of matrix metalloproteinases in proliferative vitreoretinopathy. *Invest Ophthalmol Vis Sci* 1998;39:1524-9.
- Webster L, Chignell AH, Limb GA. Predominance of MMP-1 and MMP-2 in epiretinal and subretinal membranes of proliferative vitreoretinopathy. *Exp Eye Res* 1999;68:91-8.
- Gilbert C, Hiscott P, Unger W, *et al.* Inflammation and the formation of epiretinal membranes. *Eye* 1988;(Suppl)2:S140-56.
- Hiscott P, Waller HA, Grierson I, *et al.* The extracellular matrix of reparative tissue in the vitreous: fibronectin production in proliferative diabetic retinopathy membranes. *Eye* 1993;7:288-92.

- 17 Jerdan JA, Pepose JS, Michels RG, *et al*. Proliferative vitreoretinopathy membranes. An immunocytochemical study. *Ophthalmology* 1989;**96**:801–10.
- 18 Limb GA, Chignell AH, Green W, *et al*. Distribution of TNF α and its reactive vascular adhesion molecules in fibrovascular membranes of proliferative diabetic retinopathy. *Br J Ophthalmol* 1996;**80**:168–73.
- 19 Guo L, Hussain AA, Limb GA, *et al*. Age-dependent variation in metalloproteinase activity of isolated human Bruch's membrane and choroid. *Invest Ophthalmol Vis Sci* 1999;**40**:2676–82.
- 20 Partridge CA, Jeffrey JJ, Malik AB. A 96kDa gelatinase induced by TNF- α contributes to increased microvascular endothelial permeability. *Am J Physiol* 1993;**265**:L438–47.
- 21 Ebihara I, Nakamura T, Shimada N, *et al*. Increased plasma metalloproteinase-9 concentrations precede development of microalbuminuria in non-insulin dependent diabetes mellitus. *Am J Kidney Dis* 1998;**32**:544–50.
- 22 Skyler JS. Diabetic complications. *Endocrin Metab N Am* 1996;**25**:243–55.
- 23 Hunt RC, Fox A, Pakalnis VA, *et al*. Cytokines cause cultured retinal pigment epithelial cells to secrete metalloproteinases and to contract collagen gels. *Invest Ophthalmol Vis Sci* 1993;**34**:3179–86.
- 24 Alexander JP, Bradley JMB, Gabourel JD, *et al*. Expression of matrix metalloproteinases and inhibitor by human retinal pigment epithelium. *Invest Ophthalmol Vis Sci* 1990;**31**:2520–8.
- 25 Gottschall PE, Yu X. Cytokines regulate gelatinase A and B (matrix metalloproteinase 2 and 9) activity in cultured rat astrocytes. *J Neurochem* 1995;**64**:1513–20.
- 26 Franks WA, Limb GA, Stanford MR, *et al*. Cytokines in human intra-ocular inflammation. *Curr Eye Res* 1992;**11**:187–91.
- 27 Grant MB, Caballero S, Tarnuzzer RW, *et al*. Matrix metalloproteinase expression in human retinal microvascular cells. *Diabetes* 1998;**47**:1311–17.
- 28 Schwartz DJ, Monea S, Marcus SG, P *et al*. Soluble factor(s) released from neutrophils activates endothelial cell matrix metalloproteinase-2. *J Surg Res* 1998;**76**:79–85.
- 29 Hawrami K, Hitman GA, Rema M, *et al*. An association in non-insulin-dependent diabetes mellitus subjects between susceptibility to retinopathy and tumor necrosis factor polymorphism. *Human Immunol* 1996;**46**:49–54.
- 30 Pociot F, Briant L, Jongeneel CV, *et al*. Association of tumor necrosis factor (TNF) and class II major histocompatibility alleles with secretion of TNF α and TNF β by human monocular cells: a possible link to insulin-dependent diabetes mellitus. *Eur J Immunol* 1993;**23**:224–31.

Contributors please note:

Communications from **all countries except the UK and Republic of Ireland** should be sent to Professor C Hoyt, Editor, *British Journal of Ophthalmology*, University of California, Department of Ophthalmology, 10 Kirkham Street, K 301, San Francisco, CA 94143-0730, USA (tel: 001 415 502-6871; fax: 001 415 514-1521).

Manuscripts from the **UK and the Republic of Ireland** should be sent to Professor Andrew Dick, UK Editor, *British Journal of Ophthalmology*, Division of Ophthalmology, University of Bristol, Lower Maudlin Street, Bristol BS1 2LX (tel: +44 (0) 0117 929-4496; fax: +44 (0)117 929-4607).