Surgical removal of subfoveal choroidal neovascularisation in highly myopic patients

José M Ruiz-Moreno, Concepción de la Vega

Abstract

Aim—To analyse the visual results obtained in the treatment of subretinal choroidal neovascularisation (CNV) in patients with high myopia by vitrectomy and extraction of the neovascular membrane.

Methods—22 eyes of 22 patients with high myopia (>-6 dioptres and/or axial length >26 mm) with subfoveal CNV treated by extraction of the CNV were analysed retrospectively. The patients' mean age was 60.27 (SD 16.41) years (range 32–83 years). The mean follow up was 29.3 (9.9) months (range 12–42 months).

Results—Best corrected visual acuity (BCVA) preoperatively was 0.09 (0.07) (range 0.01–0.3). After treatment BCVA was 0.12 (0.10) (range 0.01–0.4), a difference without statistical significance ($p=0.03$, Student's t test paired data). In four cases the CNV recurred, in three cases cataract developed that required extraction, and in one case retinal detachment occurred in the early postoperative period. Topical treatment was necessary in two cases to lower the intraocular pressure.

Conclusions—The treatment of subfoveal CNV in highly myopic patients by surgical removal by vitrectomy does not achieve any significant improvement of the BCVA. (*Br J Ophthalmol* 2001;85:1041–1043)

High myopia (more than -6 dioptres and/or longer than 26 mm of axial length), is the seventh most common cause of legal blindness in adults in the USA.¹ It is present in 0.5% of the

European population² and it is estimated to represent 2% of all types of myopia.³

The excessive elongation of the anterior-posterior axis of the eye is associated with complications⁴ such as choroidal neovascularisation (CNV), frequently with ruptures of Bruch's membrane,⁵ that appear in 4.3% of the patients with high myopia.⁶

CNV varies from 4% to 11% of highly myopic patients.⁷ Its presence is confirmed by fluorescein angiography, demonstrating the typical pattern of neovascular membranes.

One proposed therapy is macular surgery with removal of the CNV.⁸ We present in this paper the results obtained in the retrospective review of a consecutive series of patients with high myopia and subfoveal CNV following such surgery.

Patients and methods

For this study we have retrospectively reviewed 22 eyes of 22 consecutive patients with high myopia, in which subfoveal CNV was diagnosed, being treated by vitrectomy and surgical removal of the CNV.

As criteria of high myopia we used the existence of a refractive error greater than -6 dioptres and/or an axial length greater than 26 mm.

Selection criteria:

- Preoperative best corrected visual acuity (BCVA) less than or equal to 0.3 (decimal scale). A progressive loss of BCVA during two consecutive examinations was documented and the decline in acuity was believed to be due to the growth of the CNV
- No other ocular pathologies associated with high myopia
- No previous treatment by laser in the foveal avascular zone
- Clinical evidence of CNV, confirmed by fluorescein angiography
- Informed written consent
- All patients with clinical signs of age related macular degeneration (ARMD), such as soft drusen, presumed ocular histoplasmosis syndrome (POHS), as well as other pathologies that may develop CNV in their evolution, were excluded.

The average age of the patients was 60.27 (SD 16.41) years (range 32–83 years), of which nine were female and 13 male. In 14 patients the right eye was operated on and in eight patients the left. The average follow up was 29.3 (7.9) months (range 12–42 months).

The surgical technique was performed as described previously^{9 10} under local anaesthesia and as outpatients.

Face down positioning during the first 24 hours after surgery was maintained and topical medication begun with steroids and antibiotics.

Department of
Ophthalmology,
Miguel Hernández
University School of
Medicine, Alicante,
Spain
J M Ruiz-Moreno

Instituto
Oftalmológico de
Alicante
(Vitreoretinal Unit)
J M Ruiz-Moreno
C de la Vega

Correspondence to:
José M Ruiz-Moreno,
División de Oftalmología,
Universidad Miguel
Hernández, Campus de San
Juan, 03550, Alicante, Spain
jm.ruiz@umh.es

Accepted for publication
27 March 2001

Table 1 Characteristics of myopic patients with CNV

Patient No	Age (years)	Sex	Eye	Initial BCVA	Final BCVA	Follow up (months)	Complications
1	69	Female	Right	0,03	0,02	29	RD
2	66	Male	Right	0,16	0,04	24	Cataract
3	34	Male	Right	0,05	0,1	37	
4	41	Male	Right	0,05	0,16	41	
5	70	Female	Right	0,05	0,3	31	Recurrent CNV
6	52	Male	Left	0,1	0,16	37	Glaucoma
7	74	Male	Left	0,3	0,3	30	
8	83	Male	Right	0,1	0,2	12	
9	78	Female	Right	0,04	0,03	20	
10	38	Male	Right	0,16	0,4	13	Recurrent CNV
11	50	Male	Right	0,1	0,1	28	Recurrent CNV
12	47	Male	Left	0,1	0,16	26	Glaucoma
13	71	Female	Left	0,03	0,03	26	Recurrent CNV
14	58	Female	Right	0,2	0,1	26	
15	82	Male	Left	0,01	0,01	32	
16	74	Male	Right	0,2	0,16	39	
17	61	Female	Right	0,1	0,1	35	Cataract
18	70	Male	Right	0,02	0,03	32	Cataract
19	76	Male	Left	0,03	0,02	42	
20	64	Female	Left	0,04	0,04	35	
21	32	Female	Right	0,01	0,1	27	
22	36	Male	Left	0,1	0,1	24	

BCVA = best corrected visual acuity. RD = retinal detachment.

Table 2 Postoperative evolution of best corrected visual acuity

Follow up	Average	Standard deviation	Minimum.	Maximum	No
2 months	0.10	0.09	0.01	0.4	22
8 months	0.11	0.10	0.01	0.4	22
14 months	0.09	0.08	0.01	0.3	20
20 months	0.09	0.07	0.01	0.3	20
26 months	0.1	0.1	0.01	0.3	17

Results

We analysed the results of 22 patients with high myopia who underwent surgery for removal of subfoveal CNV (Table 1). The mean BCVA before surgery was 0.09 (0.07) (range 0.01–0.3); at the end of the follow up period the mean BCVA was 0.12 (0.1) (range 0.01–0.4), the difference not being statistically significant ($p=0.113$, Student's t test paired data).

If we study the evolution of BCVA after surgery we find that the average BCVA from 2 months postoperatively until 26 months does not decrease with time ($p>0.05$; Anova test, Table 2).

We found an improvement of BCVA after surgery in nine (40.9%) patients, a worsening in six (27.3%), and no variation in another seven (31.8%).

CNV recurred in four patients who were treated by green argon laser photocoagulation.

Of the 22 patients three developed cataracts which required surgery, and in one patient retinal detachment appeared 3 months after surgery. This patient was operated on with scleral buckling without further complications. We needed to use chronic topical medication to control the intraocular pressure in two cases.

Discussion

High myopia is the first aetiological cause of CNV in young people.¹¹ In a study of 308 eyes from 202 patients with high myopia, CNV appeared in 5.2% of the eyes.¹² The natural evolution of CNV in high myopia is very poor final visual acuity.^{13, 14}

One of the possible treatments of extrafoveal and juxtafoveal CNV is direct laser photocoagulation. In high myopia a progressive growth of the scar after photocoagulation has been described, without relation to the wavelength of the laser, the axial length of the eye, or to other factors.¹⁵ Comparing this it has been noticed that the growth of the scar following CNV photocoagulation in ARMD was 28.7%, while in high myopia it was 109.2%.¹⁶

Avila¹⁷ affirms that the results in patients treated with laser cannot be distinguished from patients without treatment, finding a drop in visual acuity in 57% in those treated and in 46% of those not treated. It is difficult to compare the results of this study with others carried out, but it is certainly clear that natural evolution and the results of laser treatment of the subfoveal CNV in myopic eyes giving very poor visual acuity.⁹

One of the therapeutic alternatives was surgery to extract the CNV. At the present time two new treatment possibilities include macular translocation and photodynamic therapy.

The majority of the reports regarding submacular surgery are related to the CNV associated with ARMD or to POHS. Adelberg⁹ published a paper about the surgical extraction of CNV in 17 eyes, five of which were highly myopic. The visual acuity after surgery improved in two of the five eyes, with recurrence in one.

For Thomas,¹⁸ the result in 10 eyes after surgical removal of CNV in myopic patients, is a lessening of visual acuity, this being 0.2 or lower in all eyes and with a recurrent CNV in two of the 10 eyes. In another later paper, of 23 eyes, eight (35%) had a final visual acuity of 20/40 or better, five (22%) between 20/50 and 20/100, and 10 (43%) between 20/200 and 20/400.¹⁹ Other authors have described positive results from surgery with improvement in 45% of eyes and no change in 37%.²⁰

The 22 patients selected for this paper represent a group of patients with high myopia who developed CNV as a result of the degenerative processes associated with their ocular disease.

We found an improvement of BCVA in nine (40.9%) of the 22 patients studied. We observed no variation in seven (31.8%), and a deterioration in six (27.2%). The difference between the preoperative (0.09) (0.07) and postoperative (0.12) (0.10) BCVA was not statistically significant ($p=0.03$, Student's t test paired data).

In our patients the BCVA did not decrease with time (Table 2), whereas the visual results obtained after treatment of myopic CNV with photocoagulation did.¹⁵

In four cases the CNV recurred, three cases developed cataracts requiring surgery and in one case retinal detachment occurred.

In our series in only one patient was the final BCVA 0.4, compared with other results. It is probably due to the average age (60 years) and the inclusion criterion of waiting for two consecutive examinations to confirm progressive loss of vision, that may have allowed the CNV to disrupt the retinal pigmentary epithelium (RPE).

It is not known why in some eyes the visual prognosis worsens and in others it improves after surgery, probably being related to the amount of RPE extracted. Clinical and laboratory studies suggest that the integrity of RPE and the choriocapillaris is the principal factor which will determine the final prognosis after submacular surgery.⁹

In our experience RPE is compromised in the majority of cases and it is extracted, leaving in some patients a denuded area of Bruch's membrane and a defect in the RPE after surgery. Laboratory studies confirm that these denuded areas induce the progressive secondary atrophy of the choriocapillaris and external layers of the retina. These changes have been investigated in animal research²¹ and are transferable to the phenomena that appear after submacular surgery in our patients.²²

In conclusion, and in agreement with our results and with previous papers,^{9, 18, 19} surgical removal of the CNV in highly myopic patients stabilises the evolution in some patients

without inducing a significant improvement in visual acuity. Randomised, prospective studies will be required to evaluate the role of surgical removal of the CNV in highly myopic patients. Comparison of outcomes with other techniques such as macular translocation and photodynamic therapy will also be required in order to determine which is the best treatment for subfoveal CNV in high myopia.

- 1 National Society for the Prevention of Blindness. *NSPB fact book. Estimated statistics of blindness and vision problems*. New York: NSPB, 1966:44.
- 2 Stocker FW. Pathologic anatomy of the myopic eye with regard to newer theories of etiology and pathogenesis of myopia. *Arch Ophthalmol* 1943;**30**:476–88.
- 3 Michael DD. *Visual optics and refraction. A clinical approach*. 2nd ed. St Louis: CV Mosby, 1980:1985.
- 4 Curtin BJ. Physiologic vs pathologic myopia: genetics vs environment. *Ophthalmology* 1979;**86**:681–91.
- 5 Johnson DA, Yannuzzi LA, Shakin JL, et al. Lacquer cracks following laser treatment of choroidal neovascularization in pathologic myopia. *Retina* 1998;**18**:118–24.
- 6 Curtin BJ, Karlin DB. Axial length measurements and fundus changes of the myopic eye. *Am J Ophthalmol* 1971;**71**:42–53.
- 7 Curtin BJ. The posterior staphiloma of pathologic myopia. *Trans Am Ophthalmol Soc* 1977;**75**:68–86.
- 8 De Juan E Jr, Machemer R. Vitreous surgery for hemorrhagic and fibrous complications of age-related macular degeneration. *Am J Ophthalmol* 1988;**105**:25–29.
- 9 Adelberg DA, Del Priore LV, Kaplan HJ. Surgery for subfoveal membranes in myopia, angioid streaks, and other disorders. *Retina* 1995;**15**:198–205.
- 10 Ruiz-Moreno JM, Pérez-Santonja JJ. Dissection of the posterior hyaloid. *Arch Ophthalmol* 1998;**116**:1392–3.
- 11 Cohen SY, Laroche A, Leguen Y, et al. Etiology of choroidal neovascularization in young patients. *Ophthalmology* 1996;**103**:1241–4.
- 12 Grossniklaus HE, Green WR. Pathologic findings in pathologic myopia. *Retina* 1992;**12**:127–33.
- 13 Hampton GR, Kohen D, Bird AC. Visual prognosis of disciform degeneration in myopia. *Ophthalmology* 1983;**90**:923–6.
- 14 Hotchkiss ML, Fine SL. Pathologic myopia and choroidal neovascularization. *Am J Ophthalmol* 1981;**91**:177–83.
- 15 Brancato R, Pece A, Avanza P, et al. Photocoagulation scar expansion after therapy for choroidal neovascularization in degenerative myopia. *Retina* 1990;**10**:239–43.
- 16 Oshima Y, Harino S, Tano Y. Scanning laser ophthalmoscope micropertimetric assessment in patients with successful laser treatment for juxtafoveal choroidal neovascularization. *Retina* 1998;**18**:109–17.
- 17 Avila MP, Weiter JJ, Jalkh AE, et al. Natural history of choroidal neovascularization in degenerative myopia. *Ophthalmology* 1984;**91**:1573–81.
- 18 Thomas MA, Dickinson JD, Melberg NS, et al. Visual results after surgical removal of subfoveal choroidal neovascular membranes. *Ophthalmology* 1994;**101**:1384–96.
- 19 Uemura A, Thomas MA. Subretinal surgery for choroidal neovascularization in patients with high myopia. *Arch Ophthalmol* 2000;**118**:344–50.
- 20 Bottoni F, Perego E, Airaghi P, et al. Surgical removal of subfoveal choroidal neovascular membranes in high myopia. *Graefes Arch Clin Exp Ophthalmol* 1999;**237**:573–82.
- 21 Del Priore LV, Hornbeck R, Kaplan HJ, et al. Debridement of the pig retinal pigment epithelium in vivo. *Arch Ophthalmol* 1995;**113**:939–44.
- 22 Nasir M, Zarbin MA. Choriocapillaris atrophy as a complication of surgical excision of choroidal neovascular membranes. *Invest Ophthalmol Vis Sci* 1993;**34** (suppl): 834.