LABORATORY SCIENCE

Expression and distribution of matrix metalloproteinases and their inhibitors in the human iris and ciliary body

J Lan, RK Kumar, N Di Girolamo, P McCluskey, D Wakefield

Br J Ophthalmol 2003;**87**:208–211

See end of article for authors' affiliations

Correspondence to: Professor Denis Wakefield, Department of Pathology, The University of New South Wales, Sydney 2052, Australia; d.wakefield@unsw.edu.au

Accepted for publication 27 August 2002

Aim: To determine the expression and distribution of matrix metalloproteinases (MMPs) and their inhibitors (TIMPs) in the normal human iris and ciliary body.

Methods: Seven postmortem human eyes were fixed with formalin. The iris and ciliary body were dissected out and embedded in paraffin. The expression of MMPs -1, 2, 3, and 9, and TIMPs 1–4 in the iris and ciliary body was determined by a novel immunofluorescence technique and the results graded by masked observers.

Results: Positive staining for MMPs and TIMPs was observed in all regions of the anterior uvea, and was more intense in the ciliary body than in the iris. Most MMPs and TIMPs showed similar patterns in their distribution. In the ciliary body, staining was strongest in the epithelium, and was localised to the epithelial cell cytoplasm, except for TIMP-3 which was strongly expressed in the basement membranes. In the iris, staining was most noticeable in the anterior border and anterior epithelial layer. Blood vessels in the stroma of the iris and ciliary body also stained moderately for MMPs and TIMPs.

Conclusion: Both MMPs and TIMPs are widely expressed in the anterior uvea, with a positive correlation between their expressions. Their differential localisation in the ciliary body suggests they may have a role in maintaining homeostasis in the uveal tract.

The matrix metalloproteinases (MMPs) are a family of over 20 zinc dependent enzymes that cleave various components of the extracellular matrix (ECM).¹ MMPs are associated with a variety of physiological and pathological conditions that involve matrix modelling and remodelling.^{2–4} They are highly expressed in adult reproductive tissues undergoing dramatic reductions in tissue mass, including the endometrium during menstruation and the involution of the breast, uterus, and prostate.⁵ MMPs have also been shown to contribute to human fetal development and are present in postnatal tissues.⁶ They are also found in the human intervertebral disc.⁷ In wound healing, the expression of MMPs is pronounced and depends on the phase of healing.⁸ The tissue destruction that occurs in diseases such as rheumatoid arthritis,^{9–10} macular degeneration,¹¹ periodontitis,¹² and tumour cell invasion¹³ are apparently mediated by members of the MMP family. The generation of synthetic inhibitors of MMPs for therapeutic use in these diseases is thus of great interest.^{14–16}

MMPs have been classified into four broad categories based on their substrate specificity and domain organisation.¹³ They include collagenases (MMP-1, 8, 13), stromelysins (MMP-3, 10, and 11), gelatinases (MMP-2 and 9), and membrane types (MMPs 14–17, 24, 25). MMPs are regulated at several levels including transcription, secretion, activation, and inhibition. Regulation by the latest of these mechanisms is via endogenous inhibitors, known as tissue inhibitors of metalloproteinases (TIMPs), and the balance between levels of active enzymes and free TIMPs is thought to determine overall MMP activity.^{17–19} Disturbance of this equilibrium is a critical determinant of proteolysis and tissue invasion.^{20–23}

In the eye, upregulated MMPs have been found in corneal ulceration,²⁴ scleritis,²⁰ uveitis,^{21–25} and pterygia.^{26–28} These observations suggest a potential role for MMPs in the pathogenesis of inflammatory eye diseases. Despite reports on the relation between MMPs and uveal diseases,^{21–25} and growing interest in modulating MMPs/TIMPs as new therapeutic strategies for the treatment of ocular diseases,^{29–30} very little is known about the distribution of MMPs and TIMPs in the

human uveal tract. This basic knowledge is important for understanding the role of these molecules in ocular physiology, in disease, and for the development of new therapeutic strategies. In the present study, we localised the major MMPs (1, 2, 3, and 9) involved in destruction and remodelling of collagenous connective tissues, as well as their principal antagonists (TIMPs 1–4) in the human anterior uveal tract. Fluorochrome staining³¹ allowed us to examine the distribution of MMPs and TIMPs in these heavily pigmented tissues, in which conventional immunohistochemistry has been proved to be unreliable.³²

MATERIALS AND METHODS

Antibodies and reagents

Mouse anti-human antibodies to individual MMPs (MMP-1, 2, 3, 9) and TIMPs 1–4 were obtained from ICN Biomedicals, Australia. A control mouse IgG₁ antibody was run in parallel (Sigma, Australia). Goat anti-mouse immunoglobulins conjugated to Alexa 568, (excitation/emission maxima of 573/596 nm) were purchased from Molecular Probes, Eugene, OR, USA.

Dissection of human iris and ciliary body

Seven postmortem human eyes from donors (45–80 years of age) were obtained from the Lions Eye Bank (Sydney, Australia) within 24 hours after death. The eyes were fixed in 10% formalin for 24 hours. The anterior uveal tissues, iris, and ciliary body area were dissected under a microscope and embedded in paraffin.

Immunohistochemistry

Expression of MMPs and TIMPs in the iris and ciliary body was examined using mouse anti-human MMPs/TIMPs monoclonal antibodies, followed by the addition of a secondary antibody conjugated to the red emitting fluorochrome Alexa 568. This method, which has previously been validated for immunohistochemistry on paraffin sections³¹ minimises interference by autofluorescence. Tissue was serially sectioned at 4 µm thickness and placed on aminopropyltriethoxysilane

Table 1 Distribution and intensity of immunoreactivity for MMPs and TIMPs in the iris. Grading is presented as medians and interquartile ranges

Abs	Iris (n=7)			
	AB	AE	IS	PostE
MMP-1	3 (3)	2 (2)	2 (2)	1 (1)
MMP-2	3 (3)	3 (3)	2 (2)	2 (1–2)
MMP-3	3 (3)	3 (2–3)	2 (1–2)	2 (1–2)
MMP-9	2 (2–3)	2 (2)	1 (1–2)	1 (0–1)
TIMP-1	2 (2–3)	2 (1–2)	1 (1–2)	1 (1)
TIMP-2	3 (2–3)	2 (2–3)	2 (1–2)	1 (1)
TIMP-3	2 (2)	2 (2)	1 (1)	1 (1)
TIMP-4	1 (1–2)	1 (1–2)	1 (0–1)	1 (1)

Abs=antibodies; AB=anterior border; AE=anterior epithelium; IS=iris stroma; PostE=posterior epithelium.

coated slides. Sections were deparaffinised in xylene and rehydrated with ethanol. Slides were boiled in antigen retrieval solution (citrate buffer) for 3 minutes in a microwave. This procedure was repeated three times, followed by three washes in phosphate buffered saline (PBS). Sections were incubated in 3% BSA in PBS for 30 minutes to block non-specific binding. They were then incubated with mouse anti-human MMPs and TIMPs Abs (1:150 dilution in 3% BSA) overnight at 4°C. Sections were washed three times and incubated for 1 hour at room temperature with goat anti-mouse-Alexa 568 (10 µg/ml). After further washing, slides were mounted in Vectashield (Vector Laboratories, Burlingame, CA, USA) and sealed with clear nail lacquer.

Negative controls included incubation of sections with an isotype matched monoclonal antibody (mouse IgG₁) and omission of the primary antibody. As a positive control, human synovial tissue from a patient with rheumatoid arthritis was stained for MMP-1. To facilitate morphological orientation, adjacent sections were stained with haematoxylin and eosin and examined by light microscopy.

Fluorescent microscopy and grading

Slides were examined with an Olympus BX60 microscope equipped with a 100 W mercury burner for epifluorescence illumination and wide band interference filters for green excitation/red emission (520–550 nm band pass, 590 nm long pass). For both fluorescence and bright field microscopy, images were captured with a spot cooled colour digital camera (Diagnostic Instruments, Sterling Heights, MI, USA). The intensity of immunofluorescent staining in each area of each section was graded as follows: 0: no staining; 1: weak staining; 2: moderate staining; 3: strong staining. Assignment of a grade was based both on the intensity of the staining and on the magnification at which it could be visualised. Grading was performed by two independent observers masked as to the identity of the samples. Data are expressed as medians and interquartile ranges.

RESULTS

Staining for MMPs and TIMPs was apparent throughout the anterior uvea, although there was marked variation in the intensity of staining in different regions (Tables 1 and 2). Interestingly, except for TIMP-3, most MMPs and TIMPs showed similar patterns in their distribution. In general, the intensity of staining in the ciliary body was stronger than in the iris. In the ciliary body, the patterns of intensity of the staining was non-pigmented epithelium > ciliary muscle > pigmented epithelium > stromal cells for all the antibodies. Whereas staining for MMP-1, 2, 3, 9, and TIMP-1, 2, 4 was mainly in the cytoplasm of both the non-pigmented and pigmented epithelial cells (Fig 1C), staining for TIMP-3 was mostly in the basement membrane of the epithelium (Fig 1D).

Table 2 Distribution and intensity of immunoreactivity for MMPs and TIMPs in the ciliary body

Abs	Ciliary body (n=7)			
	NPE	CM	PE	CS
MMP-1	3 (3)	2 (1)	2 (2)	2 (2)
MMP-2	3 (3)	3 (2–3)	2 (1–2)	2 (1–2)
MMP-3	3 (2–3)	2 (2)	2 (1–2)	2 (2)
MMP-9	2 (2–3)	2 (2)	1 (1–2)	1 (1–2)
TIMP-1	2 (2)	2 (2)	1 (1)	1 (1)
TIMP-2	3 (2–3)	3 (2–3)	2 (2)	2 (1–2)
TIMP-3	3 (3)	2 (2)	2 (2)	2 (2)
TIMP-4	2 (2)	1 (1–2)	1 (1–2)	1 (1)

NPE=non-pigmented epithelium; CM=ciliary muscles; PE=pigmented epithelium; CS=stroma in ciliary body.

In the iris, staining intensity generally followed this pattern: anterior border > anterior epithelium > stromal cells > posterior epithelium for all the antibodies tested (Table 1, Fig 1A, B). Prominent staining of blood vessel walls was also noted. Figure 1A–D shows representative images of fluorescent staining for MMP-1 and TIMP-3 in the iris and ciliary body.

No staining was observed in the negative controls (Fig 1E left). However, there was intensive staining for MMP-1 in human rheumatoid synovial tissue that was used as a positive control (Fig 1E right).

DISCUSSION

To our knowledge this is the first report demonstrating the distribution of MMP-1, 2, 3, 9, and TIMPs 1–4 in the normal human iris and ciliary body. Because of the considerable quantity of melanin in the pigment epithelium in the iris and ciliary processes, conventional immunohistochemistry is not suitable for localisation of MMPs and TIMPs in these tissues.³² This may be one reason for the lack of detailed studies on the localisation of MMPs/TIMPs in the anterior uvea. The use of a red emitting dye successfully minimises the problem of autofluorescence in paraffin sections.^{31–33} In the present study, we used this immunofluorescence staining technique to overcome the problem, which allowed us to localise MMPs and TIMPs in anterior uvea in detail.

Our results revealed the widespread distribution of MMPs 1–3 and 9 and TIMPs 1–4 in the anterior uvea. Interestingly, the intensity of the staining of MMPs and TIMPs revealed a similar pattern in their distribution. For example, in the iris the strongest immunoreactivity was found in the anterior border, followed by the anterior epithelium and stromal cells, whereas the posterior epithelium was weakly reactive. The positive correlation in their distribution of MMPs and TIMPs is in agreement with previous reports indicating that the inhibition is stoichiometric—that is, one molecule of TIMP inhibits one molecule of active MMP.³² Furthermore, this suggests that the balance between MMPs and TIMPs may be important in maintaining homeostasis in the anterior uvea.

Physiological functions of MMPs include remodelling of the extracellular matrix, cell migration, proliferation, and apoptosis. Their function and specific roles may be regulated by their naturally occurring inhibitors. Although MMPs and TIMPs have been demonstrated in the eye and have the potential to impact on ocular physiology,^{35–37} relatively little is known about their function. Recent studies have suggested a role in the maintenance of normal aqueous humour outflow,³⁰ in maintaining vitreous integrity,³⁸ and on the ageing of the eye.³⁹

In ocular disease, cytokines such as TNF- α and IL-1 α , adhesion molecules, and growth factors such as fibroblast growth factor^{40–41} may alter the activity of MMPs or lead to an imbalance between MMPs and TIMPs.^{20–23}

Figure 1 Expression of MMP-1 and TIMP-3 in the iris and ciliary body. Paraffin sections of the anterior uvea were stained for MMP-1 or TIMP-3, and then visualised with fluorochrome-Alexa 568. (A) MMP-1 in the iris: anterior border (ab); iris stroma (is); anterior epithelium (ae); posterior epithelium (poste). (B) TIMP-3 in the iris: anterior border (ab); iris stroma (is); anterior epithelium (ae); posterior epithelium (poste). (C) MMP-1 in the ciliary body: positive staining mainly located in the cytoplasm of the epithelium, non-pigmented epithelium (npe); pigmented epithelium (pe); ciliary body stroma (cs). (D) TIMP-3 in the ciliary body: basement membrane of non-pigmented epithelium (bnpe); basement membrane of pigmented epithelium (bpe); ciliary body stroma (cs). Negative control (E, left): the primary Ab was omitted; positive control (E, right): rheumatoid synovial tissue stained for MMP-1.

Previous reports from our laboratory showed that elevated MMPs (1, 2, 3, 7, 9) were found in necrotising scleritis,²⁰ pterygial tissue,^{26–28} in the aqueous humour (AH) and infiltrating cells (macrophages, T lymphocytes and neutrophils) of patients with uveal inflammation and animals

Authors' affiliations

J Lan, RK Kumar, N Di Girolamo, P McCluskey, D Wakefield, The Inflammation Research Unit, School of Medical Sciences, Department of Pathology, University of New South Wales, Sydney 2052, Australia

REFERENCES

- 1 Woessner JF. Matrix metalloproteinases and their inhibitors in connective tissue remodeling. *FASEB J* 1991;5:2145–54.
- 2 McCawley LJ, Matrisian LM. Matrix metalloproteinases: multifunctional contributors to tumor progression. *Mol Med Today* 2000;6:149–56.
- 3 Vu TH, Werb Z. Matrix metalloproteinases: effectors of development and normal physiology. *Genes Dev* 2000;14:2123–33.
- 4 Tomita T, Nakase T, Kaneko M, et al. Expression of extracellular matrix metalloproteinase inducer and enhancement of the production of matrix metalloproteinases in rheumatoid arthritis. *Arthritis Rheum* 2002;46:373–8.
- 5 Hulboy DL, Rudolph LA, Matrisian LM. Matrix metalloproteinases as mediators of reproductive function. *Mol Hum Reprod* 1997;3:27–45.
- 6 Stahle-Backdahl M, Sandstedt B, Bruce K, et al. Collagenase-3 (MMP-13) is expressed during human fetal ossification and re-expressed in postnatal bone remodeling and in rheumatoid arthritis. *Lab Invest* 1997;6:717–28.
- 7 Kanemoto M, Hukuda S, Komiya Y, et al. Immunohistochemical study of matrix metalloproteinase-3 and tissue inhibitor of metalloproteinase-1 human intervertebral discs. *Spine* 1996;21:1–8.
- 8 Wong TL, Sethi C, Daniels JT, et al. Matrix metalloproteinases in disease and repair processes in the anterior segment. *Surv Ophthalmol* 2002;47:239–56.
- 9 Ribbens C, Porras MM, Franchimont N, et al. Increased matrix metalloproteinase-3 serum levels in rheumatic diseases: relationship with synovitis and steroid treatment. *Ann Rheum Dis* 2002;61:161–9.
- 10 Alenius GM, Jonsson S, Wallberg Jonsson S, et al. Matrix metalloproteinase 9 (MMP-9) in patients with psoriatic arthritis and rheumatoid arthritis. *Clin Exp Rheumatol* 2001;19:760–9.
- 11 Plantner JJ, Jiang C, Smine A. Increase in interphotoreceptor matrix gelatinase A (MMP-2) associated age-related macular degeneration. *Exp Eye Res* 1998;67:637–45.
- 12 Kubota T, Nomura T, Takahashi T, et al. Expression of mRNA for matrix metalloproteinases and tissue inhibitors of metalloproteinases in periodontitis-affected human gingival tissue. *Arch Oral Biol* 1996;41:252–62.

- 13 **Kahari VM**, Saarialho-kere U. Matrix metalloproteinases and their inhibitors in tumor growth and invasion (Review). *Ann Med* 1999;**31**:34–5.
- 14 **Elliott S**, Cawston T. The clinical potential of matrix metalloproteinase inhibitors in the rheumatic disorder. *Drugs Aging* 2001;**18**:87–99.
- 15 **Hamada T**, Arima N, Shindo M, *et al*. Suppression of adjuvant arthritis of rats by a novel matrix metalloproteinase-inhibitor. *Br J Pharmacol* 2000;**131**:1513–20.
- 16 **Sykes AP**, Bhogal R, Brampton C, *et al*. The effect of an inhibitor of matrix metalloproteinases on colonic inflammation in a trinitrobenzenesulphonic acid rat model of inflammatory bowel disease. *Aliment Pharmacol Ther* 1999;**13**:535–42.
- 17 **Mackay AR**, Ballin M, Pelina MD, *et al*. Effect of phorbol ester and cytokines on matrix metalloproteinase and tissue inhibitor of metalloproteinase expression in tumor and normal cell lines. *Invasion Metastasis* 1992;**12**:168–84.
- 18 **Hanemaaijer R**, Koolwijk P, le Clercq L, *et al*. Regulation of matrix metalloproteinase expression in human vein and microvascular endothelial cells. Effects of tumour necrosis factor alpha, interleukin 1 and phorbol ester. *Biochem J* 1993;**296**:803–9.
- 19 **Stetler-Stevenson WG**, Kruttsch HC, Liotta LA. Tissue inhibitor of metalloproteinase-2 (TIMP-2). A new member of the metalloproteinase inhibitor family. *J Biol Chem* 1989;**264**:17374–8.
- 20 **Di Girolamo N**, Lloyd A, McCluskey P, *et al*. Increased expression of matrix metalloproteinases in vivo in scleritis tissue and in vitro in cultured human scleral fibroblasts. *Am J Pathol* 1997;**150**:653–66.
- 21 **Smith VA**, Hoh HB, Easty DL. Role of ocular matrix metalloproteinases in peripheral ulcerative keratitis. *Br J Ophthalmol* 1999;**83**:1376–83.
- 22 **Tuter G**, Kurtis B, Serdar M. Effects of phase I periodontal treatment on gingival crevicular fluid levels of matrix metalloproteinase-1 and tissue inhibitor of metalloproteinase-1. *J Periodontol* 2002;**73**:487–93.
- 23 **Ishiguro N**, Ito T, Oguchi T, *et al*. Relationships of matrix metalloproteinases and their inhibitors to cartilage proteoglycan and collagen turnover and inflammation as revealed by analyses of synovial fluids from patients with rheumatoid arthritis. *Arthritis Rheum* 2001;**44**:2503–11.
- 24 **Di Girolamo N**, Verma MJ, McCluskey PJ, *et al*. Increased matrix metalloproteinases in the aqueous humor of patient and experimental animals with uveitis. *Curr Eye Res* 1996;**5**:1060–8.
- 25 **Cuello C**, Wakefield D, Di Girolamo N. Neutrophil accumulation correlates with type IV collagenases/gelatinase activity in endotoxin induced uveitis. *Br J Ophthalmol* 2002;**86**:290–5.
- 26 **Di Girolamo N**, McCluskey P, Lloyd A, *et al*. Expression of MMPs and TIMPs in human pterygia and cultured pterygium epithelial cells. *Invest Ophthalmol Vis Sci* 2000;**41**:671–9.
- 27 **Di Girolamo N**, Wakefield D, Coroneo MT. Differential expression of matrix metalloproteinases and their tissue inhibitors at the advancing pterygium head. *Invest Ophthalmol Vis Sci* 2000;**41**:4142–9.
- 28 **Di Girolamo N**, Coroneo MT, Wakefield D. Active matrilysin (MMP-7) in human pterygia: potential role in angiogenesis. *Invest Ophthalmol Vis Sci* 2001;**42**:963–8.
- 29 **Wallace GR**, Whiston RA, Stanford MR, *et al*. The matrix metalloproteinase inhibitor BB-1101 prevents experimental autoimmune uveoretinitis. *Clin Exp Immunol* 1999;**118**:364–70.
- 30 **Gaton DD**, Sagara T, Lindsey JD, *et al*. Increased matrix metalloproteinases 1, 2, and 3 in the monkey uveoscleral outflow pathway after topical prostaglandin F (2 alpha)-isopropylester treatment. *Arch Ophthalmol* 2001;**119**:1165–70.
- 31 **Kumar RK**, Walsh C, Cooke BE. Reduced nonspecific fluorescence of paraffin sections by use of a novel red-emitting dye. *J Histotechnol* 2000;**23**:99–102.
- 32 **Klisovic DD**, O'Dorisio MS, Katz SE, *et al*. Somatostatin receptor gene expression in human ocular tissues: RT-PCR and immunohistochemical study. *Invest Ophthalmol Vis Sci* 2001;**42**:2193–201.
- 33 **Wang N**, Lindsey JD, Angert M, *et al*. Latanoprost and matrix metalloproteinase-1 in human choroids organ cultures. *Curr Eye Res* 2001;**22**:198–207.
- 34 **Nagase H**, Suzuki K, Cawston TE, *et al*. Involvement of a region near valine-69 of tissue inhibitor of metalloproteinases (TIMP)-1 in the interaction with matrix metalloproteinase 3. *Biochem J* 1997;**35**:163–7.
- 35 **Sivak JM**, Fini ME. MMPs in the eye: emerging roles for matrix metalloproteinases in ocular physiology. *Prog Retin Eye Res* 2002;**21**:1–14. (Review.)
- 36 **Fini ME**, Girard MT. Expression of collagenolytic/gelatinolytic metalloproteinases by normal cornea. *Invest Ophthalmol Vis Sci* 1990;**31**:1779–88.
- 37 **Fariss RN**, Apte SS, Olsen BR. Tissue inhibitor of metalloproteinases-3 is a component of Bruch's membrane of the eye. *Am J Pathol* 1997;**150**:323–8.
- 38 **Brown D**, Hamdi H, Bahri S, *et al*. Characterization of an endogenous metalloproteinase in human vitreous. *Curr Eye Res* 1994;**13**:639–47.
- 39 **Kamei M**, Hollyfield JG. TIMP-3 in Bruch's membrane: changes during aging and in age-related macular degeneration. *Invest Ophthalmol Vis Sci* 1999;**40**:2367–75.
- 40 **Fini ME**, Strissel KJ, Girard MT, *et al*. Interleukin 1 alpha mediates collagenase synthesis stimulated by phorbol 12-myristate 13-acetate. *J Biol Chem* 1994;**269**:11291–8.
- 41 **Mohan R**, Sivak J, Ashton P, *et al*. Curcuminoids inhibit the angiogenic response stimulated by fibroblast growth factor-2, including expression of matrix metalloproteinase gelatinase B. *J Biol Chem* 2000;**275**:10405–12.