Unilateral visual impairment and health related quality of life: the Blue Mountains Eye Study

E-M Chia, P Mitchell, E Rochtchina, S Foran, J J Wang

Br J Ophthalmol 2003;**87**:392–395

See end of article for authors' affiliations

Correspondence to:
Paul Mitchell, MD, PhD,
University of Sydney
Department of
Ophthalmology (Centre for
Vision Research,
Westmead Hospital),
Hawkesbury Road,
Westmead, NSW,
Australia, 2145;
paul_mitchell@
wmi.usyd.edu.au

Accepted for publication
6 January 2003

Background: There has been considerable recent interest in the impact of unilateral visual impairment on functional status and wellbeing, particularly in relation to second eye cataract surgery.

Aim: To determine if unilateral visual impairment has a measurable impact on health related quality of life (HRQOL) in an older community, as assessed by the generic, multidimensional 36 item short form health survey (SF-36).

Methods: All participants of the second cross sectional Blue Mountains Eye Study (n = 3508) were invited to attend comprehensive eye examinations and complete an SF-36 questionnaire. Unilateral visual impairment was defined as visual acuity (VA) <6/12 in the worse eye and ≥6/12 in the better eye. Mild visual impairment was defined as VA <6/12 but ≥6/24, moderate as VA <6/24 but ≥6/60, and severe (blindness) as VA <6/60 in the worse eye. Cases with amblyopia (n = 48) were excluded.

Results: Complete data were available for 3108 participants; 227 (7.3%) had unilateral visual impairment (148 mild, 29 moderate, 50 severe). Moderate to severe non-correctable unilateral impairment was associated with poorer SF-36 profiles. After adjusting for age and sex, this group had significantly poorer scores than the unimpaired group in three of eight domains (p<0.05); limitations as a result of physical problems, social function, limitations because of emotional problems, and in the mental component score. Mental domains were more affected than physical domains. Unilateral impairment from undercorrected refraction did not measurably affect HRQOL.

Conclusions: Moderate to severe non-correctable unilateral visual impairment caused by eye diseases such as cataract had a measurable impact on HRQOL.

The impact of bilateral visual impairment on health related quality of life (HRQOL) and independent living has been well documented¹ including an increased mortality risk,^{2–4} earlier nursing home placement,⁵ falls,⁶ and increased use of community support services.⁷ The impact of unilateral visual impairment on functional status and wellbeing is more subtle and less well understood. Its impact, however, has been generating increasing interest in recent years, particularly in the context of second eye cataract surgery.⁸

The estimated cost of cataract surgery in the United States in 1991 was US\$3.4 billion.⁹ In the United Kingdom, it has been estimated that approximately one third of cataract surgery is performed on second eyes.¹⁰ Owing to constraints on healthcare budgets, there have been suggestions that public funding for second eye cataract surgery could be rationed. Many recent studies have indicated that this would be a detrimental move which could adversely affect functional status, independence, and HRQOL.^{11–20}

The Blue Mountains Eye Study (BMES) is a well described cohort study of vision and common eye diseases. We aimed in this population sample to determine whether unilateral visual impairment had a measurable impact on HRQOL, as assessed by the generic, multidimensional 36 item short form health survey (SF-36).²¹ This instrument is a validated health outcome measure that has been widely used to assess the impacts on quality of life in many different medical conditions.^{22–26}

PATIENTS AND METHODS

The BMES is a well described population based survey in an urban population aged 49 years or older resident in two post-code areas of the Blue Mountains region, west of Sydney, Australia. Methods used in this survey have already been

described.²⁷ The present report uses findings from the second cross section of this population.

BMES I identified 4433 eligible non-institutionalised permanent residents in a door to door census conducted during 1991, of whom 3654 (82.4%) participated in the examinations during 1992–4. After 5 years, BMES I participants were invited to attend repeat examinations during 1997–9 (BMES IIA). By the time these commenced, 543 people had died (14.9%) and 2335 (75.1%) were re-examined. A repeat door to door census conducted in 1999 of the same area identified 1378 additional eligible permanent residents who had moved into the area or were now aged 50 or older. We examined 1174 (85.2%) of this group during 1999–2000 (BMES IIB). Cross section 2 thus comprised 3509 participants from BMES IIA and IIB.

Participants were invited to attend a comprehensive eye examination that included measurement of distance logMAR visual acuity, before and after subjective refraction using the ETDRS method,²⁷ automated perimetry, and lens and stereo retinal photography.

Before attending, we asked participants to complete a detailed questionnaire, which was mailed after appointments were made. Included were questions about visual function and diet, and the SF-36 quality of life, self administered questionnaire.²¹ This instrument contains 36 items, measuring eight dimensions of health and wellbeing: “physical functioning,” “role limitation due to physical problems,” “bodily pain,” “general health perceptions,” “vitality,” “social functioning,” “role limitation due to emotional problems,” and “mental health.” Each dimension was scored from 0 (worst possible health state) to 100 (best possible health state) by coding, summing, and transforming relevant item scores. Missing values were substituted by averaging other items if the number of missing items was less than half the total number

Figure 1 Flow chart showing number of people with bilateral and unilateral visual impairment (VI), based on presenting and best corrected visual acuity: the Blue Mountains Eye Study second cross section.

Figure 2 Sex adjusted mean SF-36 scores in the Blue Mountains Eye Study (second cross section) by age group.

of items in that dimension. No substitution was used for dimensions with only two items. The eight dimensions comprise two domains "physical" and "mental," scored as physical component scores (PCS) and mental component scores (MCS). These are summary measures calculated using factor analysis and Australian normalised scores.²⁸

Unilateral visual impairment was defined as visual acuity (VA) $<6/12$ in the worse eye and $\geq 6/12$ in the better eye. Mild visual impairment was defined as VA $<6/12$ but $\geq 6/24$, moderate impairment as VA $<6/24$ but $\geq 6/60$, and severe impairment (blindness) as VA $<6/60$. People with unilateral impairment attributed to amblyopia were excluded, as their inclusion would have diluted assessment of the impact of changes in unilateral visual acuity. The impact from unilateral visual impairment which was correctable by refraction was also assessed.

Figure 3 Age and sex adjusted mean SF-36 scores in the Blue Mountains Eye Study (second cross section) with moderate to severe ($n = 79$), mild ($n = 148$), and no ($n = 2822$) non-correctable unilateral visual impairment (VI).

Statistical Analysis System (sas, version 6.12 for Windows; SAS Institute, Cary, NY, USA) was used for statistical analyses. Age and sex adjusted values are presented.

RESULTS

Participants had a mean age of 66.7 years (range 49–98 years) and 57% were female. Of the 3508 participants in the second BMES cross section, complete data sets that included SF-36 questionnaires were available for 3153 participants (89.9%). Bilateral visual impairment, based on presenting VA, was present in 237 participants (7.5%), unilateral impairment was present in 543 subjects (17.2%), and 2373 (75.3%) had no visual impairment, as shown in Figure 1. After refraction, 67 participants (2.1%) were still visually impaired in both eyes, while 264 (8.4%) remained impaired in one eye only. After further exclusion of subjects with amblyopia, there were 497 people (16.0%) with unilateral visual impairment at presentation and 227 (7.3%), after refraction. Of these 227 subjects, 148 participants had mild and 29 had moderate unilateral visual impairment while 50 were unilaterally blind. Owing to the relatively small numbers, cases with moderate and severe unilateral visual impairment were pooled for the analysis of SF-36 profiles.

Increasing age was strongly associated with poorer SF-36 profiles. This worsening age related trend was demonstrated for all SF-36 dimensions (Fig 2) and for the two component scores (data not shown; $p < 0.001$). After adjusting for age, women had slightly poorer SF-36 profiles than men (data not shown). This sex difference, however, was statistically significant for all dimensions ($p < 0.05$), apart from "general health" ($p = 0.41$) and "role limitation due to emotional problems" ($p = 0.14$).

Table 1 shows that there were no overall differences in mean age and sex adjusted SF-36 scores between people with and without unilateral visual impairment. Unilateral visual impairment correctable by refraction had no measurable impact

Table 1 Mean age and sex adjusted SF-36 scores of Blue Mountains Eye Study participants with non-correctable, correctable, and no unilateral visual impairment (VI)

Unilateral visual impairment	Physical function	Role limitation physical	Bodily pain	General health	Vitality	Social function	Role limitation emotional	Mental health	Physical component score	Mental component score
None	73.3	68.7	70.6	68.3	62.4	84.2	81.5	78.5	45.4	51.9
Correctable	72.0	67.1	71.5	68.7	62.6	84.7	81.7	79.0	45.1	52.2
Non-correctable	73.5	62.6	72.7	67.2	62.1	84.4	77.9	77.8	45.4	51.5
p Value*	0.9264	0.0421	0.2713	0.4897	0.8684	0.9168	0.1709	0.5971	0.9919	0.6464
p For trend	0.6301	0.1171	0.4887	0.7292	0.9602	0.9417	0.3809	0.7399	0.8701	0.7475

*Non-correctable compared with no visual impairment.

Table 2 Mean age and sex adjusted SF-36 scores of Blue Mountains Eye Study participants with moderate to severe, mild, and no non-correctable unilateral visual impairment (VI)

Non-correctable unilateral VI	Physical function	Role limit physical	Bodily pain	General health	Vitality	Social function	Role limit emotional	Mental health	Physical component score	Mental component score
None	72.6	67.9	70.7	68.3	62.3	84.1	81.3	78.6	45.2	52.0
Mild	70.4	64.7	72.5	66.8	61.6	81.2	77.5	77.7	45.2	51.1
Moderate to severe	70.3	57.7	67.1	63.9	60.1	78.2	71.0	75.5	43.5	49.5
p Value*	0.3932	0.0317	0.2286	0.0920	0.3774	0.0340	0.0126	0.1283	0.1827	0.0495
p For trend	0.4072	0.0744	0.3228	0.1937	0.6431	0.0483	0.0260	0.2839	0.4105	0.1090

*Moderate to severe non-correctable VI compared with no non-correctable VI.

on the mean scores. Unilateral non-correctable impairment modestly affected one dimension; "role limitation due to physical problems" ($p = 0.04$), but there was no impact on overall physical or mental summary scores. Stratification of the non-correctable group, however, demonstrated a gradient with poorer SF-36 profiles seen for increasing severity of unilateral visual impairment, as shown in Figure 3. Table 2 shows that, while relatively modest, the impact was statistically significant only for people with moderate to severe impairment. After adjusting for age and sex, this group had significantly poorer mean scores than people with no visual impairment in three of eight dimensions ($p < 0.05$); "role limitation due to physical problems" (15% lower), "social functioning" (7% lower), "role limitation due to emotional problems" (13% lower), and in the mental component score. There was a greater effect on the mental domain than the physical domain.

DISCUSSION

In the past decade, there has been considerable interest in examining the effects of unilateral visual impairment, particularly in relation to second eye cataract surgery in the context of constrained healthcare expenditure.⁸ Our study demonstrates a modest, but statistically significant impact on a general health measure of quality of life and independent functioning in the presence of moderate to severe non-correctable unilateral visual impairment. This was present in only three of the eight SF-36 domains, after age-sex adjustment and only for moderate to severe impairment. There was no measurable impact from correctable unilateral impairment.

Our data confirm findings from many recent reports that have demonstrated a range of benefits from second eye cataract surgery.^{8, 13, 14, 16, 20} These have included improvements in subjective visual function,²⁰ binocular and stereo acuity,^{13, 14} and better performance on questionnaire instruments that target general functioning (for example, the Sickness Impact Profile¹⁶), eye specific functioning (for example, the Visual Function 14 item questionnaire^{8, 16}), or quality of life.¹³

In a randomised trial of the effectiveness of second eye cataract surgery, Laidlaw *et al*¹³ reported an 11%–30% difference in visual symptoms and quality of life measures between routine and expedited surgery. Javitt *et al*²⁰ showed that patients with bilateral cataract induced visual impairment who underwent cataract surgery in both eyes had significantly improved subjective visual function than those with unilateral cataract surgery alone. Talbot and Perkins¹⁴ showed that second eye cataract surgery increased the proportion of older people who satisfied driving licence requirements from 52% to 86%. Guthauser and Flammer²⁹ reported a subgroup of patients with central visual field impairment who had the greatest threshold for recovery after cataract surgery.

The functional improvement in vision from second eye cataract surgery could be explained by better monocular and binocular contrast sensitivity.^{14, 19} Talbot and Perkins¹⁴ also showed that 66% of their participants had improved binocular

visual acuity and stereoacuity from 32% to 90%, previous ocular dominance was restored in 44%, and there were improvements to binocular horizontal and vertical field of vision. Laidlaw and Harrad¹⁹ have also stated that glare reduction may also play a part, with a reduction from 86% to 41%, in the proportion of patients with symptomatic glare after second eye cataract surgery, despite previous reports of increased glare associated with posterior chamber lens implants.³⁰

There are well described objective improvements to visual acuity following second eye cataract surgery in terms of stereopsis and visual acuity that can be explained by binocular summation. Objective tests, however, have not demonstrated significant binocular inhibition from second eye cataract.¹⁹ Thus, the disability resulting from visual impairment does not stem from the loss of visual acuity or subjective aspects of visual function such as glare alone. Both visual^{8, 31} and functional^{8, 13, 19} effects need to be addressed independently. Our study has attempted to examine the impacts of unilateral visual impairment on HRQOL, as measured by the SF-36, an instrument that includes both physical and mental domains. We excluded cases where unilateral visual impairment was due to amblyopia, as their inclusion could dilute our findings and this disorder has uncertain public health importance.

Data from the United Kingdom indicate that cataract surgery will improve objective vision with minimal complications when well established guidelines are used.³² Most past studies have looked at objective changes in vision such as binocular acuity, contrast sensitivity, and stereopsis. The trial reported by Laidlaw *et al*¹³ showed benefits from second eye cataract surgery in reducing self reported difficulties, despite only marginal improvements in binocular visual acuity. A large observational UK study of cataract surgery showed significant gains in health and vision related quality of life outcome measures from first eye surgery with additional significant gains at 1 year, if second eye surgery was also performed.¹⁶

The SF-36 questionnaire has been shown to discriminate well between Australians with and without various health conditions, in a validity study conducted on 555 subjects in a National Heart Foundation survey.³³ Our age and sex specific SF-36 findings (in part, shown in Fig 2) are also very close to the SF-36 norms for a much larger population sample, provided by the 1995 Australian Bureau of Statistics National Health Survey.²⁸ The number of domains with significant age-sex adjusted differences between people with and without moderate to severe unilateral non-correctable visual impairment (three of eight domains, plus the mental component score) is similar to findings from a number of recent reports using this instrument.³⁴ In our study, moderate to severe bilateral non-correctable visual impairment was associated with a substantially greater impact than unilateral impairment, and was of a magnitude similar to that from major systemic diseases, such as heart disease, stroke, diabetes, and cancer (data not shown). Changes in SF-36 scores that differ between groups by two or more points on the scale of 0 to 100 have been shown to be clinically or socially meaningful.³⁵ In our study, at least this magnitude of difference was demonstrated

for all eight domains (in Table 2). Importantly, the completion rate (89.9%) of the SF-36 questionnaire by participants in our study is substantially higher than in most other reports using the self administered version, mailed to respondents (for example, 79.2%³⁶).

Our study has a number of limitations, including: (1) insufficient power to examine the impacts on HRQOL from the different causes of visual impairment (for example, cataract versus macular degeneration) and (2) lack of concurrent comparison with questionnaires designed to measure visual functioning. The strengths are its population base and relatively high completion rate (89.9%) by study participants.

In summary, this study has demonstrated modest, but statistically significant detrimental effects on a general measure of HRQOL associated with unilateral non-correctable visual impairment in an older population sample. Importantly, this was only found for moderate to severe levels of visual impairment and for three of the eight SF-36 dimensions, plus the mental component score. Nevertheless, our study highlights the potential for significant morbidity from unilateral eye diseases, such as unoperated cataract, particularly in the psychosocial domain, and the potential for improvement with treatment. These data provide support for the notion that cataract surgery on the second eye of patients with bilateral cataract is likely to provide significant and measurable functional benefits.

ACKNOWLEDGEMENT

This study was supported by the Australian National Health and Medical Research Council (grant Nos 974159 and 991407) and the Westmead Millennium and Save Sight Institutes, University of Sydney.

Authors' affiliations

E-M Chia, P Mitchell, E Rochtchina, S Foran, J J Wang, Department of Ophthalmology, University of Sydney, Sydney, Australia

REFERENCES

- 1 Wang JJ, Mitchell P, Smith W. Vision and low self-rated health: the Blue Mountains Eye Study. *Invest Ophthalmol Vis Sci* 2000;**41**:49–54.
- 2 Wang JJ, Mitchell P, Simpson JM, et al. Visual impairment, age-related cataract, and mortality. *Arch Ophthalmol* 2001;**119**:1186–90.
- 3 Taylor HR, McCarty CA, Nanjan MB. Vision impairment predicts five-year mortality. *Trans Am Ophthalmol Soc* 2000;**98**:91–6.
- 4 Lee DJ, Gomez-Marín O, Lam BL, et al. Visual acuity impairment and mortality in US adults. *Arch Ophthalmol* 2002;**120**:1544–5.
- 5 Wang JJ, Mitchell P, Cumming RG, et al. Visual impairment and nursing home placement in older Australians: the Blue Mountains Eye Study. *Ophthalm Epidemiol* 2003 (in press).
- 6 Ivers RQ, Cumming RG, Mitchell P, et al. Visual impairment and falls in older adults: the Blue Mountains Eye Study. *J Am Geriatr Soc* 1998;**46**:58–64.
- 7 Wang JJ, Mitchell P, Smith W, et al. Impact of visual impairment on use of community support services by elderly persons: the Blue Mountains Eye Study. *Invest Ophthalmol Vis Sci* 1999;**40**:12–19.
- 8 Javitt JC, Steinberg EP, Sharkey P, et al. Cataract surgery in one eye or both. A billion dollar per year issue. *Ophthalmology* 1995;**102**:1583–92.
- 9 Steinberg EP, Javitt JC, Sharkey PD, et al. The content and cost of cataract surgery. *Arch Ophthalmol* 1993;**111**:1041–9.
- 10 Courtney P. The National Cataract Surgery Survey: I. Method and descriptive features. *Eye* 1992;**6**:487–92.
- 11 Lundström M, Stenevi U, Thorburn W. Quality of life after first and second-eye cataract surgery: five-year data collected by the Swedish National Cataract Register. *J Cataract Refract Surg* 2001;**27**:1553–9.
- 12 Castells X, Alonso J, Ribo C, et al. Comparison of the results of first and second cataract eye surgery. *Ophthalmology* 1999;**106**:676–82.
- 13 Laidlaw DA, Harrod RA, Hopper CD, et al. Randomised trial of effectiveness of second eye cataract surgery. *Lancet* 1998;**352**:925–9.
- 14 Talbot EM, Perkins A. The benefit of second eye cataract surgery. *Eye* 1998;**12**(Pt 6):983–9.
- 15 Elliott DB, Patla A, Bullimore MA. Improvements in clinical and functional vision and perceived visual disability after first and second eye cataract surgery. *Br J Ophthalmol* 1997;**81**:889–95.
- 16 Desai P, Reidy A, Minassian DC, et al. Gains from cataract surgery: visual function and quality of life. *Br J Ophthalmol* 1996;**80**:868–73.
- 17 Mizener JB, Podhajsky P, Hayreh SS. Ocular ischemic syndrome. *Ophthalmology* 1997;**104**:859–64.
- 18 Claridge KG, Francis PJ, Bates AK. Should second eye cataract surgery be rationed? *Eye* 1995;**9**:47–9.
- 19 Laidlaw A, Harrod R. Can second eye cataract extraction be justified? *Eye* 1993;**7**:680–6.
- 20 Javitt JC, Brenner MH, Curbow B, et al. Outcomes of cataract surgery. Improvement in visual acuity and subjective visual function after surgery in the first, second, and both eyes. *Arch Ophthalmol* 1993;**111**:686–91.
- 21 Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;**30**:473–83.
- 22 Hackett ML, Duncan JR, Anderson CS, et al. Health-related quality of life among long-term survivors of stroke: results from the Auckland Stroke Study, 1991–1992. *Stroke* 2000;**31**:440–7.
- 23 Wu SY, Sainfort F, Tomar RH, et al. Development and application of a model to estimate the impact of type 1 diabetes on health-related quality of life. *Diabetes Care* 1998;**21**:725–31.
- 24 Kinney SE, Sandridge SA, Newman CW. Long-term effects of Meniere's disease on hearing and quality of life. *Am J Otol* 1997;**18**:67–73.
- 25 Lawrence WF, Fryback DG, Martin PA, et al. Health status and hypertension: a population-based study. *J Clin Epidemiol* 1996;**49**:1239–45.
- 26 Solomon GD, Skobieranda FG, Genzen JR. Quality of life assessment among migraine patients treated with sumatriptan. *Headache* 1995;**35**:449–54.
- 27 Attebo K, Mitchell P, Smith W. Visual acuity and the causes of visual loss in Australia. The Blue Mountains Eye Study. *Ophthalmology* 1996;**103**:357–64.
- 28 Australian Bureau of Statistics. *National health survey: SF-36 population norms*. Canberra: ABS, 1997:4399.0.
- 29 Guthauser U, Flammer J. Quantifying visual field damage caused by cataract. *Am J Ophthalmol* 1988;**106**:480–4.
- 30 Nadler DJ, Jaffe NS, Clayman HM, et al. Glare disability in eyes with intraocular lenses. *Am J Ophthalmol* 1984;**97**:43–7.
- 31 D'Ambrosio FA Jr. Assessing disability in the patient with cataracts. *Curr Opin Ophthalmol* 1999;**10**:42–5.
- 32 Desai P, Minassian DC, Reidy A. National cataract surgery survey 1997–8: a report of the results of the clinical outcomes. *Br J Ophthalmol* 1999;**83**:1336–40.
- 33 McCallum J. The new "SF-36" health status measure: Australian validity tests. 1–12. ACT: National Centre for Epidemiology and Population Health, 1994.
- 34 Cohen HJ, Feussner JR, Weinberger M, et al. A controlled trial of inpatient and outpatient geriatric evaluation and management. *N Engl J Med* 2002;**346**:905–12.
- 35 Bacon CG, Giovannucci, Testa M, et al. The association of treatment-related symptoms with quality-of-life outcomes for localized prostate carcinoma patients. *Cancer* 2002;**94**:862–71.
- 36 McHorney CA, Kosinski M, Ware JE Jr. Comparisons of the costs and quality of norms for the SF-36 health survey collected by mail versus telephone interview: results from a national survey. *Med Care* 1994;**32**:551–67.