SCIENTIFIC REPORT

The management of orbital cysts associated with congenital microphthalmos and anophthalmos

C J McLean, N K Ragge, R B Jones, J R O Collin

Br J Ophthalmol 2003;87:860–863

Aims: To study the management of the orbital cysts present in a group of patients with anophthalmos and microphthalmos.

Methods: A retrospective study of 34 patients (40 orbits) treated for orbital cyst associated with microphthalmos and anophthalmos.

Results: The two largest treatment groups comprised 17 orbits (42.5%) where the cyst was removed surgically and 17 orbits (42.5%) where the cyst was retained and conformers were used. The remaining cases comprised two orbits (5%) where the cyst was aspirated initially; two orbits (5%) with large cysts which will need to be excised after further orbital growth; one orbit (2.5%) in which a silicone expander was used initially, and one orbit (2.5%) in which a mildly microphthalmic eye had some vision and was monitored but required no surgery.

Conclusion: In this study 33 out of 34 patients had a good cosmetic result which illustrates that the orbital cyst in microphthalmos or anophthalmos performs a useful role in socket expansion and that the majority of patients with this condition can expect a good cosmetic outcome.

Congenital microphthalmos and anophthalmos are rare conditions with prevalence rates of 1.4–3.5 per 10 000 births for microphthalmos^{1–3} and 0.3–0.6 per 10 000 births for anophthalmos.^{1–4} In one study looking at both anophthalmos and microphthalmos the combined prevalence rate was 1.0 per 10 000 births.⁵ In one large study⁶ of 29 patients with anophthalmos and 48 with microphthalmos, two patients had bilateral microphthalmos and cyst and one had unilateral microphthalmos and cyst. There is no agreed management of the cyst which may accompany these conditions.

In this retrospective analysis of 34 patients treated for an orbital cyst associated with microphthalmos and anophthalmos we describe the management strategies in our hospital.

METHODS

All patients with either microphthalmos and cyst or anophthalmos and cyst seen at Moorfields Eye Hospital between 1970 and 1998 were included in this study.

The patients were examined by the same surgeon and a consultant paediatrician in a specially designated anophthalmic clinic. Patients underwent imaging, most commonly computed tomograph (CT) or magnetic resonance imaging (MRI).

Once a cyst had been diagnosed, the aim of the subsequent management was to keep the cyst for as long as necessary to ensure adequate socket size. If the socket became too large the cyst was removed immediately. If the socket was growing satisfactorily with the cyst, it was retained until the optimum socket size was reached. If the cyst was not causing problems and a satisfactory artificial eye could be fitted, the cyst was retained. If an artificial eye could not be fitted without removing the cyst, the cyst was removed once an adequate socket

Figure 1 A 7 year old patient (case number 9) with left microphthalmos and a large orbital cyst which was removed to improve cosmesis, (A) before surgery and (B) after surgery with a new ocular prosthesis.

size had been achieved. Lid surgery was only carried out at an early stage if a conformer or artificial eye could not be retained. Elective lid surgery was otherwise left until after puberty and the patients were discharged in their mid-teens, when orbital growth was complete, if no other orbital or lid surgery was planned. They continued to have annual prosthetic appointments.

RESULTS

Thirty four patients were treated for orbital cyst associated with anophthalmos or microphthalmos (Table 1). In six of these cases the cysts were bilateral (three cases of anophthalmos and three cases of microphthalmos) giving a total of 40 orbital cysts. Fifteen patients were male and 19 were female. The mean age at presentation was 23 months (range 1–204 months). Of the unilateral orbital cysts 17 were on the left side and 11 were on the right side. Seven patients also had significant systemic abnormalities (Table 1).

The cases fell into six treatment groups: (A) the cyst was aspirated but fluid reaccumulation led to surgical excision; (B) the cyst was excised surgically; (C) conformer therapy only; (D) large orbital cyst preventing conformer use—future cyst excision planned; (E) mildly microphthalmic eye with some vision—no conformer used; (F) in this patient a silicone orbital expander was used initially followed by conformer treatment. The two largest treatment groups were 15 patients (17 orbits) requiring surgical excision of an orbital cyst (group B) and 17 patients treated with conformers only (group C). The orbital cysts were excised at a mean age of 41 months (range 6–128 months). In two cases the orbital cysts were initially aspirated, but owing to reaccumulation of fluid, the cysts required surgical excision. A large orbital cyst was present in two cases, which prevented the use of a conformer. These children are due to have cyst excision when they are older.

Table 1 Analysis of the 34 patients in this study and the presentation and management of their orbital cysts

Number	Age at presentation (months)	Sex	Diagnosis	Side with cyst	Systemic abnormalities	Ocular abnormalities	Primary treatment	Age at time of primary treatment (months)	Secondary lid or socket surgery	Cosmetic outcome	Follow up (months)
1	2	OM	UMC	Right	Nil	Nil	B	44	Nil	Good	78
2	6	F	BMUC	Left	Nil	Right microphthalmos	C	6	Nil	Good	27
3	24	F	MAC	Left	Oculo-cerebro-cutaneous syndrome	Right microphthalmos	A	3	MM LUF	Good	108
4	6	M	BMBC	Bilateral	Haemangiomatous branchial clefts	Nil	B (Right)	36	Nil	Good	144
5	72	F	BMBC	Bilateral	Nil	Nil	B (Left) B (Right) C (Left)	72	Nil	Good	108
6	6	M	BMUC	Left	Nil	Right colobomatous optic disc	C	6	Nil	Good	64
7	3	F	UMC	Left	Atrial septal defect	Left colobomatous iris and optic disc	C	3	Nil	Good	72
8	30	F	BAUC	Right	Nil	Nil	B	33	Nil	Good	108
9	84	F	UMC	Left	Nil	Right iris and retinal colobomas	B	85	Nil	Good	60
10	122	F	UMC	Right	Nil	Nil	B	128	Nil	Good	12
11	7	M	UMC	Left	Nil	Nil	C	7	Nil	Good	10
12	9	M	BMUC	Right	Aortic coarctation	Left microphthalmos	B	11	Nil	Good	6
13	12	M	BABC	Bilateral	Nil	Nil	F (Right) B (Left)	36	MM LUF	Good	50
14	11	M	UAC	Left	Nil	Nil	B	26	Nil	Good	120
15	72	M	UMC	Left	First Arch Syndrome	Nil	B	72	MM LLF	Good	18
16	4	F	BMBC	Bilateral	Nil	Nil	C (Right) A (Left)	24	Nil	Good	25
17	3	M	UMC	Left	Nil	Nil	C	3	Nil	Good	96
18	50	F	MAC	Left	Developmental delay	Right microphthalmos with colobomatous iris and retina	E (Right) D (Left)	–	Nil	Good	12
19	6	M	UMC	Left	Nil	Nil	B	20	Nil	Good	38
20	3	F	BAUC	Left	Nil	Nil	C	3	Nil	Fair	18
21	1	F	UMC	Right	Gastroschisis, cleft lip	Right lower lid coloboma	C	1	Lid coloboma repaired at 7 months	Good	86
22	1	F	UAC	Right	Cleft lip and palate, atrial septal defect, choanal atresia	Nil	C	1	Nil	Good	7
23	6	M	UAC	Left	Nil	Nil	C	6	Nil	Good	8
24	1	F	UMC	Left	Nil	Nil	D	–	Nil	Good	15
25	2	F	MAC	Right	Nil	Left retinal coloboma	C	2	Nil	Good	19
26	204	M	UAC	Left	Nil	Right congenital nystagmus	C	6	Nil	Good	132
27	1	F	BMUC	Right	Nil	Left macular coloboma	B	12	Nil	Good	396
28	2	F	BABC	Bilateral	Congenital hip dislocation	Nil	B (Right) B (Left)	33	Nil	Good	50
29	2	M	BABC	Bilateral	Nil	Nil	C	2	Nil	Good	57
30	9	F	UMC	Left	Nil	Nil	C	9	Nil	Good	56
31	1	M	BAUC	Right	Fistula in ano	Nil	C	1	Nil	Good	43
32	4	F	UMC	Left	Nil	Nil	B	8	Nil	Good	169
33	2	M	UAC	Right	Nil	Nil	B	6	Nil	Good	63
34	2	F	BMUC	Right	Nil	Nil	C	2	Nil	Good	124

UAC = unilateral anophthalmos with cyst; UMC = unilateral microphthalmos with cyst; BMBC = bilateral microphthalmos with bilateral cysts; MAC = microphthalmos (no cyst) and anophthalmos with cyst; BABC = bilateral anophthalmos with bilateral cysts; BMUC = bilateral microphthalmos with unilateral cyst; BAUC = bilateral anophthalmos with unilateral cyst; A = cyst aspirated but required surgical excision later; B = cyst excised surgically; C = conformer therapy only; D = large cyst preventing conformer fitting - for future excision; E = mildly microphthalmic eye with vision - no conformer used; F = silicone expander used initially, cyst later excised surgically; MM LLF = mucous membrane graft left lower fornix; MM LUF = mucous membrane graft left upper fornix.

A good cosmetic result was achieved in 33 out of 34 patients (see Fig 1). This was based on a subjective assessment by patient, parents, paediatrician and surgeon. One patient was judged to have a fair cosmetic result. No patients had a poor cosmetic result.

DISCUSSION

Cysts associated with microphthalmos and anophthalmos may present in a characteristic fashion. However, as they represent two points on the spectrum of colobomatous eye disorders, clinical distinction may be difficult and discernible only upon postmortem examination.⁷

Anophthalmos with cyst often presents at birth as a bluish orbital mass which pushes the upper lid forward.⁸⁻⁹ The cyst, which varies in size, represents the primary optic vesicle and histologically it may be solid⁸⁻⁹ or cystic¹⁰ and a maldeveloped lens may be present.¹⁰ Anophthalmos with cyst is usually associated with a normal eye in the contralateral orbit,⁸⁻⁹ although there have been cases associated with microphthalmos and cyst in the fellow orbit,¹¹ such as cases 3, 18, and 25 in this study.

Duke Elder¹² described three categories of cyst associated with microphthalmos: a relatively normal eye with a small cyst, not apparent clinically; an obvious cyst associated with a grossly deformed eye; a large cyst which has pushed the globe backwards so that it is not visible clinically. The cyst usually presents as a bluish bulge in the lower eyelid and is firm on palpation. If the cyst is visible through the conjunctiva it transilluminates readily. There have been a few reports where the cyst has presented as a bulge at the upper eyelid.¹¹⁻¹³ Most cases are diagnosed at birth although late presentation is possible.¹⁴

Both microphthalmos and anophthalmos can be associated with systemic abnormalities.¹⁵⁻¹⁸ We found systemic abnormalities associated with microphthalmos were probably no more common in cases with bilateral cysts compared to cases with unilateral cysts.

The diagnosis of orbital cyst associated with microphthalmos and anophthalmos can be made clinically in some cases, such as a moderately microphthalmic eye associated with a large orbital cyst pressing on the lower lid. However, in many cases some doubt often remains as to the nature of the orbital lesion and whether there is a microphthalmic eye or ocular remnant present hidden behind a large cyst. In these cases additional imaging has been valuable. Orbital ultrasound (both A¹¹ and B¹⁸⁻²⁰ modes) have been used to identify orbital cysts. More recently both CT¹⁴⁻¹⁹ and MRI²¹⁻²² scanning have proved invaluable for diagnosis.

Following the diagnosis of orbital cyst associated with anophthalmos and microphthalmos a review by a paediatrician is mandatory to search for any other associated systemic abnormalities. The classification proposed by Warburg²³ is useful for identifying syndromes associated with microphthalmos and cyst.

Genetic disorders responsible for microphthalmos and cyst include autosomal dominant, autosomal recessive, and X linked syndromes. Once the condition has been categorised, the family should be given genetic counselling.

Consideration should next be given to the visual potential in cases of microphthalmos. Elder²⁴ found that microphthalmic eyes with a corneal diameter of 5 mm or less at birth were associated with a visual acuity of no perception of light in 81% cases. If doubt persists as to the level of vision then electrodiagnostic tests should be considered. Eyes with some residual vision can be treated with clear conformers which do not obscure the visual axis.

Once the visual potential of the microphthalmic eye has been assessed, further treatment depends upon the age of the patient at presentation and the volume of the orbital contents (microphthalmic eye plus orbital cyst or just orbital cyst in cases of anophthalmia).

Patients can be categorised into two groups by age at presentation. These are birth to 5 years and 5 years or older. Within these groups the orbital volume is assessed clinically using the Duke-Elder classification¹²; (A) a relatively normal eye with a small cyst; (B) an obvious cyst associated with a grossly deformed eye; (C) a large cyst which has pushed the globe backward so that it is not visible clinically.

It has been noted already that patients with poor orbital volume can achieve very good cosmetic results if treated with conformers as early as possible.²⁵⁻²⁶ Several clinical and experimental studies have confirmed that enucleation in childhood compromises orbital growth.²⁷⁻²⁹ The earlier the enucleation, the greater the reduction in orbital bone growth, especially if no orbital implant is used. This has resulted in the recommendation that elective enucleation should be delayed until orbital bone growth is complete.²⁹⁻³⁰

In cases of microphthalmos and anophthalmos with a noticeable cyst, parents are often keen to have the cyst removed to improve cosmesis. It is then necessary to discuss with the parents the timing of the cyst removal. The relevant considerations are that the orbit is not fully grown until puberty and that any small or medium sized cyst (with an associated ocular remnant if present) will help to stimulate orbital expansion, much more effectively than an artificial implant. However, an equally important consideration is that of cosmesis, which becomes more relevant as the child starts school. A reasonable compromise is to consider orbital cyst removal around the age of 5 years. Although not fully developed, the orbital growth for both males and females at this stage is about 90% of that of an adult.³¹ Thus, some orbital expansion may be sacrificed for cosmesis as the child begins schooling. In cases of very large orbital cysts prolapsing through the palpebral fissure, they may cause bony expansion¹³ as well as being cosmetically unacceptable to parents. For this reason they are often excised at an early age.

At Moorfields Eye Hospital, we follow a policy of aggressive socket expansion, with conformers, at first presentation of anophthalmos or severe microphthalmos without a sizeable cyst. Ideally, this should start as soon after birth as possible. At the first visit to the prosthetist, it may be difficult to fit a rigid conformer owing to the very shallow conjunctival fornices. In these cases the use of a HEMMA expander is very helpful. This hydrophilic material swells within the socket as it absorbs moisture and gently stretches the fornices, which enables a larger conformer to be inserted after 24-48 hours. With the availability of the HEMMA conformer we have not found it necessary to suture conformers into the socket.⁷

With new patients (particularly those with poor orbital volume) follow up is on a weekly basis for the first few months as, with aggressive treatment with increasingly larger conformers, there should be a steady progression through the range of conformers. As the socket size increases, so the standard conformers fit less readily. At this stage a mould of the socket can be taken using Orthoprint, under general anaesthetic and a custom made prosthesis fitted.

When the decision to remove the orbital cyst has been made, the method of removal must be considered. The easiest is to aspirate fluid from within the cyst cavity.¹¹⁻¹⁹ In the majority of reported cases, the fluid within the cyst returned and aspiration had to be repeated. There are a few cases of permanent collapse of the cyst following aspiration.¹¹⁻¹⁹

By far the most common method for removing orbital cysts is surgical excision. Most authors have recommend decompressing large cysts before removal as they otherwise need large incisions to aid extraction, although it may be possible in some cases to remove a large cyst intact.²¹

Once the cyst has been removed, if the patient is anophthalmic or there is no significant microphthalmic remnant, an orbital implant can be inserted to maintain the orbital volume and aid cosmesis and orbital growth. Both silicone¹⁸⁻³² and hydroxyapatite²¹ have been used as orbital implants.

The presence of an orbital cyst with anophthalmos or microphthalmos aids orbital expansion and thus most cases can be treated successfully with aggressive conformer therapy alone. In cases of anophthalmos or microphthalmos with cyst where the orbital volume is poor other orbital volume expansion techniques should be considered if conformer therapy is thought to be inadequate on its own.

Dermis fat grafts have been proposed,³³ as they have the advantage of being autogenous and have an ability to enlarge as the child grows. Fat atrophy with this implant in paediatric patients is rare. Tissue expanders have also been proposed for enlargement in children with poor orbital volume.^{15 18 34} These comprise a silicone balloon with a reservoir. They are inserted and usually left for 6 months. Periodically, additional saline is injected into the reservoir to expand the balloon.

Craniofacial surgery is required infrequently in patients with microphthalmos or anophthalmos and cyst, as the presence of the cyst prevents severe micro-orbitism.¹⁸ However, a few patients have such contracted sockets that aggressive conformer therapy and orbital volume expansion do not result in satisfactory cosmesis. In these patients good results have been reported using three dimensional orbital bone expansion.^{15 25}

It has been suggested^{7 15} that any secondary surgery to the lids (such as medial and lateral canthoplasty, expansion of the palpebral fissures) or mucous membrane grafting to the socket are best left until maximal expansion has been achieved by conformer therapy. Early treatment may lead to cicatricial tissue formation which may ultimately lead to a poorer cosmetic result. Exceptions to this rule might be those cases where a lower lid coloboma is present⁹ which prevents the fitting of a stable conformer, such as in case 21 in this study.

In this study we have outlined the current treatment regimen used at Moorfields Eye Hospital which is based primarily on early and aggressive use of conformers. Grossly enlarged cysts (with proptosis through the palpebral fissure) were excised soon after presentation and replaced with an orbital implant and ocular prosthesis. Large cysts (with no proptosis from the palpebral fissure) were left to encourage orbital growth. Between the ages of 3–5 years they were removed when orbital growth was near completion. Small orbital cysts (and any microphthalmic remnant) were left and the socket treated with aggressive expansion with moulded conformers. In all cases regular follow up with frequent changes in conformer size, especially in the first weeks of life, maximised orbital growth and the subsequent cosmetic result. In this study 33 out of 34 patients had a good cosmetic result which illustrates that the orbital cyst in microphthalmos or anophthalmos performs a useful role in socket expansion and that the majority of patients with this condition can expect a good cosmetic outcome.

Authors' affiliations

C J McLean, N K Ragge, R B Jones, J R O Collin, Moorfields Eye Hospital, City Road, London, UK

Correspondence to: Mr J R O Collin, Moorfields Eye Hospital, City Road, London EC1V 2PD, UK; richard.collin@moorfields.nhs.uk

Accepted for publication 13 August 2002

REFERENCES

- 1 DeWals P, Iechat MF. Surveillance of congenital anomalies, years 1980–1984. *EUROCAT Rep* 1987;**2**:57–9.
- 2 Stoll C, Alembik Y, Doff B, et al. Epidemiology of congenital eye malformations in 131 760 consecutive births. *Ophthalmic Paediatr Genet* 1992;**13**:179–86.
- 3 Myrianthopoulos ND. Malformations in children from one to seven years. New York: Alan R Liss, 1985:329.
- 4 Clementi M, Turolla L, Mammi I, et al. Clinical anophthalmia: an epidemiological study in northeast Italy based on 368 256 consecutive births. *Teratology* 1992;**46**:551–3.
- 5 Dolk H, Busby A, Armstrong BG, et al. Geographical variation in anophthalmia and microphthalmia in England, 1988–94. *BMJ* 1998;**317**:905–9.
- 6 Tucker S, Jones B, Collin JRO. Systemic anomalies in 77 patients with congenital anophthalmos or microphthalmos. *Eye* 1996;**10**:310–14.
- 7 Kohn R. *Textbook of ophthalmic plastic and reconstructive surgery*. Philadelphia: Lea & Febiger 1988:71.
- 8 Dollfus MA, Marx P, Langlois J, et al. Congenital cystic eyeball. *Am J Ophthalmol* 1968;**66**:504–9.
- 9 Helveston EM, Malone E, Lashmet MH. Congenital cystic eye. *Arch Ophthalmol* 1970;**84**:622–4.
- 10 Baghdassarian SA, Tabbara KF, Matta CS. Congenital cystic eye. *Am J Ophthalmol* 1973;**76**:269–75.
- 11 Waring GO, Roth AM, Rodrigues MM. Clinicopathologic correlation of microphthalmos with cyst. *Am J Ophthalmol* 1976;**82**:714–21.
- 12 Duke-Elder S. Normal and abnormal development. Congenital deformities. In: *System of ophthalmology*. St Louis: CV Mosby, 1963;3:483.
- 13 Meyer E, Zonis S, Gdal-On M. Microphthalmos with orbital cyst: a clinicopathological report. *J Pediatr Ophthalmol* 1977;**14**:38–41.
- 14 Mafee MF, Jampol LM, Langer BG, et al. Computed tomography of optic nerve colobomas, morning glory anomaly, and colobomatous cyst. *Radiol Clin N Am* 1987;**25**:693–9.
- 15 O'Keefe M, Webb M, Pashby RC, et al. Clinical anophthalmos. *Br J Ophthalmol* 1987;**71**:825–32.
- 16 Zeiter HJ. Congenital microphthalmos. *Am J Ophthalmol* 1963;**55**:910–22.
- 17 Foxman S, Cameron JD. The clinical implications of bilateral microphthalmos with cyst. *Am J Ophthalmol* 1984;**97**:632–8.
- 18 Mouriaux F, Audo I, Defoort-Dhellemmes S, et al. Prise en charge des microphthalmies et anophthalmies congénitales. *Fr Ophthalmol* 1997;**20**:583–91.
- 19 Leatherbarrow B, Kwartz J, Noble JL. Microphthalmos with cyst in monozygous twins. *J Pediatr Ophthalmol Strabismus* 1990;**27**:294–8.
- 20 Fledelius HC. Ultrasonic evaluation of microphthalmos and coloboma. *Acta Ophthalmol (Scand)* 1996;**74**(Suppl 219):23–6.
- 21 Harris GJ, Dolman PJ, Simons KB. Microphthalmos with cyst. *Ophthalmic Surg* 1992;**23**:432–3.
- 22 Wright DC, Yuh WTC, Thompson HS, et al. Bilateral microphthalmos with orbital cysts: MR findings. *J Comput Assist Tomogr* 1987;**11**:727–9.
- 23 Warburg M. Classification of microphthalmos and coloboma. *J Med Genet* 1993;**30**:664–9.
- 24 Elder MJ. Aetiology of severe visual impairment and blindness in microphthalmos. *Br J Ophthalmol* 1994;**78**:332–4.
- 25 Marchac D, Cophignon J, Achard E, et al. Orbital expansion for anophthalmia and micro-orbitism. *Plast Reconstr Surg* 1977;**59**:486–91.
- 26 Price E, Simon JW, Calhoun JH. Prosthetic treatment of severe microphthalmos in infancy. *J Pediatr Ophthalmol Strabismus* 1986;**23**:22–4.
- 27 Pfeiffer RL. The effect of enucleation on the orbit. *Trans Am Acad Ophthalmol* 1945;**49**:236–9.
- 28 Kennedy RE. The effect of early enucleation on the orbit in animals and humans. *Am J Ophthalmol* 1965;**60**:277–306.
- 29 Apt L, Isenberg S. Changes in orbital dimensions following enucleation. *Arch Ophthalmol* 1973;**90**:393–5.
- 30 Whitnall SE. *Anatomy of human orbit and accessory organs of vision*. London: Humphrey Milford, 1932:98–108.
- 31 Farkas LG, Posnick JC, Hreczko TM, et al. Growth patterns in the orbital region: a morphometric study. *Cleft Palate Craniofac J* 1992;**29**:315–8.
- 32 Morax S, Smida R, Hurbli T, et al. Kyste colobomateux de l'orbite au cours d'une microphthalmie congénitale. *J Fr Ophthalmol* 1995;**18**:475–80.
- 33 Handler LF, Heher KL, Katowitz JA. Congenital and acquired anophthalmia. *Curr Opin Ophthalmol* 1994;**5**:84–90.
- 34 Berry FD. A modified tissue expander for socket enlargement in clinical anophthalmos. *Ophthalm Plast Reconstr Surg* 1991;**7**:41–7.