PERSPECTIVE

Cannabinoids and glaucoma

I Tomida, R G Pertwee, A Azuara-Blanco

Br J Ophthalmol 2004;88:708–713. doi: 10.1136/bjo.2003.032250

Glaucoma is one of the leading causes of blindness in the world. In spite of the diverse therapeutic possibilities, new and better treatments for glaucoma are highly desirable. Cannabinoids effectively lower the intraocular pressure (IOP) and have neuroprotective actions. Thus, they could potentially be useful in the treatment of glaucoma. The purpose of this article is to provide the reader with an overview of the latest achievements in research into the potential use of cannabinoids for glaucoma.

without medical supervision. Recently, several states have legalised the medical use of cannabinoids. In the United Kingdom, cannabis is registered as a schedule I—class C drug.^{1–4} The medical use of cannabinoids is currently restricted to dronabinol (Marinol, synthetic Δ^9 -tetrahydrocannabinol) and nabilone; these drugs are administered orally as anti-emetics and appetite stimulants for patients with AIDS or on chemotherapy. Other potential clinical applications for cannabinoids, including glaucoma therapy, are currently under intense investigation (see below).

PHARMACOLOGY

The cannabis plant has more than 480 chemical constituents.⁵ Of these, there is a group of at least 66 compounds that contain only carbon, hydrogen, and oxygen and are known collectively as cannabinoids. The remaining constituents of cannabis include nitrogenous compounds such as amino acids, proteins and glycoproteins, sugars, hydrocarbons, alcohols, aldehydes, ketones, simple acids, fatty acids, esters and lactones, steroids, terpenes, non-cannabinoid phenols, flavonoids, pigments (carotene and xanthophylls), and vitamin K. Terpenes are thought to be responsible for the characteristic odour of cannabis plants (fig 1). The main psychoactive constituent of cannabis is the cannabinoid, Δ^9 -tetrahydrocannabinol (Δ^9 -THC), the structure of which was determined by Gaoni and Mechoulam in the 1960s (fig 2).⁵ Most research into the pharmacological properties of plant cannabinoids has focused on this cannabinoid and, indeed, with the exception of cannabidiol and cannabidiol (figs 2 and 3), other plant cannabinoids have been subjected to little or no pharmacological research.^{6,7}

It is now generally accepted that Δ^9 -THC produces many of its effects by acting through cannabinoid receptors of which there are at least two subtypes, CB₁ and CB₂.⁸ Both subtypes are coupled through G_{i/o} proteins, negatively to adenylate cyclase and positively to mitogen activated protein kinase. In addition, CB₁ receptors are positively or negatively coupled through G_{i/o} proteins to certain calcium and potassium channels. CB₁ receptors, which were cloned in 1990, are present in brain, spinal cord, and certain peripheral tissues that include lung, heart, urogenital and gastrointestinal tracts, and the eye.^{9–11} Many CB₁ receptors are present on central and peripheral neurons, one of their functions being the modulation of neurotransmitter release. CB₂ receptors, cloned in 1993, seem to be located especially in cells and tissues associated with the immune system, such as the tonsils, spleen, and different types of

Cannabis/marijuana is the most frequent illicit drug used today for recreational purposes. Yet it is not widely known that the cannabis plant (*Cannabis sativa*; Latin for “planted hemp”) (fig 1) is one of the oldest drugs used for medical purposes. Its therapeutic use was first recorded in a classical medicine book by the Chinese emperor Shen Nung in 2737 BC. The medical use of cannabis was also known in other ancient cultures throughout India, Assyria, Greece, Africa, South America, Egypt, and the Roman Empire.¹

Cannabis was introduced on a larger scale into Western medicine during the 19th century, primarily by British doctors. They accumulated experience in the use of “Indian hemp” while working in the colonies, recommending it as an appetite stimulant, analgesic, muscle relaxant, anticonvulsant, and hypnotic. In 1839, Dr William Brooke O’Shaughnessy, an Irish physician at the Medical College of Calcutta, published a detailed report “On the preparations of the Indian Hemp or Gunjah.” After performing animal studies, he determined that cannabis preparations were safe and effective in treating rabies, rheumatism, epilepsy, and tetanus. The Ohio Medical Society of Physicians reported in 1860 successful treatment of “stomach pain and gastric distress,” psychosis, chronic cough, gonorrhoea, and neuralgia with cannabis. The plant was difficult to store, its extracts were variable in potency, and the effects of oral ingestion were not constant. Other new drugs were becoming available in the early 1900s with more reliable effects, and cannabis began to be misused for recreational purposes. The American Marijuana Tax Act of 1937 intended to prevent non-medical use, but made cannabis difficult to obtain for medical purposes too, and it was subsequently removed from the US pharmacopoeia in 1942.

In the United States cannabis is now classified as a schedule I drug, regarded as having high potential for abuse, and to be unsafe to take

See end of article for authors’ affiliations

Correspondence to:
A Azuara-Blanco, PhD,
FRCS(Ed), The Eye Clinic,
Aberdeen Royal Infirmary,
Aberdeen AB25 2ZN, UK;
Aazblanco@aol.com

Accepted for publication
4 October 2003

Figure 1 Photograph of the *Cannabis sativa* plant (provided by GW Pharmaceuticals, Wiltshire, UK)

leucocytes.^{8,9} One role of these receptors is modulation of cytokine release.

Endogenous agonists for cannabinoid receptors have also been discovered; this system of “endocannabinoids” and receptors constituting what is now generally known as the endocannabinoid system.¹² The endocannabinoids that have been identified to date are all analogues of arachidonic acid. They include arachidonoyl ethanolamide (anandamide; AEA), 2-arachidonoyl glycerol (2-AG) which bind more or less equally well to CB₁ and CB₂ receptors, and 2-arachidonoyl glyceryl ether (noladin) which is CB₁ selective (fig 4). AEA behaves as a partial cannabinoid receptor agonist with less CB₂ than CB₁ efficacy. Within the nervous system, endocannabinoids are synthesised and released by neurons on demand, functioning as neurotransmitters or neuromodulators. There is also evidence that endocannabinoids serve as

retrograde synaptic messengers.¹² Following their release, the effects of at least some endocannabinoids are thought to be rapidly terminated by cellular uptake and intracellular enzymatic hydrolysis.¹³

As detailed elsewhere^{6,8} some established cannabinoid receptor agonists bind more or less equally well to CB₁ and CB₂ receptors, important examples being WIN55212-2, which has marginally greater CB₂ than CB₁ affinity, and Δ^9 -THC, HU-210, CP55940, and nabilone (fig 3). As to CB₁ selective agonists other than noladin, these include the anandamide analogues, methanandamide, O-1812, arachidonyl-2'-chloroethylamide (ACEA), and arachidonylcyclopropylamide (ACPA). Examples of CB₂ selective agonists are the THC analogues, L-759633, L-759656, JWH-133 and HU-308. Selective CB₁ and CB₂ receptor antagonists have also been developed, the most important of these being the CB₁ selective SR141716A, AM251, AM281 and LY320135, and the CB₂ selective SR144528 and AM630.⁸ There is evidence, however, that these compounds are not “silent” antagonists.¹⁴ Thus, as well as attenuating the effects of CB₁ or CB₂ receptor agonists, they can by themselves elicit responses in some cannabinoid receptor-containing tissues that are opposite in direction from those elicited by CB₁ or CB₂ receptor agonists. While some of these “inverse cannabimimetic effects” may be attributable to a direct antagonism of responses elicited at cannabinoid receptors by released endocannabinoids, there is evidence that this is not the only possible mechanism and that these compounds are in fact all inverse agonists that can reduce CB₁ or CB₂ receptor constitutive activity.

Figure 2 Chemical structure of typical classic cannabinoids Δ^9 -tetrahydrocannabinol, cannabinol, and HU-211.

Figure 3 Chemical structure of nabilone, cannabidiol, and cannabigerol.

Figure 4 Chemical structure of endogenous cannabinoids anandamide (AEA), 2-arachidonoyl glycerol (2-AG), and 2-arachidonoyl glyceryl ether (noladin).

AEA activates vanilloid (TRPV1) receptors in addition to CB₁ and CB₂ receptors and there is also growing evidence for the existence of non-CB₁, non-CB₂, non-vanilloid pharmacological targets for AEA and/or other cannabinoids.^{8,15} There is evidence too that classic cannabinoids such as Δ^9 -THC, HU-211, and CBD have antioxidant properties (see below).^{7,16–18} In addition, Δ^9 -THC can act through presynaptic CB₁ receptors in the CNS to inhibit glutamic acid release and that HU-211 can block glutamate (NMDA) receptors.^{8,18}

As well as showing therapeutic potential as neuroprotective agents, Δ^9 -THC and CBD have other potential clinical applications, as indeed does the CB₁ receptor antagonist/inverse agonist, SR141716A.^{7,18–20} Δ^9 -THC, for example, is already used in the clinic as an appetite stimulant and antiemetic.¹⁸ In addition, this cannabinoid will most likely prove to be useful as an anti cancer drug²¹ and for the management of pain,^{22,23} and of various kinds of motor dysfunction that include the muscle spasticity, spasm, or tremor associated with multiple sclerosis and spinal cord injury,²⁴ the tics and psychiatric signs and symptoms of Tourette's syndrome, and the dyskinesia that is produced by L-dopa in patients with Parkinson's disease.^{25–27} As now discussed, one other important potential clinical application for cannabinoids, is the management of glaucoma.

CANNABINOIDS AND GLAUCOMA

In 1971, Hepler and Frank reported a 25–30% IOP lowering effect of smoking marijuana in a small number of subjects.²⁸ The duration of action of marijuana after smoking was relatively short, about 3–4 hours, and there seemed to be a dose-response relation.^{28,29} Other ocular effects were observed such as conjunctival hyperaemia, reduced tear production, and change in pupil size.²⁹ Acute systemic side effects induced by marijuana smoking included reduction of systemic blood pressure and tachycardia.⁴ Psychotropic effects were very variable and included euphoria or

dysphoria, disruption of short term memory, cognitive impairments, sense of time distortion, reduced coordination, and sleepiness.⁴

An earlier report on the effect of smoked marijuana indicated the possibility of tolerance. Thus, the IOP reduction appeared to be inversely related to the duration of marijuana use.³⁰ In contrast, Dawson *et al*³¹ reported on their ophthalmological findings comparing non-users with long term users of marijuana (10 years or more). After applying the water loading test to both groups, the reduction of IOP associated with marijuana treatment was similar between users and non-users.

Since these early observations numerous studies have been conducted confirming that different cannabinoids, including cannabidiol, cannabigerol, endogenous cannabinoids, and some synthetic cannabinoids, can reduce the IOP when administered systemically and topically (see below). Obviously, smoking of marijuana is not advisable as a long term treatment. In addition to the acute side effects, long term marijuana smoking is associated with emphysema-like lung changes, and possible increase in the frequency of lung cancer.³² Oral administration has been evaluated. However, there is a poor and variable absorption with this route,³³ at least for the cannabinoid formulations that have been investigated so far.

Mechanism of IOP reduction

The mechanism of action of cannabinoids in the human eye is not fully understood. Until recently, the effect of cannabinoids on IOP was assumed to be mediated through the CNS. Studies involving unilateral topical application of cannabinoids^{34,35} showed a large difference between the treated and untreated eye, suggesting a localised action. The experiments of Liu *et al*³⁶ revealed evidence pointing in the same direction: bolus administration of Δ^9 -THC into the cerebral ventricles, as well as ventriculocisternal perfusion with Δ^9 -THC in rabbits, in contrast with intravenous administration, did not change the IOP. Thus, the main site of action of cannabinoids on IOP is not in the central nervous system.

Pharmacological and histological studies support the direct role of ocular CB₁ receptors in the IOP reduction induced by cannabinoids. Straiker *et al*¹¹ detected CB₁ receptors in ocular tissues of the human eye, including the ciliary epithelium, the trabecular meshwork, Schlemm's canal, ciliary muscle, ciliary body vessels, and retina. Porcella *et al*¹⁰ found high levels of CB₁ mRNA in the ciliary body. The anatomical distribution of cannabinoid receptors suggests a possible influence of endogenous cannabinoids on trabecular and uveoscleral aqueous humour outflow and on aqueous humour production. In addition to the proved IOP lowering effect of CB₁ receptor agonists, Pate *et al*³⁷ could antagonise the IOP lowering effect of CP-55,940 (a synthetic CB₁ agonist) by pretreating the animals with SR 141716A (a CB₁ receptor antagonist). Similarly, Song *et al*³⁸ found that the IOP lowering effect of topical WIN-55,212-2 was significantly reduced by topically administered SR141716A.

Using the synthetic cannabinoid WIN-55,212-2, Chien *et al*³⁹ could demonstrate an 18% reduction in the aqueous humour production in monkeys but without significant change in the trabecular outflow facility. As this percentage appeared not sufficient to account for the total IOP lowering effect, other additional mechanisms were thought to be involved.

The IOP reducing effect does not seem to be related to a systemic reduction of arterial blood pressure.⁴⁰ However, a direct effect on the ciliary processes, and specifically a reduction in capillary pressure, leading to changes in aqueous humour dynamics, has been proposed.⁴¹ Green *et al*⁴² showed

that Δ^9 -THC decreased the secretion of ciliary processes and led to a dilatation of the ocular blood vessels through a possible β adrenergic action. In addition, Sugrue⁴³ indicated that cannabinoids may inhibit calcium influx through presynaptic channels and in this way reduce the noradrenaline release in the ciliary body, leading to a decrease in the production of aqueous humour. Porcella *et al*¹⁰ proposed that cannabinoids might be acting as vasodilators on blood vessels of the anterior uvea, thus improving the aqueous humour uveoscleral outflow.

Green *et al*^{44, 45} postulated that some cannabinoids may influence the IOP through a prostaglandin mediated mechanism. For example, topically applied AEA is hydrolysed to arachidonic acid, which is a COX pathway precursor of prostaglandins.^{46, 47}

The topical application of the CB₂ receptor agonist JWH-133 used in in vivo experiments by Laine *et al*⁴⁸ did not have any effect on IOP compared to vehicle treatments, indicating that CB₂ receptor agonists may not be involved in the regulation of IOP.

Topical application of cannabinoids

To minimise possible systemic adverse side effects and maximise the dose at the site of action, topical application would be the ideal form of administration. However, natural cannabinoid extracts as well as synthetic forms are highly lipophilic and have low aqueous solubility, creating practical difficulties for this mode of administration.

After instillation of an eye drop of any medication, loss of the instilled solution via the lacrimal drainage system and poor drug penetration results in only <5% of an applied dose reaching the intraocular tissues. The cornea is usually the major pathway for intraocular penetration of topical medications. The corneal epithelium is highly lipophilic and its penetration is a rate limiting step for lipophilic drugs. Aqueous solubility is another drug property important for efficacy of delivery, as the surface of the eye is constantly moistened by tear fluid. Additional factors affecting corneal absorption include the molecular size, charge, and degree of ionisation.⁴⁹

Previous experiments with topical cannabinoid solutions involved the use of light mineral oil as a vehicle, but proved to be irritant to the human eye.^{50, 51} Recently, different microemulsions and cyclodextrins (macrocylic oligosaccharides) have been shown to improve the corneal penetration of cannabinoids. These formulations successfully induced an unilateral IOP lowering effect.^{52–58} Cyclodextrins have already been used efficiently by Porcella *et al*⁵⁶ to administer the synthetic cannabinoid WIN-55,212-2 topically to glaucoma patients.

Neuroprotective and vascular actions of cannabinoids

In glaucoma, the final pathway leading to visual loss is the selective death of retinal ganglion cells through apoptosis. Apoptosis is initiated by axonal injury at the optic disc, either by compression and/or by ischaemia. In ischaemia, glutamate is released and activates NMDA receptors. NMDA receptor activation appears to be one of several pathways that result in apoptotic cell death. After activation of NMDA receptors there is an influx of calcium into the cells and free radicals are generated. Substances that prevent this cascade of events and inhibit the retinal ganglion cell death are currently under investigation.⁵⁹

Recent studies have documented the neuroprotective properties of cannabinoids. There is evidence that Δ^9 -THC can inhibit glutamic acid release by increasing K⁺ and decreasing Ca²⁺ permeability and that the synthetic cannabinoid HU-211 can block glutamate (NMDA) receptors.^{8, 9, 18, 59–62} These actions are mediated by presynaptic CB₁ receptors. Yoles *et al*, using a calibrated crush injury to adult

rat optic nerve (optic nerve axotomy), showed a beneficial effect of HU-211 on injury induced metabolic and electrophysiological deficits.⁶³ However, the optic nerve crush model may not resemble the mechanisms responsible for glaucomatous nerve damage.

Classic cannabinoids such as Δ^9 -THC, HU-211, and CBD have antioxidant properties that are not mediated by the CB₁ receptor. As a result, they can prevent neuronal death by scavenging toxic reactive oxygen species produced by overstimulation of receptors for the excitatory neurotransmitter, glutamic acid.^{7, 16–18, 64}

Cannabinoids have vasorelaxant properties and so might be able to increase the ocular blood flow. The mediator endothelin-1, produced by vascular endothelial cells, has a significant role in the regulation of local circulation, producing vasoconstriction and being involved in the pathophysiological processes of ischaemic and haemorrhagic stroke, Raynaud's phenomenon, ischaemic heart disease and pulmonary arterial hypertension, among others.⁶⁵ The possible role of endothelin-1 in the pathogenesis of glaucoma has been suggested. For example, patients with open angle glaucoma may have an abnormal increase in plasma endothelin-1 in response to vasospastic stimuli.^{66, 67} Mechoulam *et al*⁶¹ could demonstrate that 2-arachidonoylglycerol, an endogenous cannabinoid, was able to reduce endothelin induced Ca²⁺ mobilisation, inhibiting vasoconstriction. Thus, cannabinoids may have beneficial properties in ischaemia induced optic nerve damage.

FUTURE DIRECTIONS

Cannabinoids have the potential of becoming a useful treatment for glaucoma, as they seem to have neuroprotective properties and effectively reduce intraocular pressure. However, several challenges need to be overcome, including the problems associated with unwanted systemic side effects (psychotropic, reduction in systemic blood pressure), possible tolerance, and the difficulty in formulating a stable and effective topical preparation. Some cannabinoids such as HU-211 and cannabidiol do not have psychotropic effects, while maintaining their IOP lowering action, so that further research on these compounds would be desirable. Tolerance may develop after repeated use of cannabinoids.³⁰ However, tolerance might be beneficial if it develops only or preferentially to unwanted side effects. There has been recent progress in the use of microemulsions and cyclodextrins to overcome the barriers in ocular penetration of topically applied cannabinoids.

Other possible applications of cannabinoids in ophthalmology could be explored. Age related macular degeneration (AMD) is the leading cause of blindness in the United Kingdom. Perhaps the potent antioxidant properties of the cannabinoids may be beneficial in AMD, offering a possible alternative to established antioxidant supplements.⁶⁸ Cannabinoids have been shown to inhibit angiogenesis, leading to a decrease in the expression of proangiogenic factors such as VEGF.⁶⁹ Evidence suggests that VEGF plays a major part in the development of choroidal neovascularisation in AMD, and clinical trials using anti-VEGF therapies are being conducted.⁷⁰ The CB₂ receptors are also under intense investigation for their possible immunomodulatory effects.⁷¹ The anti-inflammatory properties of CB₂ receptor agonists might also prove to be of therapeutic relevance in different forms of inflammatory eye disease.

Authors' affiliations

I Tomida, A Azuara-Blanco, Department of Ophthalmology, Aberdeen Royal Infirmary, University of Aberdeen, UK

R G Pertwee, Department of Biomedical Sciences, Institute of Medical Sciences, University of Aberdeen, UK

REFERENCES

- 1 Frankhauser M. Cannabis in der westlichen Welt. In: Grotenhermen F, ed. *Cannabis und cannabinoide*. 1st ed. Berlin: Verl Hans Huber, 2001:57–71.
- 2 Chalsma AL, Boyum D. *Marijuana situation assesment*. Washington DC: Office of National Drug Control Policy, 1994.
- 3 Schwab IR, Husted R, Liegner JT, et al. *Complementary therapy assessment: Marijuana in the treatment of Glaucoma*. San Francisco: American Academy of Ophthalmology, Complementary Therapy Task Force, 1999.
- 4 Grotenhermen F. Die Wirkungen von Cannabis und der Cannabinoide. In: Grotenhermen F, ed. *Cannabis und cannabinoide*, 1st ed. Berlin: Verl Hans Huber, 2001:75–86.
- 5 ElSohly MA. Chemical constituents of cannabis. In: Grotenhermen F, Russo E, eds. *Cannabis and cannabinoids. Pharmacology, toxicology and therapeutic potential*. New York: Haworth Press, 2002:27–36.
- 6 Pertwee RG. Pharmacology of cannabinoid receptor ligands. *Curr Med Chem* 1999;6:635–64.
- 7 Pertwee RG. The pharmacology and therapeutic potential of cannabidiol. In: Di Marzo V, ed. *Cannabinoids*. New York: Landes Bioscience, 2003 (in press).
- 8 Howlett AC, Barth F, Bonner TI, et al. International Union of Pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacol Rev* 2002;54:161–202.
- 9 Pertwee RG. Pharmacology of cannabinoid CB1 and CB2 receptors. *Pharmacol Ther* 1997;74:129–80.
- 10 Porcella A, Casellas P, Gessa GL, et al. Cannabinoid receptor CB1 mRNA is highly expressed in the rat ciliary body: implications for the antiglaucoma properties of marihuana. *Mol Brain Res* 1998;58:240–5.
- 11 Straiker A, Maguire C, Makie K, et al. Localization of cannabinoid CB1 receptors in the human anterior eye and retina. *Invest Ophthalmol Visual Sci* 1999;40:2442–8.
- 12 Pertwee RG, Ross RA. Cannabinoid receptors and their ligands. *Prostaglandins, Leukotrienes and Essential Fatty Acids* 2002;66:101–21.
- 13 Piomelli D, Giuffrida A, Calignano A, et al. The endocannabinoid system as a target for therapeutic drugs. *Trends Pharmacol Sci* 2000;21:218–24.
- 14 Pertwee RG. Inverse agonism at cannabinoid receptors. In: *Esteve Foundation Symposium X. Inverse Agonism*. Amsterdam: Elsevier, 2003 (in press).
- 15 Pertwee RG. Novel pharmacological targets for cannabinoids. *Current Neuropharmacology* 2003 (in press).
- 16 Hampson AJ, Grimaldi M, Axelrod J, et al. Cannabidiol and (-)-delta-9-tetrahydrocannabinol are neuroprotective antioxidants. *Proc Natl Acad Sci USA* 1998;95:8268–73.
- 17 Hampson AJ, Grimaldi M, Lolic M, et al. Neuroprotective antioxidants from marijuana. In: *Reactive oxygen species: from radiation to molecular biology*. Ann NY Acad Sci 2000;899:274–82.
- 18 Pertwee RG. Neuropharmacology and therapeutic potential of cannabinoids. *Addiction Biology* 2000;5:37–46.
- 19 Barth F. Cannabinoid receptor agonists and antagonists. *Expert Opinion On Therapeutic Patents* 1998;8:301–13.
- 20 Pertwee RG. Cannabinoid receptor ligands: clinical and neuropharmacological considerations relevant to future drug discovery and development. *Expert Opinion on Investigational Drugs* 2000;9:1553–71.
- 21 Galve-Roperh I, Sánchez C, Cortés ML, et al. Anti-tumoral action of cannabinoids: involvement of sustained ceramide accumulation and extracellular signal-regulated kinase activation. *Nature Med* 2000;6:313–19.
- 22 Pertwee RG. Cannabinoid receptors and pain. *Progress in Neurobiology* 2001;63:569–611.
- 23 Pertwee RG. Cannabinoids. In: Bountra C, Munglani R, Schmidt WK, eds. *Pain current understanding, emerging therapies and novel approaches to drug discovery*. New York: Marcel Dekker, 2003:683–706.
- 24 Pertwee RG. Cannabinoids and multiple sclerosis. *Pharmacol Ther* 2002;95:165–74.
- 25 Hemming M, Yellowlees PM. Effective treatment of Tourette's syndrome with marijuana. *J Psychopharmacol* 1993;7:389–91.
- 26 Müller-Vahl KR, Kolbe H, Schneider U, et al. Cannabis in movement disorders. *Forschende Komplementärmedizin* 1999;6(Suppl 3):23–7.
- 27 Müller-Vahl KR, Schneider U, Kolbe H, et al. Treatment of Tourette's syndrome with delta-9-tetrahydrocannabinol. *Am J Psychiatr* 1999;156:495.
- 28 Hepler RS, Frank IR. Marijuana smoking and intraocular pressure (letter). *JAMA* 1971;217:1392.
- 29 Brown B, Adams AJ, Haegerstrom-Portnoy G, et al. Pupil size after use of marijuana and alcohol. *Am J Ophthalmol* 1977;83:350–4.
- 30 Flom MC, Adams AJ, Jones RT. Marijuana smoking and reduced pressure in human eyes: drug action or epiphenomenon? *Invest Ophthalmol* 1975;14:52–5.
- 31 Dawson WW, Jiménez-Antillon CF, Perez M, et al. Marijuana and vision—after ten years' use in Costa Rica. *Invest Ophthalmol Visual Sci* 1977;16:689–99.
- 32 Hashibe M, Ford DE, Zhang ZF. Marijuana smoking and head and neck cancer. *J Clin Pharmacol* 2002;42(Suppl):103–7.
- 33 Merritt JC, McKinnon S, Armstrong JR, et al. Oral delta 9-tetrahydrocannabinol in heterogeneous glaucomas. *Ann Ophthalmol* 1980;12:947–50.
- 34 Colasanti BK, Craig CR, Allara RD. Intraocular pressure, ocular toxicity and neurotoxicity after administration of cannabidiol or cannabigerol. *Exp Eye Res* 1984;39:251–9.
- 35 Colasanti BK, Powell SR, Craig CR. Intraocular pressure, ocular toxicity and neurotoxicity after administration of delta-9-tetrahydrocannabinol or cannabichromene. *Exp Eye Res* 1984;38:63–71.
- 36 Liu JHK, Dacus AC. Central nervous system and peripheral mechanisms in ocular hypotensive effect of cannabinoids. *Arch Ophthalmol* 1987;105:245–8.
- 37 Pate DW, Järvinen K, Urtti A, et al. Effect of the CB1 receptor antagonist SR 141716A on cannabinoids induced ocular hypotension in normotensive rabbits. *Life Sci* 1998;63:2181–8.
- 38 Song ZH, Slowey CA. Involvement of cannabinoids receptors in the intraocular pressure lowering effects of WIN-55,212-2. *Pharmacol Exp Ther* 2000;292:136–9.
- 39 Chien FY, Wang RF, Mittag TW, et al. Effect of WIN-55,212-2, a cannabinoid receptor agonist, on aqueous humor dynamics in monkeys. *Arch Ophthalmol* 2003;121:87–90.
- 40 Korzyn AD. The ocular effects of cannabinoids. *Gen Pharmac* 1980;101:591–3.
- 41 McDonald TF, Cheeks L, Slagle T, et al. Marijuana-derived material induced changes in monkey ciliary processes differ from those in rabbit ciliary processes. *Curr Eye Res* 1991;4:305–12.
- 42 Green K, Pederson JE. Effect of delta-9-tetrahydrocannabinol on aqueous dynamics and ciliary body permeability in the rabbit. *Exp Eye Res* 1973;15:499–507.
- 43 Sugrue MF. New approaches to antiglaucoma therapy. *J Med Chem* 1997;40:2793–809.
- 44 Green K, Pados SM. Antagonism of arachidonic acid-induced ocular effects by delta-9-tetrahydrocannabinol. *Invest Ophthalmol Vis Sci* 1974;13:422–9.
- 45 Green K, Krease EC, McIntyre OL. Interaction between delta-9-tetrahydrocannabinol and indometacin. *Ophthalmic Res* 2001;33:217–20.
- 46 Pate DW, Järvinen K, Urtti A, et al. Effects of topical anandamides on intraocular pressure in normotensive rabbits. *Life Sci* 1996;58:1849–60.
- 47 Pate DW, Järvinen K, Urtti A, et al. Effects of topical alpha-substituted anandamides on intraocular pressure in normotensive rabbits. *Pharm Res* 1997;14:1738–43.
- 48 Laine K, Järvinen K, Järvinen T. Topically administered CB2-receptor agonist, JWH-133, does not decrease intraocular pressure (IOP) in normotensive rabbits. *Life Sci* 2003;72:837–42.
- 49 Järvinen T, Pate DW, Laine K. Cannabinoids in the treatment of glaucoma. *Pharmacology and Therapeutics* 2002;95:203–220.
- 50 Green K, Roth M. Ocular effects of topical administration of delta-9-tetrahydrocannabinol in man. *Arch Ophthalmol* 1982;100:263–265.
- 51 Jay WM, Green K. Multiple drop study of topically applied 1% delta-9-tetrahydrocannabinol in human eyes. *Arch Ophthalmol* 1983;101:591–3.
- 52 Järho P, Urtti A, Pate DW, et al. Increase in aqueous solubility, stability and in vivo corneal permeability of anandamide by hydroxypropyl-beta-cyclodextrin. *Int J Pharm* 1996;137:209–17.
- 53 Järho P, Pate DW, Brenneisen R, et al. Hydroxypropyl-beta-cyclodextrin and its combination with hydroxypropyl-methylcellulose increases aqueous solubility of delta-9-tetrahydrocannabinol. *Life Sci* 1998;63:381–4.
- 54 Kearse EC, Green K. Effect of vehicle upon in vivo transcorneal permeability and intracorneal content of delta-9-tetrahydrocannabinol. *Curr Eye Res* 2000;29:496–501.
- 55 Green K, Kearse EC. Ocular penetration of topical delta-9-tetrahydrocannabinol from rabbit corneal or cul-de-sac application site. *Curr Eye Res* 2000;21:566–70.
- 56 Porcella A, Maxia Ch, Gessa GL, et al. The synthetic cannabinoids WIN-55,212-2 decreases the intraocular pressure in human glaucoma resistant to conventional therapies. *Eur J Neurosci* 2001;13:409–12.
- 57 Laine K, Järvinen K, Mechoulam R, et al. Comparison of the enzymatic stability and intraocular pressure effects of 2-Arachidonylglycerol and noladin ether, a novel putative endocannabinoid. *Invest Ophthalmol Vis Sci* 2002;43:3216–22.
- 58 Naveh N, Weissman C, Muchtar S, et al. A submicron emulsion of HU-211, a synthetic cannabinoid, reduces intraocular pressure in rabbits. *Graefes Arch Clin Exp Ophthalmol* 2000;238:334–8.
- 59 Levin LA. Direct and indirect approaches to neuroprotective therapy of glaucomatous optic neuropathy. *Surv Ophthalmol* 1999;43(Suppl):98–101.
- 60 Shen M, Thayer SA. Cannabinoid receptor agonists protect cultured rat hippocampal neurons from excitotoxicity. *Mol Pharmacol* 1998;54:459–62.
- 61 Mechoulam R, Panikashvili D, Shohami E. Cannabinoids and brain injury: therapeutic implications. *Trends Mol Med* 2002;8:58–61.
- 62 Jin KL, Mao XO, Goldsmith PC, et al. CB1 cannabinoid receptor induction in experimental stroke. *Ann Neurol* 2000;48:257–261.
- 63 Yoles E, Belkin M, Schwartz M. HU-211, a nonpsychotropic cannabinoid, produces short- and long-term neuroprotection after optic nerve axotomy. *J Neurotrauma* 1996;13:49–57.
- 64 Marsicano G, Moosman B, Hermann H, et al. Neuroprotective properties of cannabinoids against oxidative stress: role of cannabinoids receptor CB1. *J Neurochem* 2002;80:448–56.
- 65 Gray GA, Battistini B, Webb DJ. Endothelins are potent vasoconstrictors, and much more besides. *Trends Pharmacol Sci* 2000;21:38–40.
- 66 Haefliger IO, Meyer P, Flammer J, et al. The vascular endothelium as a regulator of the ocular circulation: a new concept in ophthalmology? *Surv Ophthalmol* 1994;39:123–32.
- 67 Nicoletta MT, Ferrier SN, Morrison CA, et al. Effects of cold-induced vasospasm in glaucoma. The role of endothelin-1. *Invest Ophthalmol Vis Sci* 2003;44:2565–72.
- 68 Age-Related Eye Disease Study Research Group. A randomised, placebo-controlled, clinical trial of high-dose supplementation with vitamins C and E,

- beta carotene, and zinc for age-related macular degeneration and vision loss. *Arch Ophthalmol* 2001;**119**:1417–36.
- 69 **Casanova ML**, Blázquez C, Martínez-Palacio J, *et al*. Inhibition of skin tumor growth and angiogenesis in vivo by activation of cannabinoids receptors. *J Clin Invest* 2003;**111**:43–50.
- 70 **The Eye Tech Study Group**. Preclinical and phase 1A clinical evaluation of an anti-VEGF pegylated aptamer (EYE001) for the treatment of exudative age-related macular degeneration. *Retina* 2002;**22**:143–52.
- 71 **Zurier RB**. Prospects for cannabinoids as anti-inflammatory agents. *J Cell Biochem* 2003;**88**:462–6.