Retraction notice

Ramosetron compared with granisetron for the prevention of vomiting following strabismus surgery in children

Yoshitaka Fujii,¹ Hiroyoshi Tanakaa,¹ Mutsuko Itob²

¹Department of Anaesthesiology, Toride Kyodo General Hospital, Toride City, Ibaraki, Japan,

²Department of Ophthalmology

Correspondence to: Yoshitaka Fujii, Department of Anaesthesiology, University of Tsukuba Institute of Clinical Medicine, 2-1-1, Amakubo, Tsukuba City, Ibaraki 305-8576, Japan yfujii@igaku.md.tsukuba.ac.jp

Notice of formal retraction of article by Dr Y. Fujii. This article is being retracted as a result of:

- (1) Overwhelming evidence of fabrication, related to the fact that the distributions of many variables reported by Dr Fujii in these studies could not have occurred by chance;^{1 2} and
- (2) The inability of Dr Fujii's institutions to attest to the integrity of the study and/or its data conducted under their auspices, as set out in the Joint Editors-in-Chief Request for Determination of April 9, 2012.³

Br J Ophthalmol 2016;**100**:150. doi:10.1136/bjo.85.6.670ret

CrossMark

REFERENCES

- 1 Carlisle JB. The analysis of 168 randomised controlled trials to test data integrity. *Anaesthesia* 2012;67:521–37.
- 2 Kranke P, Apfel CC, Roewer N. Reported data on granisetron and postoperative nausea and vomiting by Fujii *et al.* are incredibly nice! Fujii Y, Response. *Anesth Analg* 2000;90:1004–7
- 3 http://www.oxfordjournals.org/our_journals/bjaint/fujii_joint_editorial_request_regarding_dr_yoshitaka_fujii.pdf