

Supplemental Table 2. List of Topical and Systemic Medications Used in the Present Study

Medications	NTG Eyes (n = 43)			Healthy Eyes (n = 24)		
	No. (%)	OR per $\mu\text{L}/\text{min}$ Average TA RBF (p value)	OR per $\mu\text{L}/\text{min}$ Average TV RBF (p value)	No. (%)	OR per $\mu\text{L}/\text{min}$ Average TA RBF (p value)	OR per $\mu\text{L}/\text{min}$ Average TV RBF (p value)
Prostaglandin analogs	36 (84)	0.80 (0.11)	0.88 (0.19)	0	–	–
Topical beta blockers	22 (51)	0.93 (0.48)	0.97 (0.75)	0	–	–
Topical carbonic anhydrase inhibitors	18 (42)	0.94 (0.54)	0.97 (0.67)	0	–	–
α 2-adrenergic agonists	6 (14)	0.86 (0.29)	0.83 (0.12)	0	–	–
Rho-associated protein kinase inhibitor	3 (7)	0.99 (0.94)	0.93 (0.68)	0	–	–
Calcium channel blockers	5 (12)	0.96 (0.79)	0.98 (0.87)	3 (13)	0.83 (0.51)	1.08 (0.67)
Diuretics	1 (2)	1.26 (0.50)	1.32 (0.30)	0	–	–
Angiotensin II receptor blockers	3 (7)	0.73 (0.16)	0.88 (0.45)	3 (13)	0.54 (0.13)	0.56 (0.14)
Anti-platelets	4 (9)	0.82 (0.27)	0.94 (0.69)	3 (13)	0.82 (0.49)	0.71 (0.26)
Anti-lipids	9 (21)	0.90 (0.44)	1.00 (0.98)	4 (17)	1.19 (0.41)	1.00 (0.98)
Pacemaker	2 (5)	1.10 (0.68)	0.91 (0.72)	0	–	–
Asthma medications	3 (7)	0.65 (0.07)	0.82 (0.27)	1 (4)	0.92 (0.84)	1.30 (0.36)

Abbreviations: NTG, normal tension glaucoma; OR, odds ratio; TA, temporal artery; RBF, retinal blood flow; TV, temporal vein

Calculated as logistic regression analysis, with $p < 0.05$ considered to be statistically significant