



OPEN ACCESS

# Retinal nerve fibre layer thickness measured with SD-OCT in a population-based study: the Handan Eye Study

Jian Wu,<sup>1,2</sup> Yifan Du ,<sup>1</sup> Caixia Lin,<sup>1</sup> Yingting Zhu,<sup>3</sup> Wei Chen,<sup>2</sup> Qing Pan,<sup>4</sup> Yehong Zhuo ,<sup>3</sup> Ningli Wang <sup>1</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bjophthalmol-2021-320618>).

<sup>1</sup>Beijing Institute of Ophthalmology, Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University; Beijing Ophthalmology & Visual Sciences Key Laboratory, Beijing, China

<sup>2</sup>Department of Ophthalmology, Stanford University School of Medicine, Palo Alto, California, USA

<sup>3</sup>State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, China

<sup>4</sup>Department of Epidemiology and Biostatistics, West China School of Public Health and West China Fourth Hospital, Sichuan University, Chengdu, Sichuan, China

## Correspondence to

Professor Ningli Wang, Ophthalmology, Beijing Tongren Hospital, Beijing, Beijing, China; [wningli@vip.163.com](mailto:wningli@vip.163.com)

JW and YD contributed equally.

JW and YD are joint first authors.

Received 18 October 2021  
Accepted 13 March 2022  
Published Online First  
5 April 2022



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

**To cite:** Wu J, Du Y, Lin C, et al. *Br J Ophthalmol* 2023;**107**:1156–1164.

## ABSTRACT

**Purpose** To examine the normative profile of retinal nerve fibre layer (RNFL) thickness and ocular parameters based on spectral-domain optical coherence tomography (SD-OCT) and its associations with related parameters among the Chinese population.

**Methods** This population-based cohort Handan Eye Study (HES) recruited participants aged  $\geq 30$  years. All subjects underwent a standardised ophthalmic examination. Peripapillary RNFL thickness was obtained using SD-OCT. Mixed linear models were adopted to evaluate the correlation of RNFL thickness with ocular parameters as well as systemic factors. R.V.3.6.1 software was used for statistical analysis.

**Results** 3509 subjects (7024 eyes) with the average age of  $55.54 \pm 10.37$  were collected in this analysis. Overall mean RNFL thickness measured was  $113.46 \pm 10.90$   $\mu\text{m}$ , and the thickest quadrant of parapapillary RNFL was the inferior quadrant, followed by the superior quadrant, the nasal quadrant and the temporal quadrant. In the multivariate linear regression model, thinner RNFL thickness was remarkable association with male ( $p < 0.001$ ), older age ( $p < 0.001$ ), increased body mass index ( $> 30$ ,  $p = 0.018$ ), absence of diabetes ( $p = 0.009$ ), history of cataract surgery ( $p = 0.001$ ), higher intraocular pressure ( $p = 0.007$ ), lower spherical equivalent ( $p < 0.001$ ) and increased axial length ( $p = 0.048$ ).

**Conclusions** In non-glaucoma individuals, this difference of RNFL thickness in Chinese population should be noted in making disease diagnoses. Meanwhile, multiple ocular and systemic factors are closely related to the thickness of RNFL. Our findings further emphasise the need to demonstrate ethnic differences in RNFL thickness and the specificity of associated ocular and systemic factors, as well as to develop better normative databases worldwide.

**Trial registration number** HES was registered in Chinese Clinical Trial Registry website, and the registry number was ChiCTR-EOC-17013214.

## INTRODUCTION

Glaucoma is a leading cause of blindness, with a prevalence of 3.54% worldwide. Glaucoma is characterised by degeneration of retinal ganglion cells, which results in thinning of the retinal nerve fibre layers (RNFL) and irreversible visual field damaged.<sup>1</sup> Several studies indicated that approximately half of the retinal nerve fibres may loss before occurrence of visual field defects in

## Key messages

### What is already known on this topic

⇒ The thickness of retinal nerve fibre layer (RNFL) is associated with varieties of eye diseases, which had been investigated in previous studies. However, lack of population-based studies in Asian population reported the RNFL thickness normative profile.

### What this study adds

⇒ This study explored the distribution of the RNFL thickness in the representative Chinese population.  
⇒ Meanwhile, the results conducted that RNFL thickness in our study was thicker than European population considering the ethnic difference.  
⇒ What's more, the study demonstrated that the ocular and systemic factors were closely affecting RNFL thickness, like examination instruments, age, gender and axial length.

### How this study might affect research, practice or policy

⇒ Using the normative European database of RNFL thickness to measure the Chinese population may lead to misdiagnosis, especially glaucoma and neurodegenerative disease.  
⇒ The influence of various factors on RNFL thickness should be well considered during the eye disease diagnosis procedure.  
⇒ For achieving the goal of precision medicine, the results emphasise the critical need to establish a better normative database that included a multiethnic population worldwide.

glaucoma patients.<sup>2,3</sup> Therefore, the assessment of RNFL thickness is important in the early detection and diagnosis of glaucoma.

Spectral-domain optical coherence tomography (SD-OCT) is a noninvasive imaging technique that used to evaluate the RNFL and optic disc with high resolution cross-sectional<sup>4</sup> retinal images.<sup>5</sup> As SD-OCT technology grows gradually and the sufficient resolution images conduct the automated segmentation and measurement of individual retinal layer clearly, the evaluation of RNFL thickness has been extensively applied and involved in clinical practice for glaucoma diagnosis and

follow-up. Meanwhile, as the retina is a laminated constituent of the central nervous system, it is the only part of the central nervous system which can be imaged optically with non-invasive methods, RNFL thickness measurements is also considered as a biomarker for neurodegenerative diseases, such as Parkinson's<sup>6</sup> and Alzheimer's<sup>7</sup> disease.

The normative database of RNFL thickness on SD-OCT was built in European, considering the parameters are different between each ethnic population,<sup>8,9</sup> several studies around the world have been dedicated to eye diseases. Most of the research on OCT-based examinations are usually derived from relatively small sample populations,<sup>10,11</sup> or assessed in hospital-based studies, and few of them have described RNFL thickness distribution in the healthy population-based setting, giving rise to potential selection bias.<sup>12</sup>

Since knowledge of RNFL thickness and optic nerve parameters are important for detection of glaucoma and other possible neurodegeneration diseases, the aim of this study was to investigate the distribution of peripapillary RNFL (pRNFL) thickness in the population-based Handan Eye Study (HES). In addition, by analysing the correlation of RNFL with age, gender and related ophthalmic parameters to further identify the potential risk factors of glaucomatous fundus damage.

## MATERIAL AND METHODS

### Study population

The HES is a population-based cohort study that recruited subjects with an age of 30+ years residing in North China. Our study population was made up of 6830 individuals from the baseline visit in 2006–2007, 5394 participated the 6-year follow-up visit in 2012–2013 (response rate 85.3%). Without undergoing SD-OCT assessments, the baseline visits were not included in the study. The details about the study have been illustrated previously.<sup>13</sup>

The OCT data from 4037 participants (7950 eyes) were available except the participants without OCT measurements. Of them, 425 eyes were further excluded due to poor OCT images quality with signal strength less than 45. Meanwhile, 177 eyes with glaucoma, retinal disease or history of ocular surgery were also excluded. Glaucoma was diagnosed by three aspects: the eye fundus, the visual field results, and intraocular pressure (IOP), and finally defined according to the standardised International Society of Geographical and Epidemiological Ophthalmology criteria.<sup>14</sup> Therefore, the research enrolment a total of 4766 eyes of 2638 participants.

### Ocular and systemic examination

All participants underwent a comprehensive clinical examination by ophthalmologists receiving standardised training. A detailed interviewer questionnaire was conducted to collect demographic variables, educational background, a medical history consisting of smoking and known major systemic diseases information. Body mass index (BMI) was calculated as the ratio of body weight (in kilograms) divided by body height (in metres) squared. Hypertension was defined as systolic blood pressure  $\geq 140$  mm Hg or diastolic blood pressure  $\geq 90$  mm Hg, or use of antihypertensive medications.<sup>15</sup> Diabetes was defined as fasting plasma glucose  $\geq 7.0$  mmol/L, self-reported diagnosis of diabetes or the use of antidiabetic medications.<sup>16</sup> Coronary heart disease was defined as self-reported coronary heart disease, stroke or related peripheral artery disease. Blood samples were collected for measurements of high-density lipoprotein, low-density lipoprotein and triglyceride concentration.

Slit-lamp biomicroscopy was performed by experienced ophthalmologists and IOP was measured using the Kowa applanation tonometer (HA-2, Kowa Company, Tokyo, Japan). Best-corrected visual acuity (BCVA) was tested monocularly (right eye followed by left eye) then using a log MAR chart at a distance of 4 m binocularly. A 10 MHz A/B-mode ultrasound device was used (Cine Scan, Quantel Medical, Clermont-Ferrand, France) to measure axial length (AL). Autorefractor (KR8800, Topcon, Tokyo, Japan) was applied to measure the refraction and corneal curvature on the occasion of no pupil dilation.

### OCT Imaging

pRNFL parameters in both eyes were measured by SD-OCT (RTVue 100-2, Optovue, Fremont, California, USA; V.4.0). The device uses a scanning laser diode to emit infrared light-source with wavelength of 840 nm and acquires 26 000 A-scans per second scan. The position of the aiming circle was adjusted by the experienced examiner to match the optic nerve head (ONH), and the 12 radial scans (scans ranging from 1.3 to 4.9 mm) was made covering a measurement area of 3.4 mm diameter ring that included the ONH and surroundings in all directions. Various ONH and RNFL parameters (disc, cup and rim area, cup and ONH volume, cup/disc ratio, average and vertical cup-to-disc ratios, average and per-quadrant pRNFL thickness) were measured by algorithms native of RTVue OCT automatically. The outputs parameters included (1) average RNFL thickness; (2) temporal, superior, nasal, and inferior average RNFL thickness; and (3) 16 sections of the measuring circle around the ONH (each section was 22.58). OCT scans images quality is described by the signal strength index (SSI), which is based on the intensity of the reflected light. The SSI ranged from near 0 (no signal) to approximately 90 (very strong signal), algorithm segmentation failure and obvious decentration misalignment were excluded further from the analysis with SSI less than 45 (as recommended by the manufacturer).

### Statistical methods

Data analysis was carried out by using the statistical software R (V.3.6.1; R Core Team, 2019). The quantitative data were represented in mean and SD (or median, IQR) and qualitative data was described with counts and percentage.  $\chi^2$  test, t-test, and one-way analysis of variance (ANOVA), Turkey tests as post hoc tests after ANOVA were applied to detect the difference between groups. Bonferroni correction to control for the potential false discovery in multiple comparisons. Variance inflation factor (VIF) for each covariate in the multivariate model was conducted, a linear mixed model was used to estimate the relationship between individual features and RNFL. The significant level was set as  $p < 0.05$ .

## RESULTS

### Comparisons of characteristics between the excluded and included individuals

#### Study population

The demographic characteristics in the study population are described in table 1. In total, 7024 eyes of 3509 participants were included in the analysis (1912 men and 1607 women). The age ranged from 35 to 86 with an average of  $55.54 \pm 10.37$  years. Compared with excluded group, the included subjects tended to be significantly younger ( $p < 0.001$ ), higher body height and weight ( $p < 0.001$ ), higher systolic ( $p < 0.001$ ), absence of diabetes and lower glycosylated hemoglobin (HbA1c) ( $p < 0.001$ ), less

**Table 1** Comparisons of demographic and biochemical characteristics between the included and excluded individuals

Variables	Excluded group (N=1885)	Included group (N=3509)	T/ $\chi^2$	P value
Age (years)	59.56±11.79	55.54±10.37	12.446	<0.001
Sex (men, %)	789 (41.9)	1617 (46.1)	8.688	<b>0.003</b>
High educational level				
Lower than high school	1843 (97.8)	3394 (96.7)	4.777	<b>0.029</b>
High school or above	42 (2.2)	115 (3.3)		
Smoking habits				
Yes	409 (23.6)	916 (26.6)	5.250	<b>0.022</b>
No	1326 (76.4)	2533 (73.4)		
Body mass index (kg/m <sup>2</sup> )	25.78±4.06	25.92±3.88	-1.199	0.231
Body height	158.49±8.46	159.76±8.05	-5.247	<0.001
Body weight	64.93±11.51	66.19±11.22	-3.810	<0.001
Blood pressure diastolic (mm Hg)	83.66±13.51	84.09±12.74	-1.120	0.263
Blood pressure systolic (mm Hg)	145.60±23.63	141.63±21.52	6.041	<0.001
Hypertension				
Yes	593 (35.3)	1015 (30.5)	11.855	<b>0.001</b>
No	1085 (64.7)	2317 (69.5)		
Diabetes				
Yes	123 (7.4)	128 (3.9)	27.739	<0.001
No	1538 (92.6)	3167 (96.1)		
HbA1c (%)	5.87±1.05	5.71±0.75	5.834	<0.001
High-density lipoprotein cholesterol (mmol/L)	1.20±0.27	1.22±0.28	-2.426	<b>0.015</b>
Low-density lipoproteins (mg/dL)	2.71±0.76	2.65±0.74	2.629	<b>0.009</b>
Triglycerides (mg/dL)	1.47±1.12	1.40±1.14	2.049	<b>0.041</b>
Coronary heart disease				
Yes	183 (11.0)	249 (7.6)	37.880	<0.001
No	1486 (89.0)	3027 (92.4)		
Cataract extraction (OS)				
No	1801 (97.6)	3481 (99.4)	30.600	<0.001
Yes	44 (2.4)	22 (0.6)		
Cataract extraction (OD)				
No	1797 (97.6)	3480 (99.3)	27.935	<0.001
Yes	44 (2.4)	24 (0.7)		
Intraocular pressure OS (mm Hg)	12.32±2.70	12.31±2.29	0.091	0.927
Intraocular pressure OD (mm Hg)	11.76±2.91	11.80±2.32	-0.457	0.648
Spherical equivalent OS	0.06±2.09	0.11±1.76	-1.017	0.309
Spherical equivalent OD	0.06±1.97	0.07±1.84	-0.199	0.842
Axial length OS (mm)	22.78±0.95	22.82±0.91	-1.538	0.124
Axial length OD (mm)	22.85±0.98	22.87±1.08	-0.793	0.428
BCVA OS (logMAR)	0.60±0.26	0.56±0.26	4.437	<0.001
BCVA OD (logMAR)	0.60±0.26	0.57±0.26	3.570	<0.001

Statistics presented: mean±SD; n (%).  
Values with statistical significance are shown in boldface.  
BCVA, best-corrected visual acuity; OD, Oculus Dexter; OS, Oculus Sinister.

coronary heart disease ( $p<0.001$ ), higher cataract extraction ( $p<0.001$ ) and lower BCVA ( $p<0.001$ ) (table 1).


### Distribution of RNFL thickness

The average RNFL thickness in this study was  $113.46\pm 10.90$   $\mu\text{m}$ . And the thickest quadrant of pRNFL was the inferior quadrant, followed by the superior quadrant, the nasal quadrant and the temporal quadrant. The mean RNFL thickness was 2.11  $\mu\text{m}$  thicker in female than in male ( $p<0.001$ ). These sex-specific differences were statistically significant also in all temporal, nasal, superior and inferior ( $p<0.001$ ) quadrants. All segments were thicker in female than in male, as shown in figure 1. For the optic disc parameters, the mean (SD) disc,


cup and rim area were 2.14 (0.44) mm, 0.67 (0.39) mm and 1.46 (0.44) mm, respectively. Compared with male, female had significant smaller average cup-to-disc ratio ( $p<0.001$ ), cup volume ( $p<0.001$ ), AL ( $p<0.001$ ) and larger rim area ( $p<0.001$ ) (online supplemental tables 2 and 4).

### The relationship between RNFL and age

The normative distribution of RNFL thickness by age groups is described in table 2. Participants were divided into five age groups: less than 39, 40–49, 50–59, 60–69, over 70 years. For all age-specific groups, all inferior quadrant RNFL mean thickness was thicker than superior quadrant, followed by nasal and temporal quadrant mean RNFL. The four quadrants thickness



**Figure 1** Distribution of RNFL thickness by gender. This figure shows the distribution of RNFL in 16 segments near the optic disc of different genders. It can be seen that the distribution of RNFL thickness was the same as the overall. The RNFL of 16 segments was all thicker in women than in men. RNFL, retinal nerve fibre layer.



**Figure 2** Distribution of RNFL thickness at different ages. This figure shows the distribution of RNFL in 16 segments near the optic disc at different genders. It can be seen that the thickness quadrant of RNFL is the inferior quadrant, followed by the superior, nasal and temporal quadrant. The thickness of peripheral RNFL decreases with age but this trend is gradually reflected after 50 years old. RNFL, retinal nerve fibre layer.

sectors are shown in [table 2](#). Mean RNFL thickness was highest in the less than 39-year-old age group (115.22  $\mu$ m) vs 107.70  $\mu$ m in the over 70-year-old age group. A general trend toward thinner RNFL was noted with older age in 16 sectors is show in [figure 2](#) and online supplemental table 3. Of ONH parameters, AL decreased and disc area increased significantly with age ([table 2](#)), the post hoc tests after ANOVA among the age groups were shown in online supplemental tables 7 and 8. Mixed linear regression models were used to compare the sectionalised RNFL thickness between different age and gender groups. And the fan-shaped distribution of RNFL regions of different ages and genders was shown in [figure 3](#). With the main and interaction effect from age and gender were both considered, the results demonstrated that except for RNFL, the coefficients of interaction effect were non-significantly (online supplemental tables 1 and 3).

**Linear regression analysis of influencing factors of RNFL**

Univariate and multivariate linear regression models were used to estimate the independent associations between ocular

(history of cataract surgery, IOP, spherical equivalent (SE), AL, BCVA) and systemic (age, gender, educational level, history of smoking, BMI, hypertension and diabetes status, and LDL cholesterol) parameters with average RNFL thickness ([table 3](#)). In univariate linear regression analysis, thinner RNFL thickness was significantly associated with older age ( $p < 0.001$ ), male sex ( $p < 0.001$ ), absence of smoking ( $p < 0.001$ ), increased BMI value ( $BMI < 21$ ,  $p = 0.017$ ), increased BMI value ( $BMI > 30$ ,  $p = 0.033$ ), absence of diabetes ( $p = 0.002$ ), coronary heart disease ( $p < 0.001$ ), history of cataract surgery ( $p = 0.001$ ), higher IOP ( $p = 0.043$ ), lower SE ( $p < 0.001$ ), longer AL ( $p < 0.001$ ) and BCVA ( $p < 0.001$ ) ([table 3](#)).

In the multivariate analysis model, RNFL thickness was taken as the dependent variable, the significant ( $p < 0.05$ ) variables in univariate models were further included in the multivariate mixed linear model ([table 3](#)). We then dropped the variables that were no longer significantly associated with average RNFL thickness. In the multivariate model, thinner RNFL thickness was significantly associated with older age ( $p < 0.001$ ), male sex ( $p < 0.001$ ), increased BMI value ( $BMI > 30$ ,  $p = 0.018$ ), absence


**Table 2** Comparisons of sectionalised RNFL thickness between different age groups

Segment	Age					P value
	<39 (N=352)	40-49 (N=1728)	50-59 (N=2095)	60-69 (N=1647)	$\geq 70$ (N=550)	
RNFL thickness ( $\mu$ m)						
Average	115.22 $\pm$ 10.88	115.90 $\pm$ 10.25	113.95 $\pm$ 10.34	111.47 $\pm$ 10.75	107.80 $\pm$ 12.54	<0.001
Superior	143.28 $\pm$ 19.00	145.29 $\pm$ 17.89	141.81 $\pm$ 17.76	137.96 $\pm$ 18.39	133.22 $\pm$ 19.47	<0.001
Nasal	85.05 $\pm$ 14.78	85.14 $\pm$ 14.63	84.53 $\pm$ 15.01	83.42 $\pm$ 16.05	81.23 $\pm$ 16.22	<0.001
Inferior	149.61 $\pm$ 17.31	150.14 $\pm$ 17.27	147.73 $\pm$ 16.89	144.67 $\pm$ 17.50	139.40 $\pm$ 19.56	<0.001
Temporal	83.06 $\pm$ 12.54	83.03 $\pm$ 12.27	81.76 $\pm$ 12.32	79.84 $\pm$ 12.00	76.98 $\pm$ 14.60	<0.001
ONH parameters						
Rim area ( $mm^2$ )	1.69 $\pm$ 0.38	1.68 $\pm$ 0.47	1.69 $\pm$ 0.47	1.67 $\pm$ 0.49	1.71 $\pm$ 0.55	0.615
Disc area ( $mm^2$ )	2.37 $\pm$ 0.43	2.39 $\pm$ 0.48	2.43 $\pm$ 0.51	2.44 $\pm$ 0.48	2.49 $\pm$ 0.53	<0.001
Average CDR ( $mm^2$ )	0.33 $\pm$ 0.13	0.33 $\pm$ 0.13	0.34 $\pm$ 0.13	0.34 $\pm$ 0.13	0.33 $\pm$ 0.14	0.549
Cup volume ( $mm^3$ )	0.13 $\pm$ 0.15	0.15 $\pm$ 0.19	0.15 $\pm$ 0.18	0.14 $\pm$ 0.17	0.13 $\pm$ 0.17	0.228
Axial length (mm)	22.95 $\pm$ 1.19	22.98 $\pm$ 0.94	22.83 $\pm$ 0.93	22.75 $\pm$ 0.86	22.69 $\pm$ 0.80	<0.001

Statistics presented: mean $\pm$ SD; n (%).

Values with statistical significance are shown in boldface.

CDR, cup-to-disc ratio; ONH, optic nerve head; RNFL, retinal nerve fibre layer.



**Figure 3** Fan-shaped distribution of RNFL thickness by gender and different ages. The fan-shaped distribution can show that the RNFL thickness near the optic disc is the thinnest in TL1, TU1, NI1 and NU1 and the thickest in ST1 and IT1. The fan-shaped distribution of RNFL thickness showed a thinner trend in men and older people. RNFL, retinal nerve fibre layer.

of diabetes ( $p=0.009$ ), history of cataract surgery ( $p=0.001$ ), higher IOP ( $p=0.007$ ), lower SE ( $p<0.001$ ) and longer AL ( $p=0.048$ ). In addition, we supplemented the VIF for each covariate in the multivariate model, the VIFs were all below 10 (online supplemental table 5).

**DISCUSSION**

We evaluated the RNFL thickness around the optic disc and the parametric distribution of pRNFL measured by SD-OCT among the non-glaucomatous Asian population. This is the first population-based cohort study in China and showing the

discrepancy in the standard distribution of RNFL thickness in middle-aged and older populations. Quadrant thicknesses decreasing order: inferior, superior, nasal and temporal, respectively. We demonstrated that the RNFL was significantly thicker in the normal Asian population than in the European standard database using SD-OCT. We demonstrated that the RNFL thickness tends to be influenced by races. Therefore, errors may occur in measuring Asian individuals when the RNFL thickness standard database composed of other races (such as Caucasians) in OCT measurement. Moreover, various factors may also affect the RNFL thickness evaluation. RNFL thickness thinned with age, while women showed a tendency to be thicker in RNFL measurements. And other factors such as IOP, AL, cataract surgery, diabetes and others were also shown to be associated with RNFL thickness.

This study is a population-based cohort study (all data based on HES), which was statistically analysed to yield a mean RNFL thickness for this in Chinese population, and the average RNFL thickness of Chinese population is  $113.46 \pm 10.90 \mu\text{m}$ . The distribution of RNFL thickness was consistent with the distribution of RNFL in the majority of studies. As described by Zhao *et al*<sup>17</sup> measuring RNFL in subjects without optic nerve or retinal disease, the average RNFL was significantly higher in the inferior temporal quadrant than in the superior temporal quadrant, followed by the inferior nasal quadrant, the superior nasal quadrant, the temporal region and finally the nasal quadrant, respectively. This sequence of nasal quadrants bore a definite resemblance to the Singapore-Malay study<sup>18</sup> and the Singapore Chinese Eye Study.<sup>19,20</sup> However, the mean thickness of RNFL in our study population was thicker compared with the most of other studies. In the Gutenberg Health Study,<sup>21</sup> the overall mean

**Table 3** The univariate and multivariate analysis of relationship between demographic and biochemical characteristics and average RNFL thickness

Variables	Univariate analysis			Multivariate		
	B	SE	P value	B	SE	P value
Age (years)	-0.233	0.016	<0.001	-0.260	0.024	<0.001
Gender (female vs male)	2.043	0.346	<0.001	1.772	0.495	<0.001
High educational level	-0.576	0.973	0.554			
Smoking status	1.369	0.394	<0.001	-0.319	0.552	0.563
Body mass index ( $\text{kg}/\text{m}^2$ )*						
<21	-1.531	0.643	0.017	-0.822	0.690	0.234
27-30	0.745	0.426	0.080	0.668	0.461	0.147
$\geq 30$	1.144	0.537	0.033	1.403	0.593	0.018
Diastolic >140 (mm Hg)	2.287	7.222	0.751			
Systolic >90 (mm Hg)	7.903	5.897	0.180			
Diabetes	2.888	0.929	0.002	2.576	0.992	0.009
HbA1c (%)	-0.412	0.516	0.425			
HDL (normal vs abnormal)	0.028	0.383	0.942			
LDL (normal vs abnormal)	0.414	0.368	0.260			
TG (normal vs abnormal)	0.165	0.398	0.678			
Coronary heart disease	2.248	0.677	<0.001	0.562	0.706	0.426
Cataract extraction	-4.591	1.419	0.001	-5.845	1.633	<0.001
Intraocular pressure (mm Hg)	-0.110	0.054	0.043	-0.157	0.059	0.007
Spherical equivalent	0.734	0.076	<0.001	0.862	0.108	<0.001
Axial length	-0.513	0.112	<0.001	-0.240	0.121	0.048
BCVA (logMAR)	-4.270	0.593	<0.001	-0.703	0.724	0.331

Values with statistical significance are shown in boldface.

\*21-27 ( $\text{kg}/\text{m}^2$ ) was set as the reference category.

BCVA, best-corrected visual acuity; HbA1c, glycosylated hemoglobin; HDL, high density lipoprotein; LDL, Low Density Lipoprotein; RNFL, retinal nerve fibre layer; TG, triglyceride.

Table 4 Information of RNFL thickness from previous studies

Study	Year	OCT	Country	N	Age	RNFL thickness ( $\mu\text{m}$ )				
						Average	Temporal	Superior	Nasal	Inferior
Girkin <i>et al</i> <sup>24</sup>	2010	Stratus OCT	Africa	315	45.1 $\pm$ 13.3	103.7 $\pm$ 10.7	66.5 $\pm$ 11.1	128.8 $\pm$ 17.2	84.3 $\pm$ 17.2	135.1 $\pm$ 16.3
Girkin <i>et al</i> <sup>24</sup>	2010	Stratus OCT	Europe	290	47.7 $\pm$ 15.9	100.6 $\pm$ 10.9	71.5 $\pm$ 12.6	120.9 $\pm$ 17.5	80.8 $\pm$ 16.3	129.2 $\pm$ 17.4
Celebi and Mirza <sup>25</sup>	2013	Cirrus SD-OCT	Turkey	121	38.9 $\pm$ 11.2	97.0 $\pm$ 7.4	64.7 $\pm$ 6.4	119.2 $\pm$ 11.2	74.8 $\pm$ 8.2	129.3 $\pm$ 11.7
Alasil <i>et al</i> <sup>26</sup>	2013	Spectralis OCT	Mixed	190	53.7 $\pm$ 16.3	97.3 $\pm$ 9.6	70.6 $\pm$ 10.8	117.2 $\pm$ 16.3	75.0 $\pm$ 14.0	126.0 $\pm$ 15.8
Park <i>et al</i> <sup>27</sup>	2005	Stratus OCT	Korea	121	43.2 $\pm$ 13.9	112.7 $\pm$ 15.0	85.2 $\pm$ 17.9	137.5 $\pm$ 20.0	89.5 $\pm$ 22.2	138.1 $\pm$ 20.8
Budenz <i>et al</i> <sup>28</sup>	2007	Stratus OCT	America	328	47.4 $\pm$ 15.8	100.1 $\pm$ 11.6				
Schuster <i>et al</i> <sup>29</sup>	2016	Topcon 3D-OCT	Caucasian	306	38.8 $\pm$ 10.9		87.0 $\pm$ 11.0	124.0 $\pm$ 13.0	93.0 $\pm$ 15.4	138.0 $\pm$ 13.6
Cheung <i>et al</i> <sup>19</sup>	2011	Cirrus HD-OCT	Chinese	542	53.0 $\pm$ 6.4	97.6 $\pm$ 9.1	71.6 $\pm$ 11.2	123.0 $\pm$ 15.9	69.2 $\pm$ 10.8	126.8 $\pm$ 16.2
Sung <i>et al</i> <sup>30</sup>	2009	Stratus OCT	America	226	47.5 $\pm$ 15.9	100.8 $\pm$ 10.5				
Leung <i>et al</i> <sup>31</sup>	2004	Stratus OCT	China	107	53.0 $\pm$ 11.8	103.2 $\pm$ 10.0				
Kanamori <i>et al</i> <sup>32</sup>	2003	Humphrey OCT	Japan	144	46.3 $\pm$ 18.1	123.0 $\pm$ 11.6	101.0 $\pm$ 18.5	148.0 $\pm$ 18.4	96.0 $\pm$ 19.2	146.0 $\pm$ 19.3
Fujiwara <i>et al</i> <sup>9</sup>	2019	Topcon 3D-OCT	Japan	749	58.0 $\pm$ 10.0	102.3 $\pm$ 0.9				
Zangalli <i>et al</i> <sup>33</sup>	2018	Spectralis OCT	Brazil	220	44.0 $\pm$ 13.9	103.0 $\pm$ 10.4	67.7 $\pm$ 10.6		86.0 $\pm$ 13.8	
Perez <i>et al</i> <sup>34</sup>	2018	Cirrus OCT	Vietnam	151	60.8 $\pm$ 11.1	97.9 $\pm$ 9.2	69.7 $\pm$ 10.1	119.7 $\pm$ 15.1	73.4 $\pm$ 13.8	128.6 $\pm$ 15.4
Manassakorn <i>et al</i> <sup>35</sup>	2008	Stratus OCT	Thailand	250	44.7 $\pm$ 12.2	109.3 $\pm$ 0.7	75.1 $\pm$ 0.7	136.0 $\pm$ 1.0	83.6 $\pm$ 1.0	142.4 $\pm$ 1.1
Appukuttan <i>et al</i> <sup>36</sup>	2014	Spectralis OCT	India	150	20–75	101.4 $\pm$ 8.6	72.0 $\pm$ 7.7	125.3 $\pm$ 13.7	79.7 $\pm$ 12.1	128.3 $\pm$ 14.7
Al-Sa'ad <i>et al</i> <sup>37</sup>	2018	RTVue OCT	Jordan	148	60.0 $\pm$ 12.0	99.0 $\pm$ 11.0	82.0 $\pm$ 20.0	114.0 $\pm$ 20.0	75.0 $\pm$ 16.0	125.0 $\pm$ 20.0
Varma <i>et al</i> <sup>38</sup>	2003	OCT 2000	Latino	312	51.9 $\pm$ 9.8	132.7 $\pm$ 14.4	102.5 $\pm$ 19.0	157.7 $\pm$ 17.8	109.3 $\pm$ 19.1	159.8 $\pm$ 18.9
Kang <i>et al</i> <sup>39</sup>	2016	iVue-100 OCT	China	1811	7.1 $\pm$ 0.4	102.0 $\pm$ 0.2	80.2 $\pm$ 0.2	125.1 $\pm$ 0.3	75.9 $\pm$ 0.3	126.8 $\pm$ 0.3
Méndez <i>et al</i> <sup>40</sup>	2017	Spectralis OCT	France	427	81.6 $\pm$ 4.2	86.8 $\pm$ 13.7	67.2 $\pm$ 13.4		66.2 $\pm$ 16.4	
Kanno <i>et al</i> <sup>41</sup>	2010	EG-SCANNER	Japan	460	44.0 $\pm$ 14.5	111.8 $\pm$ 10.0				
Gupta <i>et al</i> <sup>42</sup>	2015	Cirrus HD-OCT	Singapore	843	66.7 $\pm$ 10.4	92.9 $\pm$ 11.4	67.4 $\pm$ 11.4	116.1 $\pm$ 18.4	69.7 $\pm$ 11.3	118.6 $\pm$ 19.8
Rougier <i>et al</i> <sup>22</sup>	2015	Spectralis OCT	French	210	81.0 $\pm$ 3.6	91.4 $\pm$ 12.6	70.1 $\pm$ 14.7		69.1 $\pm$ 16.3	
Ho <i>et al</i> <sup>43</sup>	2019	Cirrus HD-OCT	Chinese	1371	57.4 $\pm$ 7.0	95.7 $\pm$ 9.6	71.4 $\pm$ 12.2	119.9 $\pm$ 16.7	68.1 $\pm$ 10.9	123.4 $\pm$ 16.7
Ho <i>et al</i> <sup>43</sup>	2019	Cirrus HD-OCT	Malay	1303	60.6 $\pm$ 8.6	94.9 $\pm$ 10.6	67.8 $\pm$ 11.3	118.6 $\pm$ 17.3	71.0 $\pm$ 10.9	122.4 $\pm$ 18.3
Ho <i>et al</i> <sup>43</sup>	2019	Cirrus HD-OCT	Indian	1801	60.7 $\pm$ 7.8	87.3 $\pm$ 10.6	59.2 $\pm$ 10.9	108.8 $\pm$ 16.5	69.1 $\pm$ 11.1	112.3 $\pm$ 17.3
Zhu <i>et al</i> <sup>44</sup>	2013	iVue-100 OCT	China	1955	12.3 $\pm$ 0.6	103.1 $\pm$ 9.0	83.0 $\pm$ 10.6	126.2 $\pm$ 15.2	73.8 $\pm$ 13.9	129.3 $\pm$ 14.9
Wang <i>et al</i> <sup>45</sup>	2018	Spectralis OCT	China	1440	11.9 $\pm$ 3.5	101.3 $\pm$ 9.2	85.2 $\pm$ 14.3		61.7 $\pm$ 20.4	
Nousome <i>et al</i> <sup>46</sup>	2021	Cirrus HD-OCT	Mixed	6133	60.1 $\pm$ 7.4	95.1 $\pm$ 10.1	65.6 $\pm$ 12.2	118.6 $\pm$ 16.5	71.2 $\pm$ 11.1	125.1 $\pm$ 17.3
Malik <i>et al</i> <sup>47</sup>	2012	Stratus OCT	India	150	42.6 $\pm$ 13.6	101.1 $\pm$ 10.1	65.7 $\pm$ 12.1	125.8 $\pm$ 16.5	83.6 $\pm$ 17.4	127.5 $\pm$ 15.6
Bendschneider <i>et al</i> <sup>8</sup>	2010	Spectralis OCT	America	170	20–78	97.2 $\pm$ 9.7	68.8 $\pm$ 11.1	118.0 $\pm$ 14.5	76.4 $\pm$ 15.0	123.7 $\pm$ 16.4
Wang <i>et al</i> <sup>48</sup>	2013	iVue OCT	China	1654	66.2 $\pm$ 9.9	103.2 $\pm$ 12.6	79.8 $\pm$ 12.2	126.1 $\pm$ 19.1	75.1 $\pm$ 12.6	131.4 $\pm$ 20.6
Hashemi <i>et al</i> <sup>49</sup>	2017	Cirrus HD-OCT	Iran	3084	54.3 $\pm$ 5.6	92.5 $\pm$ 0.3	65.5 $\pm$ 0.4	111.2 $\pm$ 0.5	74.8 $\pm$ 0.8	118.9 $\pm$ 0.6
Thapa <i>et al</i> <sup>50</sup>	2014	Spectralis OCT	Nepal	156	38.9 $\pm$ 17.0	102.6 $\pm$ 9.6	70.7 $\pm$ 15.5	129.5 $\pm$ 15.1	76.6 $\pm$ 12.0	134.5 $\pm$ 17.2
Ismail <i>et al</i> <sup>51</sup>	2019	Spectralis OCT	South Africa	132	41.3 $\pm$ 12.5	108.7 $\pm$ 10.7	74.8 $\pm$ 10.3		77.7 $\pm$ 14.6	
Zhao <i>et al</i> <sup>17</sup>	2014	Spectralis OCT	China	2548	63.5 $\pm$ 9.1	102.0 $\pm$ 11.0	76.0 $\pm$ 13.0		73.0 $\pm$ 15.0	
Mashige and Oduntan <sup>52</sup>	2016	iVue-100 OCT	South Africa	600	10–66	110.0 $\pm$ 7.4	73.6 $\pm$ 15.7	132.0 $\pm$ 10.5	87.2 $\pm$ 13.2	135.1 $\pm$ 9.7
Budenz <i>et al</i> <sup>53</sup>	2005	Stratus OCT	America	109	42.8 $\pm$ 14.6	104.8 $\pm$ 10.7	75.1 $\pm$ 17.2	130.9 $\pm$ 18.2	79.8 $\pm$ 17.2	133.4 $\pm$ 18.7
Hoffmann <i>et al</i> <sup>21</sup>	2018	Spectralis OCT	Germany	1974	40–80	96.0 $\pm$ 10.3	68.8 $\pm$ 12.9		72.2 $\pm$ 15.1	
Feuer <i>et al</i> <sup>54</sup>	2011	Stratus OCT	America	425	46.0 $\pm$ 15.0	104.7 $\pm$ 10.8				

RNFL, retinal nerve fibre layer; SD-OCT, spectral-domain optical coherence tomography.

thickness of RNFL was 96.0  $\mu\text{m}$ , and the Singapore Chinese Eye Study<sup>19, 20</sup> showed an overall mean RNFL thickness of 96.2  $\mu\text{m}$ . Another population-based cohort study from France found a mean RNFL thickness of 91.4  $\mu\text{m}$ ,<sup>22</sup> all these figures were obviously lower than our findings. In contrast, the overall average pRNFL thickness in the Beijing Eye Study was 103.2  $\mu\text{m}$  (SD-OCT), which shown similar results as our study, although it is also lower than our results. The differences in RNFL measurements can be partially explained by the different instruments applied and various age groups of the population covered. However, even for the same examination equipment, the RNFL thickness in our study population compared with other European or African populations is still thicker.<sup>23</sup> The RNFL thickness of other studies was shown in table 4,<sup>8, 9, 17, 19, 21, 22, 24–54</sup> it can be

roughly seen that the RNFL thickness measured by Cirrus OCT is generally lower than that measured by other OCT equipment according to this table, which may come from the differences in the built-in RNFL thickness standard database of OCT machines by various companies. The reason for the differences is that these companies shall select diverse individuals with different proportions when setting the standard database of the RNFL thickness. For instance, Heidelberg company selects all Caucasians, Topcon selects all Asians, and other companies select a wide range of people (online supplemental table 6). Thus, when the difference of RNFL thickness on account of various ethnic groups is judged, the database of the instrument itself must show the certain impact, suggesting that the instrument differences shall be taken into consideration in judging RNFL thickness

in ethnic groups. Therefore, the international standard RNFL database is not suitable for the application in the Chinese population, which may bring about a lot of missed diagnosis (taking the glaucoma as the normal one), or misdiagnosis (taking normal one as glaucoma) potentially, suggesting the need to establish additional appropriate RNFL thickness measures in the Chinese population.

Age is one of the important factors affecting the distribution of RNFL thickness, which has been confirmed in several studies. In the study conducted by Hashemi *et al*, the overall average RNFL thickness as well as the mean thickness of all quadrants went down dramatically with age,<sup>49</sup> while the decrease of RNFL thickness<sup>8 25</sup> was in connection with the ageing in other studies also. In addition to OCT studies, some studies have shown age-related RNFL thinning by histological analysis.<sup>26</sup> The RNFL thickness decreases by 2–4  $\mu\text{m}$  per 10 years of ageing has been observed in individuals over 50 years, whereas it nearly hasn't been found under 50 years old or even younger people in accordance with these results. The rationale for the effect of age on the RNFL may be due to decreasing blood supply to the fundus and senescence apoptosis of the optic nerve with advancing age.<sup>21</sup> The additional systemic factors effecting the measurement of RNFL thickness shall not be ruled out as the age grows. Nevertheless, there definitely exists the correlation of a decrease in RNFL thickness with an increase in age. Therefore, RNFL thickness for different age groups may need to be considered separately, and synthesising data from multiple studies and deriving a pattern of decay in RNFL thickness with age in order to advance the use of measuring RNFL thickness in clinical practice (figure 3).

Our study also found that women tended to be thicker than men in both overall average RNFL and RNFL thickness in all orientations. Rougier *et al* found that women tended to have a thicker RNFL than men overall and across all temporal ranges, although this was only significant in the inferior temporal segment.<sup>22</sup> Even higher RNFL values in females and even female children have been reported in some population-based studies and clinical-based studies.<sup>55</sup> However, some studies have failed to find an association between RNFL and gender in adult populations, as recently Girkin *et al*<sup>23</sup> found that pRNFL or macular parameters obtained using SD-OCT (RTVue) did not differ between females and males. Although this correlation needs to be explored in further studies, however, the gender should be taken into consideration when the RNFL is measured and analysed (figure 3).

As for the correlation analysis of systemic factors with RNFL, the lower BMI (BMI  $\geq 30$ ) and no diabetes were correlated linearly with RNFL thinning in both univariate and multivariate linear analyses. In terms of the BMI, the higher BMI is related to the increase of intracranial pressure, and the increase of intracranial pressure can lead to changes such as optic disc oedema, making the measured RNFL thicker,<sup>56</sup> then showing the lower BMI and the decline of RNFL are definitely related. Most of the studies reported thinner RNFL in patients with diabetes,<sup>57</sup> which may be on account of diabetes-induced microangiopathy and ischaemia, and the correlation between reduced RNFL visibility and elevated blood glucose concentrations was associated with loss of retinal nerve fibres in patients with diabetic retinopathy.<sup>58</sup> However, there is no evidence that diabetes is related to the thinning of RNFL in our study, which may be due to the proliferation and retinal oedema caused by diabetes. Our study did not find a linear relationship between elevated blood pressure as well as coronary artery disease and RNFL, although previous studies have also reported conflicting results regarding the effect of hypertension on RNFL.<sup>59</sup> Nevertheless, the hypertensive

patients facing with fluctuations in blood pressure or medication control may give rise to the negative results and hypertensive fundus lesions may affect RNFL thickness in our study according to our speculation. This study did not find an association between smoking and RNFL, Mauschitz *et al* still suggested such an association, although its relationship of smoking with increased or decreased RNFL is unclear.<sup>60</sup> Other central nervous system conditions similar to stroke, apoplexy, and dementia have also been shown to be associated with RNFL thickness. Quite a few systemic conditions can affect RNFL thickness measurements, which is a factor that cannot be ignored when building a standard RNFL database. Broadly speaking, however, it appears that the vast majority of these systemic factors affect the status of vascular microcirculation throughout the body.

Generally, elevated IOP, decreased SE and AL were also connected with the declining of RNFL in both univariate and multivariate linear regression analyses after the cataract surgery. As for the IOP, a host of studies suggest that the decreasing of RNFL bears the association with higher IOP, which is one of the major potential factors for glaucoma after all.<sup>10 31</sup> However, the study by Mauschitz *et al* also confirmed that high IOP was associated with a reduction in RNFL thickness even exclusion<sup>60</sup> of the patients with known glaucoma. The association between a longer AL and a thinner RNFL is another finding of this study, which has been confirmed by previous studies,<sup>61</sup> whereas the relationship between AL and RNFL thickness was not the same across quadrants. Yoo *et al* showed that increased AL was associated with decreased RNFL thickness in the upper, lower and nasal quadrants and increased RNFL thickness in the temporal quadrant.<sup>62</sup> Kang *et al* studied Chinese individuals living in Korea using SD-OCT and reported similar results except for the temporal quadrant observation.<sup>63</sup> In a European study, Savini *et al* demonstrated a remarkable association that a larger AL was significantly associated with lower RNFL thickness in all quadrants. Whereas Hirasawa *et al* used a simple regression model to show that thickness was directly correlated with AL in all quadrants, while this kind of relation was only present in the temporal quadrant in their multiple regression model.<sup>12</sup> Although the vast majority of studies have found a relationship between AL and RNFL, this relationship is nevertheless influenced by many factors. As for the SE, the decline in both SE and RNFL may always occur among the elder for the majority of them having the vision loss. Moreover, the ocular disease history is also a factor that affects RNFL and these factors have to be taken into account in the measurement of RNFL.

This study also found a relationship between cataract surgery and thinner RNFL thickness. We speculated that this should be due to the fact that the average age of people with cataract surgery (with average age 56 years) is much older than that of people without cataract surgery (with average age 70y). However, in addition to age, some studies have also proposed the relationship between cataract surgery and RNFL thickness. The macular and RNFL thickness face with a rise after cataract extraction partly on account of the quality of scanning image developing proposed by Pařová and Skorkovská,<sup>64</sup> while the RNFL and macular measurement conducted by SD-OCT shall be influential by the presence of cataract and a better OCT instrument may reduce this effect as much as possible suggested by Bambo *et al*.<sup>65</sup> Therefore, the cataract surgery makes changes on the penetration of refractive medium, then further impacting the measurement of RNFL thickness.

Our study has several advantages. First, this large population-based study involved 3509 participants all coming from the general population instead of that attending the hospital. Second,

the ocular examination excluded the participants with ocular disease, which made the RNFL measurements more accurate, and could be representative of the normal database. Third, the SD-OCT evaluation was performed by experienced technicians on the same machine following a standardised protocol, the bias of measurement has been avoided. And applied SD-OCT provided the best image quality and reproducibility currently available. Finally, we simultaneously evaluated many possible risk factors associated with RNFL thickness.

However, there are still some limitations of our study: although Chinese people were selected to explore the thickness of RNFL in this study, the thickness of RNFL also depends on other factors. As the rural population was selected in this study, the prevalence of myopia was relatively low for the participation of them in the study, and the AL (22.8 mm) was shorter than that in other studies involving Chinese population (Ho *et al*<sup>43</sup>), further affecting the evaluation of the results. we cannot exclude the possibility that subjects with refractive media opacities could influence the results. However, routine clinical examinations would also face such a situation, which is unavoidable with SD-OCT examinations. Second, the assessment of RNFL visibility and local RNFL defects is subjective and therefore it depends on the technicians. In our study, however, all inspectors had been well trained, but it is inevitable that inspectors have varying experience and examination skills. Third, individuals excluded due to poor OCT scans were older and they were more likely to have hypertension, diabetes and hyperlipidaemia. Therefore, these biases could not be completely excluded in our final sample.

In conclusion, we measured the thickness and distribution of RNFL in the Chinese population, the results in our study demonstrate that the RNFL in the Chinese population is thicker than we widely use in normal SD-OCT database. Using the European RNFL database to evaluate the Chinese population may miss some glaucoma patients. Besides, it shall not be ignored in RNFL evaluation that various factors may have some influential on RNFL thickness and these related parameters also widely applied in the diagnosis of glaucoma and other neurological disorders. Hence, our findings further emphasise the need to demonstrate ethnic differences in RNFL thickness and the specificity of associated ocular and systemic factors, as well as to develop better normative databases worldwide.

**Acknowledgements** The authors wish to thank the Beijing Tongren Hospital, Handan Eye Hospital and all survey participants are greatly appreciated for their valuable contributions.

**Contributors** JW and YD completed the design, data collection, data analysis and manuscript writing of this study. They contributed equally as co-first authors. CL participated in the data collection and method design of this study, YZ participated in the research design and statistical analysis of this study, WC participated in the statistical analysis of this study and the grammar modification of the manuscript, and QP participated in the design of the statistical method of this study, YZ participated in the later revision of this study. NW was the corresponding author of this study, is responsible for the overall content as guarantor, accepts full responsibility for the finished work and the conduct of the study, had access to the data, and controlled the decision to publish.

**Funding** The study is supported by National Natural Science Foundation of China (GZR-2012-009); Beijing Traditional Chinese Medicine Technology Development Fund Project (JJ2018-50).

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. This study obtained the ethical clearance from the Beijing Tongren Hospital Ethical Committee, according to the

Helsinki Declaration. Written informed consent was obtained from all participants. For those who were illiterate or blindness, we read the consent form to them and asked them to mark the consent form with an inked forefinger, and the consent form with an inked forefinger that obtained from illiterate participant was also approved by the Ethics committee. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available on reasonable request. Data may be obtained from a third party and are not publicly available.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

#### ORCID iDs

Yifan Du <http://orcid.org/0000-0002-6823-2719>

Yehong Zhuo <http://orcid.org/0000-0003-3247-7199>

Ningli Wang <http://orcid.org/0000-0002-8933-4482>

#### REFERENCES

- Harasymowicz P, Birt C, Gooi P, *et al*. Medical management of glaucoma in the 21st century from a Canadian perspective. *J Ophthalmol* 2016;2016:6509809.
- Hoffmann EM, Medeiros FA, Sample PA, *et al*. Relationship between patterns of visual field loss and retinal nerve fiber layer thickness measurements. *Am J Ophthalmol* 2006;141:463–71.
- Kerrigan-Baumrind LA, Quigley HA, Pease ME, *et al*. Number of ganglion cells in glaucoma eyes compared with threshold visual field tests in the same persons. *Invest Ophthalmol Vis Sci* 2000;41:741–8.
- Höhn R, Kottler U, Peto T, *et al*. The ophthalmic branch of the Gutenberg health study: study design, cohort profile and self-reported diseases. *PLoS One* 2015;10:e0120476.
- Huang D, Swanson EA, Lin CP, *et al*. Optical coherence tomography. *Science* 1991;254:1178–81.
- Gabriele ML, Wollstein G, Ishikawa H, *et al*. Optical coherence tomography: history, current status, and laboratory work. *Invest Ophthalmol Vis Sci* 2011;52:2425–36.
- Hindle JV. Ageing, neurodegeneration and Parkinson's disease. *Age Ageing* 2010;39:156–61.
- Bendschneider D, Tornow RP, Horn FK, *et al*. Retinal nerve fiber layer thickness in normals measured by spectral domain OCT. *J Glaucoma* 2010;19:475–82.
- Fujiwara K, Yasuda M, Hata J, *et al*. Glucose tolerance levels and Circumpapillary retinal nerve fiber layer thickness in a general Japanese population: the Hisayama study. *Am J Ophthalmol* 2019;205:140–6.
- Li S, Wang X, Li S, *et al*. Evaluation of optic nerve head and retinal nerve fiber layer in early and advance glaucoma using frequency-domain optical coherence tomography. *Graefes Arch Clin Exp Ophthalmol* 2010;248:429–34.
- Kostanyan T, Wollstein G, Schuman JS. New developments in optical coherence tomography. *Curr Opin Ophthalmol* 2015;26:110–5.
- Hirasawa H, Tomidokoro A, Araie M, *et al*. Peripapillary retinal nerve fiber layer thickness determined by spectral-domain optical coherence tomography in ophthalmologically normal eyes. *Arch Ophthalmol* 2010;128:1420–6.
- Cao K, Hao J, Zhang Y, *et al*. Design, methodology, and preliminary results of the follow-up of a population-based cohort study in rural area of northern China: Handan eye study. *Chin Med J* 2019;132:2157–67.
- Foster PJ, Buhrmann R, Quigley HA, *et al*. The definition and classification of glaucoma in prevalence surveys. *Br J Ophthalmol* 2002;86:238–42.
- Chobanian AV, Bakris GL, Black HR, *et al*. The seventh report of the joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *JAMA* 2003;289:2560–72.
- American Diabetes Association. 2. Classification and Diagnosis of Diabetes: *Standards of Medical Care in Diabetes-2019*. *Diabetes Care* 2019;42:S13–28.
- Zhao L, Wang Y, Chen CX, *et al*. Retinal nerve fibre layer thickness measured by spectralis spectral-domain optical coherence tomography: the Beijing eye study. *Acta Ophthalmol* 2014;92:e35–41.



- 18 Gupta P, Cheung CY, Baskaran M, *et al.* Relationship between peripapillary choroid and retinal nerve fiber layer thickness in a population-based sample of Nonglaucomatous eyes. *Am J Ophthalmol* 2016;161:4–11.
- 19 Cheung CY, Chen D, Wong TY, *et al.* Determinants of quantitative optic nerve measurements using spectral domain optical coherence tomography in a population-based sample of non-glaucomatous subjects. *Invest Ophthalmol Vis Sci* 2011;52:9629–35.
- 20 Cheng C-Y, Allingham RR, Aung T, *et al.* Association of common SIX6 polymorphisms with peripapillary retinal nerve fiber layer thickness: the Singapore Chinese eye study. *Invest Ophthalmol Vis Sci* 2014;56:478–83.
- 21 Hoffmann EM, Schmidtmann I, Siouli A, *et al.* The distribution of retinal nerve fiber layer thickness and associations with age, refraction, and axial length: the Gutenberg health study. *Graefes Arch Clin Exp Ophthalmol* 2018;256:1685–93.
- 22 Rougier M-B, Korobelnik J-F, Malet F, *et al.* Retinal nerve fibre layer thickness measured with SD-OCT in a population-based study of French elderly subjects: the Alienor study. *Acta Ophthalmol* 2015;93:539–45.
- 23 Girkin CA, McGwin G, Sinai MJ, *et al.* Variation in optic nerve and macular structure with age and race with spectral-domain optical coherence tomography. *Ophthalmology* 2011;118:2403–8.
- 24 Girkin CA, Sample PA, Liebmann JM, *et al.* African descent and glaucoma evaluation study (ADAGES): II. ancestry differences in optic disc, retinal nerve fiber layer, and macular structure in healthy subjects. *Arch Ophthalmol* 2010;128:541–50.
- 25 Celebi ARC, Mirza GE. Age-Related change in retinal nerve fiber layer thickness measured with spectral domain optical coherence tomography. *Invest Ophthalmol Vis Sci* 2013;54:8095–103.
- 26 Alasil T, Wang K, Keane PA, *et al.* Analysis of normal retinal nerve fiber layer thickness by age, sex, and race using spectral domain optical coherence tomography. *J Glaucoma* 2013;22:532–41.
- 27 Park JJ, Oh DR, Hong SP, *et al.* Asymmetry analysis of the retinal nerve fiber layer thickness in normal eyes using optical coherence tomography. *Korean J Ophthalmol* 2005;19:281–7.
- 28 Budenz DL, Anderson DR, Varma R, *et al.* Determinants of normal retinal nerve fiber layer thickness measured by Stratus OCT. *Ophthalmology* 2007;114:1046–52.
- 29 Schuster AK-G, Fischer JE, Vossmerbaeumer C, *et al.* Determinants of peripapillary retinal nerve fiber layer thickness regarding ocular and systemic parameters - the MIPH Eye&Health Study. *Graefes Arch Clin Exp Ophthalmol* 2016;254:2011–6.
- 30 Sung KR, Wollstein G, Bilionick RA, *et al.* Effects of age on optical coherence tomography measurements of healthy retinal nerve fiber layer, macula, and optic nerve head. *Ophthalmology* 2009;116:1119–24.
- 31 Leung CKS, Yung WH, Ng ACK, *et al.* Evaluation of scanning resolution on retinal nerve fiber layer measurement using optical coherence tomography in normal and glaucomatous eyes. *J Glaucoma* 2004;13:479–85.
- 32 Kanamori A, Escano MFT, Eno A, *et al.* Evaluation of the effect of aging on retinal nerve fiber layer thickness measured by optical coherence tomography. *Ophthalmologica* 2003;217:273–8.
- 33 Zangalli CES, Reis ASC, Vianna JR, *et al.* Interocular asymmetry of minimum rim width and retinal nerve fiber layer thickness in healthy Brazilian individuals. *J Glaucoma* 2018;27:1136–41.
- 34 Perez CI, Chansangpetch S, Thai A, *et al.* Normative database and Color-code agreement of peripapillary retinal nerve fiber layer and macular ganglion Cell-inner plexiform layer thickness in a Vietnamese population. *J Glaucoma* 2018;27:665–73.
- 35 Manassakorn A, Chaidaroon W, Ausayakhun S, *et al.* Normative database of retinal nerve fiber layer and macular retinal thickness in a Thai population. *Jpn J Ophthalmol* 2008;52:450–6.
- 36 Appukuttan B, Giridhar A, Gopalakrishnan M, *et al.* Normative spectral domain optical coherence tomography data on macular and retinal nerve fiber layer thickness in Indians. *Indian J Ophthalmol* 2014;62:316–21.
- 37 Al-Sa'ad MM, Shatarat AT, Amarin JZ, *et al.* Normative values of peripapillary retinal nerve fiber layer thickness in a middle Eastern population. *J Ophthalmol* 2018;2018:7238464.
- 38 Varma R, Bazzaz S, Lai M. Optical tomography-measured retinal nerve fiber layer thickness in normal Latinos. *Invest Ophthalmol Vis Sci* 2003;44:3369–73.
- 39 Kang M-T, Li S-M, Li H, *et al.* Peripapillary retinal nerve fibre layer thickness and its association with refractive error in Chinese children: the Anyang childhood eye study. *Clin Exp Ophthalmol* 2016;44:701–9.
- 40 Méndez-Gómez JL, Rougier M-B, Tellouck L, *et al.* Peripapillary retinal nerve fiber layer thickness and the evolution of cognitive performance in an elderly population. *Front Neurol* 2017;8:93.
- 41 Kanno M, Nagasawa M, Suzuki M, *et al.* Peripapillary retinal nerve fiber layer thickness in normal Japanese eyes measured with optical coherence tomography. *Jpn J Ophthalmol* 2010;54:36–42.
- 42 Gupta S, Zivadinov R, Ramanathan M, *et al.* Optical coherence tomography and neurodegeneration: are eyes the windows to the brain? *Expert Rev Neurother* 2016;16:765–75.
- 43 Ho H, Tham Y-C, Chee ML, *et al.* Retinal nerve fiber layer thickness in a multiethnic normal Asian population: the Singapore epidemiology of eye diseases study. *Ophthalmology* 2019;126:702–11.
- 44 Zhu B-D, Li S-M, Li H, *et al.* Retinal nerve fiber layer thickness in a population of 12-year-old children in central China measured by iVue-100 spectral-domain optical coherence tomography: the Anyang childhood eye study. *Invest Ophthalmol Vis Sci* 2013;54:8104–11.
- 45 Wang C-Y, Zheng Y-F, Liu B, *et al.* Retinal nerve fiber layer thickness in children: the Gobi desert children eye study. *Invest Ophthalmol Vis Sci* 2018;59:5285–91.
- 46 Nousome D, Mckean-Cowdin R, Richter GM, *et al.* Retinal nerve fiber layer thickness in healthy eyes of black, Chinese, and Latino Americans: a population-based multiethnic study. *Ophthalmology* 2021;128:1005–15.
- 47 Malik A, Singh M, Arya SK, *et al.* Retinal nerve fiber layer thickness in Indian eyes with optical coherence tomography. *Nepal J Ophthalmol* 2012;4:59–63.
- 48 Wang YX, Pan Z, Zhao L, *et al.* Retinal nerve fiber layer thickness. The Beijing eye study 2011. *PLoS One* 2013;8:e66763.
- 49 Hashemi H, Khabazkhoob M, Nabovati P, *et al.* Retinal nerve fibre layer thickness in a general population in Iran. *Clin Exp Ophthalmol* 2017;45:261–9.
- 50 Thapa M, Khanal S, Shrestha GB, *et al.* Retinal nerve fibre layer thickness in a healthy Nepalese population by spectral domain optical coherence tomography. *Nepal J Ophthalmol* 2014;6:131–9.
- 51 Ismail S, Ally N, Alli HD. Retinal nerve fiber layer thickness in a normal black South African population. *Eye* 2020;34:1426–31.
- 52 Mashige KP, Oduntan OA. Retinal nerve fibre layer thickness values and their associations with ocular and systemic parameters in black South Africans. *Afr Health Sci* 2016;16:1188–94.
- 53 Budenz DL, Michael A, Chang RT, *et al.* Sensitivity and specificity of the StratusOCT for perimetric glaucoma. *Ophthalmology* 2005;112:3–9.
- 54 Feuer WJ, Budenz DL, Anderson DR, *et al.* Topographic differences in the age-related changes in the retinal nerve fiber layer of normal eyes measured by Stratus optical coherence tomography. *J Glaucoma* 2011;20:133–8.
- 55 Balazsi AG, Rootman J, Drance SM, *et al.* The effect of age on the nerve fiber population of the human optic nerve. *Am J Ophthalmol* 1984;97:760–6.
- 56 Jonas JB, Nangia V, Gupta R, *et al.* Retinal nerve fibre layer cross-sectional area, neuroretinal rim area and body mass index. *Acta Ophthalmol* 2014;92:e194–9.
- 57 Carpineto P, Toto L, Aloia R, *et al.* Neuroretinal alterations in the early stages of diabetic retinopathy in patients with type 2 diabetes mellitus. *Eye* 2016;30:673–9.
- 58 Zhang Y, Xu L, Zhang L, *et al.* Ophthalmoscopic assessment of the retinal nerve fiber layer. The Beijing eye study. *PLoS One* 2013;8:e62022.
- 59 Topouzis F, Coleman AL, Harris A, *et al.* Association of blood pressure status with the optic disk structure in non-glaucoma subjects: the Thessaloniki eye study. *Am J Ophthalmol* 2006;142:60–7.
- 60 Mauschitz MM, Bonnemaier PWM, Diers K, *et al.* Systemic and ocular determinants of peripapillary retinal nerve fiber layer thickness measurements in the European eye epidemiology (E3) population. *Ophthalmology* 2018;125:1526–36.
- 61 Chen C-Y, Huang EJ-C, Kuo C-N, *et al.* The relationship between age, axial length and retinal nerve fiber layer thickness in the normal elderly population in Taiwan: the Chiayi eye study in Taiwan. *PLoS One* 2018;13:e0194116.
- 62 Yoo YC, Lee CM, Park JH. Changes in peripapillary retinal nerve fiber layer distribution by axial length. *Optom Vis Sci* 2012;89:4–11.
- 63 Kang SH, Hong SW, Im SK, *et al.* Effect of myopia on the thickness of the retinal nerve fiber layer measured by Cirrus HD optical coherence tomography. *Invest Ophthalmol Vis Sci* 2010;51:4075–83.
- 64 Pašová P, Skorkovská K. [The Effect of Cataract Surgery on the Reproducibility and Outcome of Optical Coherence Tomography Measurements of Macular and Retinal Nerve Fibre Layer Thickness]. *Cesk Slov Oftalmol* 2016;72:20–6.
- 65 Bambo MP, Garcia-Martin E, Otin S, *et al.* Influence of cataract surgery on repeatability and measurements of spectral domain optical coherence tomography. *Br J Ophthalmol* 2014;98:52–8.

**Supplemental table 1. Comparisons of RNFL thickness between age and gender subgroups**

Segment	Age		Gender		Age*Gender	
	B	P-value	B	P-value	B	P-value
<b>RNFL thickness (<math>\mu\text{m}</math>)</b>						
Average	-0.188	<0.001	3.597	0.047	-0.031	0.336
Superior	-0.245	0.003	5.742	0.047	-0.067	0.195
Nasal	-0.024	0.715	4.755	0.043	-0.061	0.140
Inferior	-0.278	<0.001	3.950	0.163	-0.020	0.689
Temporal	-0.208	<0.001	-0.565	0.774	0.035	0.314
<b>ONH parameters</b>						
Rim area (mm <sup>2</sup> )	0.004	0.078	0.237	0.003	-0.003	0.077
Disc area (mm <sup>2</sup> )	0.003	0.190	-0.003	0.974	<0.001	0.807
Average CDR (mm <sup>2</sup> )	-0.001	0.279	0.065	0.065	<0.001	0.227
Cup volume (mm <sup>3</sup> )	-0.001	0.085	0.003	0.003	0.001	0.164
Axial length (mm)	-0.011	0.006	0.002	0.002	0.001	0.648
<b>16 sections (<math>\mu\text{m}</math>)</b>						
TU1	-0.164	0.003	-0.539	0.784	0.032	0.360
TU2	-0.198	0.014	4.199	0.134	0.006	0.902
ST2	-0.193	0.069	10.527	0.005	-0.084	0.205
ST1	-0.388	<0.001	1.272	0.737	-0.018	0.784
SN1	-0.232	0.048	6.010	0.144	-0.080	0.273
SN2	-0.175	0.081	5.853	0.094	-0.081	0.191
NU2	-0.124	0.164	1.581	0.612	-0.025	0.648
NU1	-0.035	0.602	3.296	0.155	-0.045	0.274
NL1	0.077	0.207	6.742	0.002	-0.090	0.018
NL2	-0.014	0.863	7.576	0.008	-0.091	0.076
IN2	-0.127	0.214	3.661	0.306	-0.020	0.756
IN1	-0.221	0.073	2.676	0.536	0.001	0.988
IT1	-0.382	<0.001	3.988	0.286	-0.028	0.675
IT2	-0.431	<0.001	3.991	0.311	-0.010	0.885
TL2	-0.288	<0.001	-1.869	0.488	0.044	0.355
TL1	-0.164	<0.001	-3.129	0.050	0.046	0.102

**Supplemental table 2. Comparisons of sectionalized RNFL thickness between male and female in 16 sections.**

Segment	Female (N=3819)	Male (N=3205)	t	P-value
<b>16 sections (µm)</b>				
TU1	73.99±12.92	72.69±12.7	4.279	< 0.001
TU2	101.93±18.62	97.24±17.50	10.832	< 0.001
ST2	140.08±23.73	133.93±23.85	10.746	< 0.001
ST1	156.10±24.37	155.57±24.37	0.890	0.372
SN1	139.34±26.59	137.58±26.39	2.761	0.006
SN2	133.87±23.82	132.21±22.80	2.776	0.006
NU2	102.53±21.04	102.20±19.61	0.670	0.501
NU1	74.26±15.34	73.34±14.70	2.564	0.010
NL1	69.79±13.82	67.95±13.73	5.544	< 0.001
NL2	92.71±19.58	90.02±18.79	5.850	< 0.001
IN2	129.98±23.40	127.16±23.22	4.874	< 0.001
IN1	153.88±27.41	150.89±26.93	4.569	< 0.001
IT1	169.39±24.05	166.53±23.34	5.036	< 0.001
IT2	141.13±26.01	137.34±24.99	6.192	< 0.001
TL2	87.94±18.38	87.20±17.13	1.725	0.085
TL1	63.99±10.23	64.51±10.02	-2.145	0.032

Statistics presented: mean±SD; n (%)

**Supplemental table 3. Comparisons of sectionalized RNFL thickness between different age groups in 16 sections.**

Segment	Age					P-value
	<39 (N=352)	40-49 (N=1728)	50-59 (N=2095)	60-69 (N=1647)	≥70 (N=550)	
<b>16 sections (μm)</b>						
TU1	74.67±12.05	74.78±12.47	73.69±12.76	72.45±12.28	69.77±14.77	< 0.001
TU2	101.68±18.17	102.31±18.09	100.24±17.83	98.02±17.89	94.28±20.32	< 0.001
ST2	139.45±23.36	141.50±23.57	137.85±22.82	134.29±24.11	128.69±26.39	< 0.001
ST1	158.05±23.60	160.89±23.52	156.92±23.81	151.39±24.08	146.64±26.30	< 0.001
SN1	140.65±27.98	142.51±25.93	138.92±25.97	135.21±26.35	130.78±25.98	< 0.001
SN2	134.88±25.08	136.46±22.73	133.47±23.05	130.58±23.52	126.50±22.63	< 0.001
NU2	104.53±20.70	103.57±19.56	102.75±19.98	101.46±21.41	98.37±21.25	< 0.001
NU1	74.70±14.52	74.63±14.27	74.24±15.01	73.12±15.73	71.12±15.94	< 0.001
NL1	68.85±13.28	69.45±12.97	69.15±13.54	68.54±14.32	67.33±16.14	0.022
NL2	91.84±18.74	93.08±18.71	91.66±18.68	90.16±19.69	87.85±21.49	< 0.001
IN2	129.32±22.10	130.10±22.66	128.69±22.98	127.72±23.92	124.21±25.23	< 0.001
IN1	153.52±25.45	153.95±27.33	152.64±26.79	151.36±27.62	146.24±28.94	< 0.001
IT1	171.16±23.51	172.12±22.89	168.57±22.88	164.70±23.59	157.46±27.31	< 0.001
IT2	144.21±25.77	144.44±25.36	140.71±24.71	134.60±25.05	128.85±27.83	< 0.001
TL2	90.42±18.29	89.99±17.32	88.52±17.87	85.30±17.23	82.10±20.25	< 0.001
TL1	65.44±9.80	65.11±9.64	64.59±10.21	63.59±9.81	61.44±12.45	< 0.001

Statistics presented: mean±SD; n (%)

Supplemental table 4. Comparisons of sectionalized RNFL thickness between male and female.

**Statistics presented: mean±SD; n (%)**

Segment	Female (N=3819)	Male (N=3205)	t	P-value
<b>RNFL thickness (µm)</b>				
Average	114.43±10.96	112.32±10.72	8.120	< 0.001
Superior	142.30±18.69	139.96±18.24	5.294	< 0.001
Nasal	84.87±15.59	83.39±14.90	4.046	< 0.001
Inferior	148.63±17.74	145.50±17.37	7.447	< 0.001
Temporal	81.94±12.77	80.43±12.23	5.057	< 0.001
<b>ONH parameters</b>				
Rim area (mm <sup>2</sup> )	1.73±0.49	1.63±0.46	8.621	< 0.001
Disc area (mm <sup>2</sup> )	2.41±0.51	2.44±0.48	-2.244	0.025
Average CDR (mm <sup>2</sup> )	0.33±0.13	0.34±0.14	-5.092	< 0.001
Cup volume (mm <sup>3</sup> )	0.12±0.15	0.17±0.20	-11.367	< 0.001
Axial length (mm)	22.67±0.92	23.05±0.88	-17.805	< 0.001

**Supplemental table 5. The supplementary analysis of VIF for the multivariate model**

Variables	VIF
Age	1.584
Gender	1.815
Smoking status	1.768
Body mass index	1.050
Diabetes	1.048
Coronary heart disease	1.055
Cataract extraction	1.019
Intraocular pressure	1.031
Spherical equivalent	1.213
Axial length	1.117
BCVA	1.603

**Supplemental table 6. Different types of OCT produced by different companies.**

OCT	Time to market	Type	Company	Country	Number of standard database eyes	Race	Proportion of Asians	Is it FDA certified
Stratus OCT	2002	TD-OCT	Carl Zeiss	Germany	284	Caucasian, Asian, African, Hispanic, Indian	24%	Yes
Cirrus OCT	2007	SD-OCT	Carl Zeiss	Germany	284	Caucasian, Asian, African, Hispanic, Indian	24%	Yes
Spectralis OCT	2006	SD-OCT	Heidelberg	Germany	201	Caucasian	0%	Yes
Topcon 3D OCT	2008	SD-OCT	Topcon	Japan	800	Japanese	100%	No
RTVue OCT	2006	SD-OCT	Optovue	America	861	Caucasian, Spanish, African, Chinese, Japanese, Indian	47%	Yes

Supplemental table 7. Post hoc test for comparisons of RNFL thickness parameters among the different age groups.

Comparisons	Average RNFL thickness		Superior RNFL thickness		Nasal RNFL thickness		Inferior RNFL thickness		temporal RNFL thickness	
	<i>t</i>	SE	<i>t</i>	<i>P-value</i>	<i>t</i>	<i>P-value</i>	<i>t</i>	<i>P-value</i>	<i>t</i>	<i>P-value</i>
≥ 70 - <39 = 0	-7.753	0.728	-8.107	<0.001	3.698	0.002	-8.590	<0.001	-7.145	<0.001
40-49 - <39 = 0	0.675	0.624	1.886	0.310	0.062	1.000	0.516	0.985	-0.037	1.000
50-59 - <39 = 0	-1.271	0.614	-1.411	0.605	-0.624	0.969	-1.874	0.317	-1.817	0.348
60-69 - <39 = 0	-3.754	0.626	-4.991	<0.001	-1.853	0.328	-4.834	<0.001	-4.408	<0.001
40-49 - ≥70 = 0	8.205	0.522	13.555	<0.001	5.230	<0.001	12.592	<0.001	9.918	<0.001
50-59 - ≥70 = 0	6.259	0.511	9.853	<0.001	4.518	<0.001	9.985	<0.001	7.995	<0.001
60-69 - ≥70 = 0	3.775	0.525	5.285	<0.001	2.917	0.027	6.141	<0.001	4.648	<0.001
50-59 - 40-49 = 0	-1.946	0.347	-5.894	<0.001	-1.217	0.729	-4.250	<0.001	-3.154	0.013
60-69 - 40-49 = 0	-4.430	0.367	-11.713	<0.001	-3.264	0.009	-9.120	<0.001	-7.453	<0.001
60-69 - 50-59 = 0	-2.484	0.351	-6.431	<0.001	-2.212	0.165	-5.343	<0.001	-4.681	<0.001

**Supplemental table 8. Post hoc test for comparisons of rim area RNFL thickness among age groups**

Comparisons	Rim area		Disc area		Average CDR		Cup volume		Axial length	
	<i>t</i>	<i>P-value</i>	<i>t</i>	<i>P-value</i>	<i>t</i>	<i>P-value</i>	<i>t</i>	<i>P-value</i>	<i>t</i>	<i>P-value</i>
≥ 70 - <39 = 0	0.423	0.993	3.396	0.006	-0.271	0.999	0.116	1.000	-3.997	0.001
40-49 - <39 = 0	-0.321	0.998	0.580	0.977	-0.545	0.981	1.569	0.501	0.600	0.973
50-59 - <39 = 0	-0.298	0.998	1.893	0.306	0.370	0.996	1.586	0.490	-2.156	0.185
60-69 - <39 = 0	-0.799	0.927	2.254	0.150	0.170	1.000	1.095	0.799	-3.615	0.003
40-49 - ≥70 = 0	-0.973	0.860	-4.041	< 0.001	-0.273	0.999	1.713	0.410	6.281	<0.001
50-59 - ≥70 = 0	-0.961	0.865	-2.562	0.072	0.830	0.916	1.742	0.392	3.106	0.015
60-69 - ≥70 = 0	-1.539	0.521	-2.019	0.243	0.578	0.977	1.145	0.771	1.229	0.722
50-59 - 40-49 = 0	0.048	1.000	2.312	0.132	1.635	0.458	-0.013	1.000	-4.917	<0.001
60-69 - 40-49 = 0	-0.818	0.921	2.858	0.032	1.215	0.730	-0.798	0.927	-7.175	<0.001
60-69 - 50-59 = 0	-0.903	0.890	0.708	0.952	-0.343	0.997	-0.822	0.919	-2.681	0.053



**Point-by-point responses to the reviewers' comments:**

*We would like to express our sincerest appreciation to the Editor and reviewers for their efforts on our article.*

Reviewer(s)' Comments to Author (if any):

Reviewer: 1

**Comments to the Author**

Wu et al. investigated normative profile of RNFL thickness measured by SD-OCT in Chinese population. Compared to previous studies, the number of subjects in this study is large. Also, considering the lack of studies in Asians on this topic, the current analysis is worthy of investigation. However, conclusion made in the Abstract of the paper (i.e., RNFL in the Chinese population was thicker than that in other studies) cannot be substantiated by the data. In addition, some corrections are needed to consider publication.

**Major comments:**

1. Presentation of the results for multivariate linear regression should be consistent throughout the manuscript.

- Abstract: result according to sex was repeated.

- Abstract and page 10: "decreased BMI" and "(BMI>30)" – These two are contradictory.

- In the abstract, thinner RNFL was significantly associated with lower SE and longer AL.

However, in the Results section (page 10, line 20), thinner RNFL was significantly associated with decreased AL.

*Response: We would like to thank the reviewer for the positive comments and thorough evaluation of our manuscript, we apologized for the repetitive and unclear expression leading to misunderstanding. We have modified the corresponding part. (Line 36-37, Line 210-212)*

2. Some expressions need to be clarified.

- Abstract conclusion, "RNFL thickness in Chinese population is thicker compared to other studies": Since this study did not conduct a qualitative systematic review on this topic, it would be better to present conclusions on results directly obtained from the analysis.

*Response: We thank the reviewer for pointing out the flaw in this expression. We apologized for the improper statement due to our negligence. We have modified in the corresponding part. (Line 40-41)*

- Abstract conclusion, "Meanwhile, many ocular and systemic factors are closely related to the changes of RNFL": This study did not evaluate "changes" of RNFL.

*Response: Thanks for your suggestion, we apologized for the inappropriate description. We described the true value but not the changes, which were mostly adopted in the cohort study. Thanks for your professionalism and scrutiny. We also amended the original text accordingly. (Line 42)*

### 3. Considerations for statistics

- In the multivariate analyses, multicollinearity needs to be considered. This is because the factors included in the analysis (i.e., Diabetes and HbA1c or SE and AL) are likely to have a significant correlation with each other.

*Response: Thanks for your suggestion. We supplemented the Variance Inflation Factor (VIF) for each covariate in the multivariate model. The VIFs were all less than 10, indicating no obvious multicollinearity in the reported model. The VIFs are listed as follows (as there was no HbA1c parameter included in our multivariate model analysis, thus HbA1c is not shown in the following table). We also added the relevant description in the manuscript. (Line 154, Line 213, Supplemental table 5)*

<b>The analysis of VIF for the multivariate model</b>	
<b>Variables</b>	<b>VIF</b>
<b>Age</b>	1.584
<b>Gender</b>	1.815
<b>Smoking status</b>	1.768
<b>Body mass index</b>	1.050
<b>Diabetes</b>	1.048
<b>Coronary heart disease</b>	1.055
<b>Cataract extraction</b>	1.019
<b>Intraocular pressure</b>	1.031
<b>Spherical equivalent</b>	1.213
<b>Axial length</b>	1.117
<b>BCVA</b>	1.603

- Table 2 and Supplemental Tables: OCT sectors should be compared considering issue of multiple comparisons.

*Response: Thanks for your suggestion. We applied Bonferroni correction to control for the potential errors found in multiple comparisons.*

*The adjusted significant level was  $\alpha' = \alpha/n$ , in which  $\alpha = 0.05$ ,  $n = 10$ . Then the comparisons of OCT sectors in Table 2 were considered to be statistically significant only when the P-value was less than or equal to  $\alpha' = 0.005$ . After the adjustment of the significant level, our conclusion regarding comparisons among age groups in Table 2 remained unchanged. Besides, we did not consider Bonferroni correction for comparisons in*

*supplementary tables because these variables were not the primary effects. And results from comparisons in supplemental tables were considered as explanatory rather than confirmatory conclusions.*

- Table 2: post hoc test should be considered after ANOVA to reveal difference between each age group.

*Response: Thanks for your suggestion. We supplemented Turkey test as a post hoc test after ANOVA across age groups. The whole results are listed as follows, and we have added the integrated version as supplemental tables 7 and 8. Also, the accordingly description in the method and result part was added in the manuscript. (Line 152, 188)*

**Table. Post hoc test for comparisons of average RNFL thickness among age groups**

Comparisons	Estimate	SE	t	P-value
$\geq 70 - <39 = 0$	-7.753	0.728	-10.346	<0.001
40-49 - <39 = 0	0.675	0.624	1.083	0.805
50-59 - <39 = 0	-1.271	0.614	-2.069	0.221
60-69 - <39 = 0	-3.754	0.626	-5.996	<0.001
40-49 - $\geq 70 = 0$	8.205	0.522	15.718	<0.001
50-59 - $\geq 70 = 0$	6.259	0.511	12.252	<0.001
60-69 - $\geq 70 = 0$	3.775	0.525	7.190	<0.001
50-59 - 40-49 = 0	-1.946	0.347	-5.616	<0.001
60-69 - 40-49 = 0	-4.430	0.367	-12.064	<0.001
60-69 - 50-59 = 0	-2.484	0.351	-7.073	<0.001

**Table. Post hoc test for comparisons of superior RNFL thickness among age groups**

Comparisons	Estimate	SE	t	P-value
$\geq 70 - <39 = 0$	-10.061	1.241	-8.107	<0.001
40-49 - <39 = 0	2.005	1.063	1.886	0.310
50-59 - <39 = 0	-1.478	1.047	-1.411	0.605
60-69 - <39 = 0	-5.329	1.068	-4.991	<0.001
40-49 - $\geq 70 = 0$	12.066	0.890	13.555	<0.001
50-59 - $\geq 70 = 0$	8.583	0.871	9.853	<0.001
60-69 - $\geq 70 = 0$	4.732	0.895	5.285	<0.001
50-59 - 40-49 = 0	-3.483	0.591	-5.894	<0.001
60-69 - 40-49 = 0	-7.334	0.626	-11.713	<0.001
60-69 - 50-59 = 0	-3.851	0.599	-6.431	<0.001

**Table. Post hoc test for comparisons of nasal RNFL thickness among age groups**

Comparisons	Estimate	SE	t	P-value
$\geq 70 - <39 = 0$	-3.857	-1.043	3.698	0.002
40-49 - <39 = 0	0.055	0.893	0.062	1.000
50-59 - <39 = 0	-0.549	0.880	-0.624	0.969

<b>60-69 - &lt;39 = 0</b>	-1.662	0.897	-1.853	0.328
<b>40-49 - ≥70 = 0</b>	3.912	0.748	5.230	<0.001
<b>50-59 - ≥70 = 0</b>	3.307	0.732	4.518	<0.001
<b>60-69 - ≥70 = 0</b>	2.194	0.752	2.917	0.027
<b>50-59 - 40-49 = 0</b>	-0.604	0.496	-1.217	0.729
<b>60-69 - 40-49 = 0</b>	-1.717	0.526	-3.264	0.009
<b>60-69 - 50-59 = 0</b>	-1.113	0.503	-2.212	0.165

Table. Post hoc test for comparisons of inferior RNFL thickness among age groups

Comparisons	Estimate	SE	t	P-value
<b>≥70 - &lt;39 = 0</b>	-10.213	1.189	-8.590	<0.001
<b>40-49 - &lt;39 = 0</b>	0.526	1.019	0.516	0.985
<b>50-59 - &lt;39 = 0</b>	-1.880	1.003	-1.874	0.317
<b>60-69 - &lt;39 = 0</b>	-4.945	1.023	-4.834	<0.001
<b>40-49 - ≥70 = 0</b>	10.738	0.853	12.592	<0.001
<b>50-59 - ≥70 = 0</b>	8.333	0.835	9.985	<0.001
<b>60-69 - ≥70 = 0</b>	5.268	0.858	6.141	<0.001
<b>50-59 - 40-49 = 0</b>	-2.406	0.566	-4.250	<0.001
<b>60-69 - 40-49 = 0</b>	-5.470	0.600	-9.120	<0.001
<b>60-69 - 50-59 = 0</b>	-3.065	0.574	-5.343	<0.001

Table. Post hoc test for comparisons of temporal RNFL thickness among age groups

Comparisons	Estimate	SE	t	P-value
<b>≥70 - &lt;39 = 0</b>	-6.076	0.850	-7.145	<0.001
<b>40-49 - &lt;39 = 0</b>	-0.027	0.729	-0.037	1.000
<b>50-59 - &lt;39 = 0</b>	-1.304	0.718	-1.817	0.348
<b>60-69 - &lt;39 = 0</b>	-3.224	0.732	-4.408	<0.001
<b>40-49 - ≥70 = 0</b>	6.049	0.610	9.918	<0.001
<b>50-59 - ≥70 = 0</b>	4.772	0.597	7.995	<0.001
<b>60-69 - ≥70 = 0</b>	2.852	0.614	4.648	<0.001
<b>50-59 - 40-49 = 0</b>	-1.277	0.405	-3.154	0.013
<b>60-69 - 40-49 = 0</b>	-3.197	0.429	-7.453	<0.001
<b>60-69 - 50-59 = 0</b>	-1.921	0.410	-4.681	<0.001

Table. Post hoc test for comparisons of rim area RNFL thickness among age groups

Comparisons	Estimate	SE	t	P-value
<b>≥70 - &lt;39 = 0</b>	0.014	0.033	0.423	0.993
<b>40-49 - &lt;39 = 0</b>	-0.009	0.028	-0.321	0.998
<b>50-59 - &lt;39 = 0</b>	-0.008	0.028	-0.298	0.998
<b>60-69 - &lt;39 = 0</b>	-0.022	0.028	-0.799	0.927
<b>40-49 - ≥70 = 0</b>	-0.023	0.023	-0.973	0.860
<b>50-59 - ≥70 = 0</b>	-0.022	0.023	-0.961	0.865

<b>60-69 - <math>\geq 70 = 0</math></b>	-0.036	0.024	-1.539	0.521
<b>50-59 - 40-49 = 0</b>	0.001	0.016	0.048	1.000
<b>60-69 - 40-49 = 0</b>	-0.013	0.016	-0.818	0.921
<b>60-69 - 50-59 = 0</b>	-0.014	0.016	-0.903	0.890

Table. Post hoc test for comparisons of disc area RNFL thickness among age groups

Comparisons	Estimate	SE	t	P-value
<b><math>\geq 70 - &lt; 39 = 0</math></b>	0.114	0.034	3.396	0.006
<b>40-49 - <math>&lt; 39 = 0</math></b>	0.017	0.029	0.580	0.977
<b>50-59 - <math>&lt; 39 = 0</math></b>	0.054	0.028	1.893	0.306
<b>60-69 - <math>&lt; 39 = 0</math></b>	0.065	0.029	2.254	0.150
<b>40-49 - <math>\geq 70 = 0</math></b>	-0.098	0.024	-4.041	< 0.001
<b>50-59 - <math>\geq 70 = 0</math></b>	-0.061	0.024	-2.562	0.072
<b>60-69 - <math>\geq 70 = 0</math></b>	-0.049	0.024	-2.019	0.243
<b>50-59 - 40-49 = 0</b>	0.037	0.016	2.312	0.132
<b>60-69 - 40-49 = 0</b>	0.049	0.017	2.858	0.032
<b>60-69 - 50-59 = 0</b>	0.012	0.016	0.708	0.952

Table. Post hoc test for comparisons of average CDR RNFL thickness among age groups

Comparisons	Estimate	SE	t	P-value
<b><math>\geq 70 - &lt; 39 = 0</math></b>	-0.002	0.009	-0.271	0.999
<b>40-49 - <math>&lt; 39 = 0</math></b>	-0.004	0.008	-0.545	0.981
<b>50-59 - <math>&lt; 39 = 0</math></b>	0.003	0.008	0.370	0.996
<b>60-69 - <math>&lt; 39 = 0</math></b>	0.001	0.008	0.170	1.000
<b>40-49 - <math>\geq 70 = 0</math></b>	-0.002	0.007	-0.273	0.999
<b>50-59 - <math>\geq 70 = 0</math></b>	0.005	0.006	0.830	0.916
<b>60-69 - <math>\geq 70 = 0</math></b>	0.004	0.007	0.578	0.977
<b>50-59 - 40-49 = 0</b>	0.007	0.004	1.635	0.458
<b>60-69 - 40-49 = 0</b>	0.006	0.005	1.215	0.730
<b>60-69 - 50-59 = 0</b>	-0.002	0.004	-0.343	0.997

Table. Post hoc test for comparisons of cup volume RNFL thickness among age groups

Comparisons	Estimate	SE	t	P-value
<b><math>\geq 70 - &lt; 39 = 0</math></b>	0.001	0.012	0.116	1.000
<b>40-49 - <math>&lt; 39 = 0</math></b>	0.016	0.010	1.569	0.501
<b>50-59 - <math>&lt; 39 = 0</math></b>	0.016	0.010	1.586	0.490
<b>60-69 - <math>&lt; 39 = 0</math></b>	0.011	0.010	1.095	0.799
<b>40-49 - <math>\geq 70 = 0</math></b>	0.015	0.009	1.713	0.410
<b>50-59 - <math>\geq 70 = 0</math></b>	0.015	0.008	1.742	0.392
<b>60-69 - <math>\geq 70 = 0</math></b>	0.010	0.009	1.145	0.771
<b>50-59 - 40-49 = 0</b>	<0.001	0.006	-0.013	1.000
<b>60-69 - 40-49 = 0</b>	-0.005	0.006	-0.798	0.927

<b>60-69 - 50-59 = 0</b>	-0.005	0.006	-0.822	0.919
--------------------------	--------	-------	--------	-------

**Table. Post hoc test for comparisons of axial length RNFL thickness among age groups**

Comparisons	Estimate	SE	t	P-value
<b>≥70 - &lt;39 = 0</b>	-0.256	0.064	-3.997	0.001
<b>40-49 - &lt;39 = 0</b>	0.033	0.055	0.600	0.973
<b>50-59 - &lt;39 = 0</b>	-0.116	0.054	-2.156	0.185
<b>60-69 - &lt;39 = 0</b>	-0.199	0.055	-3.615	0.003
<b>40-49 - ≥70 = 0</b>	0.288	0.046	6.281	<0.001
<b>50-59 - ≥70 = 0</b>	0.140	0.045	3.106	0.015
<b>60-69 - ≥70 = 0</b>	0.057	0.046	1.229	0.722
<b>50-59 - 40-49 = 0</b>	-0.149	0.030	-4.917	<0.001
<b>60-69 - 40-49 = 0</b>	-0.232	0.032	-7.175	<0.001
<b>60-69 - 50-59 = 0</b>	-0.083	0.031	-2.681	0.053

4. Discussion Page 11 line 19-55: The authors claimed that the type of SD-OCT device and age distribution might be the main reasons for RNFL thickness difference. However, in this study, it is presumed that not only these two factors but also axial length had a significant effect on RNFL thickness measurement. It should be further discussed in the manuscript.

*Response: Thanks for your suggestion. The age and type of SD-OCT were not only the primary factors affecting RNFL. There were a lot of factors affecting RNFL thickness, which were further discussed and analyzed in this study, we revised and clearly stated in the text (the axial length was discussed on Line 317-324).*

5. The average AL was about 22.8 mm, which is much shorter compared with previous study on Chinese population. This needs to be explained.

Ex) Ho et al. Ophthalmology 2019 (reference #53)

- Chinese (n=1371 participants)
- Average RNFL thickness: 95.7
- Average age: 54.7
- Average axial length: 24.0

*Response: Thanks for your advice on this study. Most of the subjects in the study by Ho et al. were urban population, while the participants in this study were rural population. Relatively speaking, the myopia proportion induced by the education level and social factors of the rural population were lower than that of the urban population, so the axial length was relatively short, while axial length was also an important influence factor for the evaluation of RNFL thickness, so the relatively short axial length in this study can result in a thicker RNFL as a whole. This factor should be taken into account. Thank you for finding the shortcomings of this study, and we have added the explanation of this part in an appropriate section of Limitation (Lines 352-357).*

6. Page 10 line 40: “We demonstrated that.... Using SD-OCT”: this sentence cannot be substantiated by the data presented in the Methods/Results sections. I believe that this conclusion cannot be drawn from a simple comparison with the numbers presented in previous studies.

*Response: Thanks for your careful observation and suggestion on the rigor of the manuscript. This study did not make a statistical comparison between Asian and European populations in RNFL thickness, so the conclusion drawn was inaccurate. We just got the information from the observation of statistical results, so we modified the expression of conclusion. It was inaccurate to use the European Standard database when applying OCT to measure Asian population (Line 222-225).*

Minor comments:

1. 62 references are indicated in the Supplement file, but only 35 are indicated in the reference lists in the manuscript.

*Response: Thank you for your question. Given the submission requirements of the journal, our study controlled references of the whole article within 35, so the references in other supplementary documents were listed only in the supplementary references. We put all the references together and relist. If the editor has any special requirements, we will further modify as required by the editor (see References).*

2. Abbreviations and acronyms should be consistently used after their first appearance. Ex) AL for axial length.

*Response: Thank you for your suggestion on the shortcomings of this study. We carefully checked the use of abbreviations and acronyms in the whole paper and made a targeted modification in the manuscript.*

3. Page 8 line 31: The authors mentioned that the RNFL thickness satisfies the “ISNT rule”, which is generally used to describe the relative thickness of the neuroretinal rim.

*Response: We thank the reviewer for pointing out this issue. This is where we neglect and state improperly. We have amended the original text accordingly.*

4. Page 15 line 36: It would be better to separate the limitation part as an independent paragraph.

*Response: Thanks for your suggestion. The limitation was an important part of the discussion. We highly appreciate your suggestion and set out Limitation as an individual paragraph (Line 352).*

Reviewer: 2

#### Comments to the Author

This study investigated the normative profile of retinal nerve fiber layer (RNFL) thickness based on spectral-domain optical coherence tomography and its associations with related parameters in a Chinese population.

Overall, this study supports the clinically important finding that the Chinese have thicker peripapillary RNFL from a population study, although this is not a novel one.

#### Specific comments:

1. 4th sentence, Abstract needs to be rephrased:

- 1) delete either male or male sex,
- 2) change 'less diabetes' to 'absence of diabetes',
- 3) change 'more history of cataract surgery' to 'history of cataract surgery'.

*Response: Thanks for the meticulous concern, we apologized for the repetitive and unclear expression leading to misunderstanding. The relevant mistakes in spelling and punctuation have been addressed. (Line 36-38, Line 210-212)*

2. Proof-reading and editing by a professional native English speaker is needed. Grammar errors are noted throughout the manuscript: e.g. Line 15-28, Introduction: With the developing of SD-OCT gradually, automated segmentation and measurement of individual retinal layers is ...

*Response: We thank the reviewer for the comment, we have amended the relevant sentence in the manuscript. (Line 75-78) And accordingly, we already went through the manuscript and corrected and polished the language mistakes in the revised manuscript. The revised manuscript has been edited and proofread by a native speaker.*

3. A separate paragraph needs to be added for better discussion of the association between cataract surgery and RNFL thickness.

*Response: Thank you for your suggestions on this study. According to the suggestion, we queried the relationship between cataract surgery and RNFL thickness and found some valuable opinions also hypotheses. The contents were listed as an individual paragraph and discussed accordingly. (Lines 333-343).*

4. Some OCT machines are known to include Asian eyes in their normative database. Please provide a table or a paragraph comparing the percentages of Asians in the normative database in different recent OCT machines.

*Response: Thank you for your suggestion on this study. According to your suggestion, we*



*summarized the common OCTs that were commercially available, and searched for information in RNFL thickness standard database related to instruments from the official website and related literature of OCT and compiled them into a table (Supplementary Table 6). As can be seen from these results, the RNFL thickness standard database of OCT instruments produced by different companies varied greatly from population to population. Hence, it is necessary to list and provide clinical reference for the measurement of RNFL thickness and the diagnosis of relevant diseases (Line 251-261).*

Editor(s)' Comments to Author (if any):

Section Editor

Comments to the Author:

(There are no comments.)

*We thank the reviewer for pointing out this issue. This is really where we neglect and state improperly.*