Supplementary Materials

Determinants of non-attendance at face-to-face and telemedicine ophthalmic consultations

```
SK Wagner MD<sup>1, 2, 3</sup>, LD Raja MD<sup>1</sup>, M Cortina Borja PhD <sup>4</sup>, J Huemer MD<sup>1</sup>, RR Struyven MD<sup>3</sup>, <sup>5</sup>, PA Keane MD<sup>1, 2, 3</sup>, K Balaskas MD<sup>1, 2, 3</sup>, DA Sim PhD<sup>1, 2</sup>, PBM Thomas PhD<sup>1, 2</sup>, JS Rahi PhD<sup>2</sup>, <sup>4, 6, 7</sup>, AL Solebo PhD<sup>2, 4, 6, 7</sup>, S Kang MD<sup>1, 2, 3*</sup>
```

*Correspondence and reprint requests: Swan Kang MBBS MBioch FRCOphth, Moorfields Eye Hospital NHS Foundation Trust, 162 City Road, London, United Kingdom. Tel: +44 207 253 3411 Email: swan.kang1@nhs.net

¹Moorfields Eye Hospital NHS Foundation Trust, London, UK

²Institute of Ophthalmology, University College London, London, UK

³NIHR Biomedical Research Centre at Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology, London, UK

 $^{^4\}mathrm{Great}$ Ormond Street Institute of Child Health, University College London, London, UK

⁵Centre for Medical Image Computing, Department of Medical Physics and Biomedical Engineering, University College London, London, UK


⁶Great Ormond Street Hospital NHS Foundation Trust, London, UK

⁷Ulverscroft Vision Research Group, University College London, London, UK


Methods

Index of Multiple Deprivation (IMD) was derived from the patient's residential postcode and cross-referenced to national data tables for 2019 provided by the UK Ministry of Housing, Communities & Local Government[1]. The Digital Exclusion Risk Index (DERI) covers twelve indicators across three equally-weighted domains - deprivation (33%), age (33%) and home broadband access and speed (33%) (full data available here: https://github.com/GreaterManchesterODA/Digital-Exclusion-Risk-Index)[2]. We suspected and statistically confirmed collinearity between the DERI and IMD (Pearson r = 0.68) and therefore opted to solely use the broadband component of the DERI. This comprises three indicators: i) the proportion of homes unable to receive a 30MBit/s connection, ii) the proportion of connections receiving less than 10MBit/s broadband and iii) the average download speed. The broadband component of the DERI did not correlate with IMD (r=0.04).

All analyses were conducted in R version 4.1.0 (R Core Team, 2021. R Foundation for Statistical Computing, Vienna, Austria). Multiple imputation was conducted using the mice package and visualisations were generated using the ggplot and jtools packages. Reporting is in line with the guidelines set by the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) and its extension, the REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) statements (Checklist: Supplementary materials) [3,4].



Supplementary Figure 1: Distribution of first appointments of newly registered patients by attendance status across the study period.



Supplementary Figure 2: Flow diagram illustrating the reasons for exclusion/ineligibility and division into mode of healthcare delivery. Pre-pandemic is defined as the period between January 1st 2019 and March 23rd 2021 inclusive, the date of the first UK lockdown. Pandemic is defined as March 24th 2021 to October 31st 2021 inclusive.

Supplemental material

Characteristic		F2F Pre-Pandemic	F2F Pandemic	Asynchronous	Synchronous – Telephone	Synchronous – Audiovisual
	F2F Pre-Pandemic					
	F2F Pandemic	1				
Sex	Asynchronous	<0.001	0.0124			
	Synchronous – Telephone	0.51	0.20	0.0030		
	Synchronous – Webcamera	<0.001	<0.001	<0.001	1	
	F2F Pre-Pandemic					
	F2F Pandemic	<0.001				
Age	Asynchronous	<0.001	<0.001			
	Synchronous – Telephone	<0.001	<0.001	<0.001		
	Synchronous – Webcamera	<0.001	<0.001	<0.001	<0.001	
	F2F Pre-Pandemic					
	F2F Pandemic	<0.001				
Asian (South) ethnicity	Asynchronous	<0.001	<0.001			
	Synchronous – Telephone	0.0360	1	1		
	Synchronous –Webcamera	<0.001	0.0083	1	1	
	F2F Pre-Pandemic					
	F2F Pandemic	<0.001				
Black ethnicity	Asynchronous	1	<0.001			
	Synchronous – Telephone	0.0014	1	0.0016		
	Synchronous – Webcamera	<0.001	0.08	<0.001	1	
	F2F Pre-Pandemic					
	F2F Pandemic	<0.001				
Other ethnicity	Asynchronous	<0.001	<0.001			
	Synchronous – Telephone	<0.001	<0.001	1		
	Synchronous – Webcamera	<0.001	<0.001	<0.001	<0.001	
White otherisity	F2F Pre-Pandemic					
White ethnicity	F2F Pandemic	<0.001				

	Asynchronous	<0.001	1			
	Synchronous – Telephone	<0.001	<0.001	<0.001		
	Synchronous – Webcamera	<0.001	0.33	1	<0.001	
	F2F Pre-Pandemic					
	F2F Pandemic	<0.001				
Unknown ethnicity	Asynchronous	<0.001	<0.001			
	Synchronous – Telephone	<0.001	<0.001	<0.001		
	Synchronous – Webcamera	0.0019	<0.001	<0.001	<0.001	
	F2F Pre-Pandemic					
	F2F Pandemic	1				
SES	Asynchronous	0.0028	0.0271			
	Synchronous – Telephone	0.0013	0.0035	0.37		
	Synchronous – Webcamera	<0.001	<0.001	<0.001	0.15	
	F2F Pre-Pandemic					
Non-attendance	F2F Pandemic	<0.001				
	Asynchronous	<0.001	0.0051			
	Synchronous – Telephone	1	1	0.19		
	Synchronous – Webcamera	<0.001	<0.001	<0.001	0.0207	

Supplementary Table 2. Baseline characteristics of the cohort failing to self-report ethnicity.

	Characteristic U		Reported N=32,137	<i>p-</i> value	
Sex n (%)	Female	29,551 (54.9%)	17,185 (53.5%)	<0.001	
Age median (IQR)	Years	56 (14.5)	53 (15.5)	<0.001	
SES ² median (IQR)	Decile (1=most deprived)	5 (2)	5 (2)	0.016	
Broadband Access ³ median (IQR)	Index (1=most at risk of digital exclusion)	3.40 (0.2)	3.41(0.2)	<0.001	
Interpreter n (%)	Yes	451 (0.8%)	281 (0.9%)	0.61	
Diabetes n (%)**	Yes	2,992 (5.6%)	1,200 (3.7%)	<0.001	
	Early Morning	24,015 (44.6%)	14,548 (45.2%)		
Time of Appointment	Late Morning	4,749 (8.8%)	2,711 (8.4%)	<0.001	
n (%)	Early Afternoon	19,748 (36.7%)	11,275 (35.1%)		
	Late Afternoon	5,275 (9.8%)	3,603 (11.2%)		
Sight-Impaired n (%)	Yes	318 (0.6%)	172 (0.5%)	0.31	
	No	44,067 (81.9%)	28,195 (87.7%)		
Previous cancellation n (%)	By hospital	7,037 (13.1%)	2,743 (8.5%)	<0.001	
	By patient	2,683 (5.0%)	1,199 (3.7%)		
	Adnexal	5,484 (10.2%)	1,538 (4.8%)		
	Cataract	10,034 (18.7%)	5,085 (15.8%)		
Specialty n (%)	General	18,559 (34.5%)	16,654 (51.8%)	<0.001	
	Glaucoma	8,672 (16.1%)	3,304 (10.3%)		
	Medical Retina	11,038 (20.5%)	5,556 (17.2%)		
Attendance status n (%)	Non-attendance	5,948 (11.1%)	2,340 (7.3%)	<0.001	

Supplementary Table 3: Adjusted odds ratios for non-attendance derived from multivariable logistic regression stratified by delivery mode with multiple imputation (predominantly ethnicity, but also SES and Broadband)

		Pre-pandem	ic F2F	Pandemic F2F Asynchronous		Synchronous (telephone)		Synchronous (audiovisual)			
		OR (95% CI)	<i>p</i> -value	OR (95% CI)	<i>p</i> -value	OR (95% CI)	<i>p</i> -value	OR (95% CI)	<i>p</i> -value	OR (95% CI)	<i>p</i> -value
Age	(decade)	0.92 (0.90-0.94)	<0.001	0.90 (0.88-0.92)	<0.001	0.90 (0.86-0.95)	<0.001	0.83 (0.73-0.95)	0.0063	1.05 (0.98-1.13)	0.14
Sex	Female	Reference		Reference		Reference		Reference		Reference	
	Male	1.17 (1.09-1.25)	<0.001	1.11 (1.02-1.20)	0.0119	1.27 (1.11-1.45)	<0.001	1.26 (0.88-1.82)	0.20	1.31 (1.05-1.63)	0.0161
Ethnicity	White	Referen	ce	Referen	ce	Referenc	e	Referenc	e	Reference 2.43 (1.32-4.48)	,
	Asian (South)	1.05 (0.91-1.21)	0.49	1.14 (0.91-1.43)	0.23	1.44 (1.01-2.04)	0.0416	1.49 (0.48-4.57)	0.47	2.43 (1.32-4.48)	0.0052
	Black	1.28 (1.08-1.51)	0.0058	1.69 (1.30-2.20)	<0.001	1.02 (0.76-1.39)	0.88	2.58 (0.70-9.46)	0.14	4.62 (2.27-9.42)	<0.001
	Other	1.37 (1.23-1.53)	<0.001	2.04 (1.66-2.50)	<0.001	2.41 (1.90-3.06)	<0.001	2.67 (1.03-6.93)	0.0440	1.67 (1.00-2.79)	0.0050
SES (greater deprivation)	Per decile decrease	1.11 (1.10-1.13)	<0.001	1.08 (1.06-1.10)	<0.001	1.07 (1.04-1.11)	<0.001	1.03 (0.95-1.12)	0.47	1.16 (1.10-1.23)	<0.001
Broadband	Index	0.94 (0.86-1.01)	0.10	0.99 (0.90-1.09)	0.84	0.94 (0.80-1.11)	0.46	0.81 (0.52-1.26)	0.34	1.06 (0.84-1.34)	0.56
Interpreter	er No Reference		ce	Reference		Reference		Reference		Reference	
	Yes	0.52 (0.34-0.79)	0.0023	1.09 (0.72-1.67)	0.68	1.26 (0.62-2.54)	0.52	2.55 (0.75-8.73)	0.13	0.70 (0.10-5.25)	0.73
Diabetes mellitus	No	Reference		Reference		Reference		Reference		Reference	
	Yes	0.88 (0.75-1.04)	0.13	0.58 (0.48-0.70)	<0.001	0.53 (0.41-0.67)	<0.001	1.48 (0.31-7.08)	0.62	*	0.50

Subspecialty	Adnexal	Reference		Reference		*		Reference		Reference	
	Cataract	1.46 (1.24-1.72)	<0.001	1.00 (0.86-1.17)	0.98	*		1.20 (0.55-2.61)	0.65	1.16 (0.68-1.97)	0.59
	General	1.31 (1.14-1.52)	<0.001	0.45 (0.38-0.51)	<0.001	*		1.38 (0.62-3.08)	0.43	0.55 (0.44-0.71)	<0.001
	Glaucoma	1.64 (1.38-1.95)	<0.001	0.96 (0.80-1.15)	0.65	Referenc	e	*		*	
	MR	2.08 (1.74-2.37)	<0.001	1.40 (1.21-1.61)	<0.001	2.39 (2.06-2.77)	<0.001	*		*	
Registered sight- impaired	No	Reference									
	Yes	0.34 (0.18–0.64)	<0.001	0.50 (0.28-0.90)	0.0208	0.59 (0.14-2.57)	0.49	*	*	*	
Appointmen t time	Early afternoon	Reference									
	Early morning	1.09 (1.01-1.17)	0.0196	1.05 (0.97-1.15)	0.22	1.15 (0.97-1.36)	0.11	0.83 (0.51-1.35)	0.44	0.92 (0.71-1.18)	0.50
	Late morning	0.72 (0.61-0.85)	<0.001	0.97 (0.83-1.13)	0.68	1.03 (0.82-1.28)	0.82	0.59 (0.26-1.34)	0.21	0.74 (0.51-1.08)	0.12
	Late afternoon	0.63 (0.54-0.74)	<0.001	0.58 (0.47-0.73)	<0.001	1.18 (0.95-1.46)	0.13	1.66 (1.08-2.56)	0.0221	0.59 (0.41-0.83)	0.0028
Previous cancellation	No	Reference		Reference		Reference		Reference			
	By the hospital	1.32 (1.19-1.47)	<0.001	1.91 (1.72-2.12)	<0.001	2.08 (1.71-2.52)	<0.001	0.40 (0.21-0.75)	0.0042	2.19 (1.61-2.98)	<0.001
	By the patient	1.88 (1.63-2.15)	<0.001	1.84 (1.56-2.18)	<0.001	1.59 (1.26-2.01)	<0.001	1.37 (0.65-2.91)	0.41	1.63 (0.90-2.98)	0.11

^{*}Either no cases or very few leading to unstable estimates.

F2F: Face-to-face, OR: Odds ratio, SES: Socioeconomic status, MR: Medical retina.

References

- 1 Ministry of Housing, Communities, Local Government. English indices of deprivation 2019. 2019.https://www.gov.uk/government/statistics/english-indices-of-deprivation-2019 (accessed 24 Feb 2023).
- A new tool: the Digital Exclusion Risk Index. Good Things Foundation. 2021.https://www.goodthingsfoundation.org/a-new-tool-in-your-toolbox-the-digital-exclusion-risk-index/ (accessed 24 Feb 2023).
- Benchimol EI, Smeeth L, Guttmann A, *et al.* The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) Statement. *PLoS Med* 2015;**12**. doi:10.1371/journal.pmed.1001885
- 4 von Elm E, Altman DG, Egger M, *et al.* The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *PLoS Med* 2007;4. doi:10.1371/journal.pmed.0040296