- Exner, Sigm.—*Pflüger's Arch.*, Vol. I, p. 375, 1868.
 Fischer, F. P.—*Arch. f. Augenheilk.*, Vol. XCVI, p. 57, 1925.
 Garten, S.—*Graefse-Saemisch Handb. d. ges. Augenheilk.*, Bd. III, Kap. XII, Anhang., 2^{te} Aufl., 1925.
 Gescher, Jul.—*Arch. f. Augenheilk.*, Vol. XCVI, p. 419, 1925.
 Gurwitsch, A. and L.—*Arch. f. Entwicklungsmek. d. Organ.*, Vol. CVII, p. 829, 1926.
 Helmholtz, H. von —*Handb. d. physiol. Optik*, any edition.
 Hess, C. von.—*Ergeb. d. Physiol.*, Vol. XX, p. 1, 1922.
 Jago, J.—"Entoptics." Churchill, 1864.
 Joly, J.—*Phil. Mag.* 6 ser., Vol. XLI, p. 289, 1921.
 Klein, Fr.—*Arch. f. (Anat. u.) Physiol.*, p. 305, 1904; p. 140, 1905; p. 445, 1908; p. 531, 1910; Suppl. Bd., p. 294, 1910; pp. 191, 339, 1911.
 Kühne, W.—*Untersuch. a. d. physiol. Institut. Heidelb.*, Vols. I, II, 1877-82.
 Johnson, G. Lindsay.—*Phil. Trans.* 194B p. 1, 1901.
 Marshall, W.—*Jl. Pharmac. and exp. Therap.*, Vol. XXX, p. 361, 1927.
 Müller, G. E.—*Zeitschr. f. Psychol., Ergänzungsbd.*, Vol. XVII, 1930.
 Müller, H.—*Zeitschr. f. wissenschaft. Zool.*, Vol. VIII, p. 1, 1856; *Verhandl. d. phys-med. Ges. Würzb.*, Vol. V, p. 411, 1855.
 Nuel, J. P.—*Arch. de Biol.*, Vol. IV, p. 641, 1883.
 Policard, A., et Paillot, A.—*Compt. Rend.*, Vol. CLXXXI, p. 378, 1925.
 Purkyně, J. E.—*Beobacht. u. Versuch. z. Physiol. d. Sinne*, Bd. I, Prag., 1823; Bd. II, Berlin, 1825.
 Radley and Grant.—*Fluorescence Analysis in Ultra-violet Light.* 1933.
 Schanz, Fr.—*Zeitschr. f. Sinnesphysiol.*, Vol. LIV, p. 93, 1923.
 Schreerer, R.—*Klin. Monatsbl. f. Augenheilk.*, Vol. LXXIII, p. 67, 1924.
 Schultze, Max.—*Arch. f. mikros. Anat.*, Vol. II, 175, 1866.
 Smith, D. T.—*Johns Hopkins Hosp. Bull.*, Vol. XXXI, p. 239, 1920.
 Sorin, A. N.—*Arch. f. Entwicklungsmek. d. Organ.*, Vol. CVIII, p. 634, 1926.
 Tait.—*Proc. Roy. Soc. Edin.*, Vol. VII, p. 605, 1872.
 Thomsen, E.—*Skand. Arch. f. Physiol.*, Vol. XXXVII, p. 1, 1919.
 Vierordt, K.—*Arch. f. physiol. Heilk.*, Vol. XV, pp. 255, 567, 1856.
 Wheatstone, C.—*Brit. Assoc. Rep.*, p. 551, 1831-32.
 Zehender, W. von.—*Klin. Monatsbl. f. Augenheilk.*, Vol. XXXIII, pp. 73, 112, 293, 339, 379, 1895.
 Zoth, O.—*Ergeb. d. Physiol.*, Vol. XXII, p. 345, 1923.

ORBITAL TERATOMA

BY

LIEUT.-COL. E. W. O'G. KIRWAN, F.R.C.S.I., I.M.S.

PROFESSOR OF OPHTHALMOLOGY, MEDICAL COLLEGE,
CALCUTTA, INDIA

ORBITAL teratomata are very rare tumours, and in ophthalmic literature I am only able to find 12 cases which so far have been reported. They are congenital, grow very rapidly, the child as a rule dying in the first few weeks of life. They may also occur in the form of a cystic tumour. The tumour consists of the derivatives of two or three germinal layers.

The Marchand-Bonnet theory best explains the formation of these tumours. During the early development of the embryo, the blastomere severed from its connections may remain as a

FIG. 1.

A low power micro-photograph taken with the micro-planar lens. Magnification $\times 4$. A, cartilage; B, intestinal epithelium; C, bone; D, fat; E, connective tissue bands with some smooth muscle; F, blood vessel; G, neuroglial tissue; H, smooth muscle and connective tissue.

FIG. 2a.

resting germ in any part of the body and begins to grow later or develop at the same time as the normal organs. As blastomeres are still capable of producing a normal body, after the first segmentation, but are able to produce only a few parts on continued division, it can be understood that derivatives of all or of only some embryonic layers are present in these tumours according as it is near the first cleavage or distant from it. The earlier the segmentation occurs, the less frequently does the germ stay latent.

Mizuo divides orbital tumours into four groups according to their degree of development.

(1) A foetus or a teratoid foetus attached to the orbit by an umbilical cord (orbitopagus parasiticus)—Mizuo's case.

(2) Parts of the body of a second foetus hang from the orbit (Ahlfeld's case).

FIG. 2b.

Micro-photograph showing epithelial glandular structure resembling intestinal epithelium. In some of the cells, round or oval white globules are seen which appear to be secretion showing that the cells are actually functioning. (a) shows the whole area; (b) part of the same under a higher magnification.

(3) A shapeless mass growing from the orbit and found anatomically to be a teratoma (Broer and Weigert's, Courant's and von Hippel's cases).

(4) The congenital orbital tumour containing the products of two germinal layers (cysts, cartilage, etc., mixed tumour). Mixed tumours of the lacrymal gland come under this heading.

FIG. 3.

Micro-photograph showing the area of smooth muscle and connective tissue.

Teratoma sometimes contains parts of organs and portions of the body. von Hippel states that complete organs are never found in these tumours.

True intra-orbital teratomata occur in infants, consist of derivatives of all three germinal layers and are very uncommon. The case to be described presently is the second to be reported from

India—the first was described by Elliot and Ingraham in 1910. In chronological order the following cases have been recorded:—

Holmes in 1863; Broer and Weigert, 1876; Ahlfeld, 1880; Lawson, 1884; Courant, 1893; Ewetsky in 1904; von Hippel in 1906; Mizuo in 1908; Coulter and Coats, 1910; Elliot and Ingraham, 1910; Corbett, 1925; Kearney, 1926.

In Holmes' case the tumour consisted of a female twin and was seen seven weeks after birth. The eye was protruded and formed the apex of the tumour.

In Broer and Weigert's case, the baby was seen one day after birth when the tumour was found. The eyeball was on the apex of the tumour, the cornea was opaque and the tumour was in the right orbit and displaced the nose and cheek. The tumour was the size of an apple and was excised. The child died.

In Ahlfeld's case the tumour consisted of a separate foetus. The buttock and the left leg projected from the left orbit.

In Lawson's case, the baby was seen two days after birth. The eyeball was situated on the top of the tumour and the cornea was cloudy. The child died.

FIG. 4.

Micro-photograph showing a rather primitive form of cartilage which formed a large encapsulated mass in the tumour.

In Courant's case the baby was seen immediately after birth. The eyeball was completely protruded and situated on the top of the tumour about the size of an apple. The cornea was opaque. The tumour was removed and the child recovered.

FIG. 5.

A micro-photograph showing fatty tissue intersected with connective tissue bands containing blood vessels.

In Ewetsky's case, the child was seen four and a half months after birth. The tumour filled the right orbit and surrounded a rudimentary eye at its summit. The tumour was removed, but the child died.

In von Hippel's case the baby was seen five days after birth. The tumour was the size of a moderate apple, enclosed the eyeball and the optic nerve. The cornea was dull. The tumour was removed.

PLATE I.

Teratoma of the orbit. Condition at birth.

PLATE II.

A coloured drawing of the section showing fourteen different types of tissue. Magnification X8.

1. Cartilage.	6. Muscle (involuntary).	11. Liver tissue.
2. Nerve.	7. Post. root ganglion.	12. Squamous epithelium (skin).
3. Probably small intestine.	8. Intestinal wall c villi.	13. Vascular tissue c melanoblasts (choroid).
4. Lymph.	9. Heart muscle.	14. Parathyroid.
5. Fat.	10. Hair.	

In Mizuo's case, a Japanese child was seen immediately after birth. A red injected swelling of the conjunctiva was seen which grew quickly and projected beyond the eyelids. On the 50th day after birth, a small teratoid foetus and the opening in the conjunctiva grew larger and the whole foetus was expelled with rudimentary arms and legs. The child lived and some vision was retained in the left eye.

In Coulter and Coats' case, seen three days after birth, the left eye was pushed forwards outside the orbit with the eyelid stretched

FIG. 6.

A micro-photograph showing commencing bone formation from cartilage.

around its equator. The cornea was opaque and a soft elastic tumour could be felt in the orbit. The child lived 19 days and died of exhaustion.

In Elliot and Ingraham's case, a Hindu female child was seen six months after birth. A tumour about the size of a small lime was present at birth and was then the size of an orange. The left eye was not to be seen. The nose, cheek and mouth were pushed to the right side. The tumour was removed and the child lived.

In Kearney's case a female child was seen immediately after birth, the right eye was pushed forward by a red mass about the size and shape of a tomato of medium size. The cornea was the same as that of the other eye and was not opaque, both pupils were equal in size. The tumour was removed and the child lived.

The following case of intra-orbital teratoma recently came under my care:—

R. S., Bengali, female baby, first child, born at full term on August 9, 1932, quite normal. Both father and mother were in good health. The baby was brought to the hospital one week after birth, and the case was diagnosed as an intra-orbital new growth. The baby was up to normal size and weighed seven pounds. From the left orbital cavity a red mass about the size of an ordinary tomato protruded and in the middle of the mass

FIG. 7.

A micro-photograph showing a "wild riot" of fibrous tissue bundles passing in all directions.

the anterior half of the eye could be seen, the posterior half being buried in the growth. The cornea was irregularly opaque rendering the examination of the fundus impossible. The tension of the eyeball was normal. The size of the cornea appeared to be about the same as in the normal eye. The upper and lower eyelids were tremendously distended. The mass was growing principally on the inner side of the orbital cavity and was pressing upon the lateral side of the nose obliterating the left nasal cavity. The

FIG. 8.

growth was hard and elastic, and the consistence was the same all over.

An exenteration of the orbit was carried out on August 16, 1932, a week after birth. The tumour was found to be adherent to the eyeball, but was separated by blunt dissection. After removal the mass was incised and found to contain a cyst holding about half an ounce of serous-like fluid. The rest consisted of solid growth about the size of a large walnut. The operation was carried out very rapidly and the baby had an uneventful recovery. The case was last seen by me on April 13, 1934, and a picture of the child is shown in the accompanying figure (Fig. 8). The palpebral fissure and the eyelids are now the same as in the right eye. There is only a small conjunctival sac. The vertical and the horizontal meridia of the bony orbit are somewhat larger than in the right eye. In other respects the child is quite normal.

Pathological report.—Serial sections show all the different types of tissues to be found in the body which originate in the three

germinal layers, namely, cartilage, nerve, lymphoid tissue, fat, involuntary and heart muscle, skin, hair follicles, vascular tissue with melanoblasts, small intestine, liver, and parathyroid. The eyeball and the optic nerve are normal and are independent of the origin of the tumour.

Diagnosis.—True filial teratoma of the left orbit.

In this case certain cells of the developing embryo had taken on independent growth after being segregated in the left orbit and this must have taken place relatively late in intra-uterine life as the development of the eye and optic nerve had already taken place.

I am indebted to Lt.-Col. H. Shortt, I.M.S., Director of the Central Research Institute, India, for the micro-photographs, and to Capt. S. K. Roy, Surgeon to the Mayo Hospital, Calcutta, for bringing the case to me.

BIBLIOGRAPHY

- Mizuo.—*Ber. u. d. 35st Versamml. d. Ophth. Gesellschaft. z., Heidelberg*, p. 347, 1908. *Arch. f. Augenheilk.*, Vol. LXV.
 Ahlfeld.—*Die Missbildungen d. Menschen. I Abschnitt*, Leipzig, Taf. VI, Fig. 11, p. 52, 1880.
 Broer and Weigert.—*Virch. Arch.*, Vol. LXVII, p. 518, 1876.
 Courant.—*Centralbl. f. Gynäkologie*, Vol. XVII, p. 740, 1893.
 von Hippel.—*Arch. f. Ophthal.*, Vol. LXIII, p. 1, 1906.
 Elliot and Ingraham.—*Ophthalmoscope*, December, 1910.
 Holmes.—*Trans. Path. Soc. Lond.*, Vol. XIV, p. 248, 1863.
 Lawson.—*Trans. Path. Soc. Lond.*, Vol. XXXV, p. 379, 1884.
 Ewetsky.—*Moscow Ophthal. Soc.*, Wratsch, Vol. XXI, p. 690, 1904; *Nagel's Jahresbericht*, p. 303, 1904.
 Coulter and Coats.—*Roy. Lond. Ophthal. Hosp. Repts.*, January, 1910.
 Corbett.—*Boston Med. and Surg. J.*, Vol. CXCII, pp. 484-488, 1925.
 Kearney.—*Amer. J. of Ophthal.*, Vol. IX, p. 416, 1926.

POSTERIOR LENTICONUS (REPORT OF A CASE)*

BY

T. H. LUO, M.D.

CHINA

AMONG its congenital or developmental anomalies, the human crystalline lens may present a conical bulging either on its anterior or posterior surface. The anomaly is a rare ocular condition, and according to the literature, the posterior type, lenticonus posterior, is more common than the anterior variety, lenticonus anterior. Posterior lenticonus was first recognized and reported by Meyer¹ in 1888. Owing to mistakes in either diagnosis or classification

* From the Department of Ophthalmology of the Peiping Union Medical College, Peiping, China.