BOOK NOTICE

Diseases of the Eye. By SIR JOHN PARSONS, F.R.S., Tenth Edition, revised with the assistance of H. B. STALLARD. P. 726. London: J. and A. Churchill. 1942. Price 25/-.

Close on five-and-thirty years ago a new star appeared in the ophthalmological book firmament. This was the first edition of Sir John Parsons' admirable book. It has been the mainstay of most English-speaking students for more than a generation, and we welcome the appearance of a new and revised edition, which gives promise of a continued long life. The reviewer was working in Sir John Parsons' clinic at Moorfields when the first edition came out and well remembers the appreciation with which it was universally received. In the early years of this century there were a number of good hand-books on diseases of the eye and students had a wide range from which to choose. Many read Swanzy, while to others the works of Jessop and Nettleship (the second edition of the latter by Holmes Spicer, was then either going or gone out of print), appealed; students who could afford a more expensive type of book indulged in Berry or Henry Juler, while those whose interest in eyes was only concerned with passing their finals relied on the excellent "aids" by the young Jonathan Hutchinson. Fuchs, then in a black cover, had not reached the undergraduate mind. Sir John's handbook immediately achieved success and this has been maintained by succeeding editions. We cannot imagine a better text-book of its size in our language. This tenth edition has undergone careful revision throughout. The section on general and local anaesthesia has been largely re-written and a special section devoted to anaesthesia and nerve block has been added. Recent advances in operative technique have been incorporated and the outstanding advances in therapeutics such as the sulphonamides and vitamins are included. Even so recent a discovery as that of the peculiar type of retinal haemorrhages found in some cases of hyperemesis gravidarum is mentioned. As the author says in his preface, "every effort has been made to retain its character as a reliable introduction to the diseases of the eye for students, general practitioners and junior ophthalmic surgeons." In our mind there is no doubt that this is so.

CORRESPONDENCE

To the Editors of THE BRITISH JOURNAL OF OPHTHALMOLOGY.

DEAR SIRS,—Miss Rosa Ford is of the same opinion still. It is scarcely surprising; the aerial bastions of her theories would hardly crumble at the first whiff of dialectical grapeshot.

I am grateful for the maternal solicitude of her reply to my recent letter in your columns; but it consists of little more than a reiteration of her familiar beliefs, and the implication that if her article of your March number has failed to convince me, her report of February might succeed. This naive suggestion is surely one of those triumphs of hope over experience which I have ventured to criticise in her views.

I regret to see that no other opinions have as yet been expressed in this controversy; it seems as though Miss Ford, who is rapidly reducing pathology to a single disease, is at the same time reducing criticism to a single obscure provincial voice.

I am anxious to be quite fair in the objections I have raised against her theories; I have not sought to press unduly the fewness of her cases and the absence of controls; I admit that we should not necessarily wait until an instructive rarity has become a commonplace before we dare to draw fair inferences from it. There are some phenomena—like the trout in the milkjug—which do not need confirmation; inescapable conclusions flow naturally from them; but I cannot agree that the facts reported by Miss Ford fall into this category. It is not primarily of the quantity of the evidence of which I complain; it is of the quality. Obviously, however, the fewer the cases, the more critical must be the examination and the more cautious the deductions drawn from them.

In my previous letter I confined myself (with some misgivings) to discussion of the conclusions she has drawn; and I tried to show that they were neither fair nor justified. But if, as Miss Ford herself suggests, her data are reviewed too—particularly those of her case reported in February—it seems that this attitude may have been over generous.

In the first place, Miss Ford clearly uses the term 'retrobulbar neuritis' in what might be called a Pickwickian sense. Surely the term implies a very definite and familiar clinical condition, characterised chiefly by the presence of a central scotoma, and with a very strong tendency to spontaneous recovery, and commonly if not invariably a herald of disseminated sclerosis. It is not difficult to visualise an early focus of this disease occurring more or less centrally in the optic nerve, and producing just such an effect. Miss Ford, on the other hand, uses the term retrobulbar neuritis as synonymous with peripheral field defects, which is an entirely different matter. To employ the same term for this condition too—without explanation or qualification—is not only to confuse the issue, but also to a considerable extent to beg the question. It is noteworthy that in her later article (*Brit. J. Ophthalm.*, March, 1942), no visual fields are given or described.

Her cases, as far as one can judge, may have been examples of peripheral interstitial neuritis or perineuritis; a rare and obscure

condition, but one into which a diagnosis of disseminated sclerosis can hardly enter. It is possible to imagine a focal lesion in the optic nerve disturbing the central fibres, but it is hardly possible to imagine an annular or spiral focus interrupting the peripheral ones. Miss Ford's terminology is so lacking in precision that it permits her to argue both ways; but if she is to call her colleagues sharply to task for diagnosing disseminated sclerosis too easily, she must at least reason from cases where such a diagnosis is likely.

The question is too lengthy to discuss in detail here; but it has been admirably summed up in "An Introduction to Clinical Perimetry" by Traquair, and in "Textbook of Ophthalmology" by Duke-Elder. In the second place, I cannot regard the evidence of nasal sinusitis in her cases as convincing; certainly not in the second one, which I have already discussed in my previous letter. With regard to the earlier case, we are told that two well-known rhinologists agreed that the nose was remarkably normal and without the slightest suspicion of any underlying sinusitis; but this pathetic stand by the Old Guard is overpowered by the triumphant advance of Miss Ford's theory. I still think it is at least possible they were right. I write here tentatively, and subject to correction; I am not knowledgeable about the nose; its dark impenetrable caverns are as unlighted for me as those of Kubla Khan; I look upon rhinologists as bold and bloody men. Perhaps I am unduly prejudiced by the fact that I have only twice entrusted myself to their care, and in each instance was fortunate to escape with my life; yet I still look upon their work with respect, and with childlike awe upon the products of their intrepid burrowings; I think twice, however, before unduly encouraging them.

It has long been recognised that the close proximity of the optic nerve to the sphenoidal air cells would seem anatomically to render it vulnerable to extension of infection from sinus disease; but that same proximity is itself an alarming prospect for the nasal surgeons. It is rumoured that the very optic nerve itself has not always been immune from the attention of their instruments; surgical interference with the deeply placed sinuses is not an operation lightly to be undertaken, but only on the clearest indications. It is perhaps significant that Miss Ford's first case relapsed after operation on the sphenoidal sinuses, but recovered after antrostomy. She does not indicate how the sinusitis is supposed to have affected the nerve, but this is not a strong confirmation of the rather attractive theory of damage produced by the direct spread of infection.

In the third place, the recovery of the visual fields after sinus drainage is inconclusive. Retrobulbar neuritis—even allowing for Miss Ford's specialised use of the term—is a disease notoriously liable to recover, whatever remedies are employed. All theories about its origin founded upon recovery after treatment are open to

the fallacy of *post hoc ergo propter hoc*, and must be critically examined in that light. It has also been quite reasonably suggested that when there is perineuritis of the optic nerve, operation on the sphenoidal sinus (even if it is not diseased) may hasten recovery through the beneficial effects of the relief of local congestion and the action of neighbouring processes of repair. An heroic remedy, if this is so.

All that is left then of Miss Ford's theory, is that a very strange sort of retrobulbar neuritis, in a few cases, is alleged to have been associated with a very peculiar type of sphenoidal sinusitis, despite the opinion of the well-known rhinologists; but under such circumstances that it is very difficult to demonstrate any connection between the two conditions, if either or both of them really existed. The whole is made plausible only by Miss Ford's technique; a few cases, a few facts, a few disputed data; these knit together by a loose terminology make up a theory to explain away the difficulties of at least three diseases which have baffled the patient work of painstaking investigators.

I hope this will serve to make clear my contentions, and that they will not appear prejudiced or unreasonably sceptical; if this is progress, I still prefer to wait. It is clearly profitless to labour the question, as the points of view of Miss Ford and myself are so diametrically opposed that they cannot be reconciled; we hardly speak the same scientific language. It remains only for me to retire from the controversial field, and to leave to her that latest word which is the privilege of her sex. Strive as I may to keep an open mind, I must confess that I am still unconvinced as to the possibility of closed nasal sinusitis—undetectable, and invisible to the most skilful eye—lurking in wait, and dealing out destruction in due time. Many and various, I know, are the weapons in the armoury of death and disease; but I cannot yet accept among them the sphenoidal submarine.

I am, Yours truly,

ALLAN H. BRIGGS.

20, MINSTER YARD,
LINCOLN, June 11, 1942.

OBITUARY

MALCOLM LANGTON HEPBURN

OLD Moorfields students in many parts of the world will have learnt with great regret of the death of Malcolm Hepburn. But by those who had the double privilege of being his colleagues and of knowing him well, his loss will indeed be deeply felt.