month by month, have contributed papers, and if the abstract section has occasionally been a little thin, it is not to be wondered at. Nor can one continue indefinitely to produce annotations of value. So far we have not descended to Mr. Pott's famous expedient, when an article on Chinese metaphysics was needed, of looking up China and metaphysics in an encyclopaedia and pooling the results, nor have we ever followed the advice offered by a very experienced editor to take one of the abstracts and put it into better English; yet at times during the past three years we have sailed rather wide of the ophthalmological mark. And if on occasions a strain of levity or of whimsical attempt at humour has crept into our lucubrations it has been done with a purpose. In times of stress we like to think that anything that raises a smile may help to enable people to carry on in spite of difficulties. Such has been our intention and if it has met with any success we are only too thankful. When peace comes we shall hope to revert to more solid contributions to the ophthalmic effort.

ABSTRACTS

MISCELLANEOUS

- (1) **Riwchun, M. H. (Buffalo, New York).—Albino with senile cataracts.** *Amer. Jl. Ophthal.*, Vol. XXIV, p. 798, 1941.

(1) **Riwchun** reports the successful extraction of senile mature cataract from each eye of a male albino, aged 59. He comments that he has been unable to find any statistics about the incidence of senile cataract in albinos and quotes Duke-Elder's text-book in which it is stated "There is no evidence available which shows that the eyes of albinos in later life are more vulnerable or prone to disease than normally pigmented eyes; there are several records of the extractions of senile cataract with good results."

H. B. STALLARD.

- (2) **Parkhill, E. M. and Benedict, W. L. (Rochester, Minnesota).—Gliomas of the retina.** *Amer. Jl. Ophthal.*, Vol. XXIV, p. 1354, 1941.

(2) **Parkhill** and **Benedict** discuss the origin, classification, grade of malignancy and prognosis of glioma retinae based on a histopathological study of 35 eyes. They suggest that these neoplasms arise from the glial cells which prevail in the inner nuclear, the ganglion-cell and nerve fibre layers and are probably the only retinal cells which retain the power of reproduction after birth. They

comment that histologically the majority of glioma retinae appear to originate in the inner nuclear layer, a few in the ganglion cell and nerve fibre layer.

They discuss the formation of rosettes as attempts at partial differentiation to a form imitating more or less closely the primitive epithelium of the neural tube rather than the highly specialized rod and cone cells. For what it is worth they agree that rosettes are sometimes found in metastatic growths and in medullo-blastomas of the central nervous system and in neuro-blastomas of the sympathetic nervous system.

The authors describe pseudo-rosettes which are aggregations of compressed glioma cells arranged around a blood vessel and sometimes bordering a clear space.

They classify glioma retinae into three types based on cytology, the degree of undifferentiation of the cells and thus the grade of malignancy. These are (1) Retino-blastoma, highly undifferentiated; (2) Neuro-epithelioma, showing slight or partial differentiation as indicated by rosettes or pseudo-rosettes. (3) Astrocytoma, well differentiated neoplasms. Rare in the retina. The cells nearly resemble normal astrocytes. The authors comment on the serious prognosis of the retino-blastoma type but in a large proportion of these cases described by them the optic nerve and extra-ocular tissues had already been invaded.

H. B. STALLARD.

CORRESPONDENCE

ANOMALOUS PUPIL REACTIONS

To the Editors of THE BRITISH JOURNAL OF OPHTHALMOLOGY.

DEAR SIRS,—It is clear that Cases 1 and 2 reported by Dr. Edleston in his interesting paper entitled "Anomalous Pupil Reactions," *Brit. Jl. of Ophthal.*, November, 1942, were suffering from tabes, but since he does not tell us whether the reaction to near vision was present or not, some doubt may exist in the minds of those who have read his paper as to whether a diagnosis of tabetic pupil is justified.

For the purpose of this comment however, I assume that the reaction to near vision was present and that the anomalous reaction, *viz.*, dilatation instead of constriction to light, was tabetic in origin.

This paradoxical reaction has been described as occasionally occurring in tabetic pupils, and at rare intervals has been noticed by ophthalmologists and neurologists. As far as I know this