without loss of aqueous. It is a state of affairs that indicates lack of presence of mind before beginning the operation, for this is the time to make sure that the edge of the knife is directed upwards. We do not say that such an event might not enable a nimble witted person to display presence of mind when he has to explain to the patient and to his relatives and friends why it was that the operation has had to be postponed, but we are not up to offering any suitable suggestions. In ophthalmological practice it is said that very occasionally the wrong eye has been excised for such a condition as choroidal sarcoma. Here no amount of presence of mind can conceal the fact that a terrible blunder has been made. We always adopted the plan of making sure which eye was affected and marking a cross on the forehead over that eye before the patient was taken to the theatre, with instruction to the nursing staff that the mark was not to be washed off. Care in the preliminaries in every case is the best guarantee that we shall not have to exhibit this inborn quality in such cases, even if it means that by lack of use, presence of mind may grow rusty.

A good example of presence of mind was exhibited on one occasion in the final examination for medicine. In the viva voce the candidate was told to examine the chest of a stout, middle-aged man. He found that the patient was completely dull to percussion, back and front, on one side, and no breath sounds were audible, while the heart was not displaced. When the examiner asked for a diagnosis the candidate was unable to offer one and said he didn't know. The examiner said brusquely "Why, hasn't he an empyema?" The candidate replied that no patient with an empyema capable of producing the signs found had any business to be an ambulant case at the examination hall, which was perfectly true and quite unanswerable. This answer did not adversely affect the result of the examination.

ABSTRACTS

MISCELLANEOUS

- (1) **Loewenstein and Steel (Glasgow).—Retinal tuberous sclerosis (Bourneville's disease).** *Amer. Jl. Ophthalm.*, Vol. XXIV, p. 731, 1941.

(1) **Loewenstein and Steel** describe the case of a boy aged 3½ years who had one eye excised for "pseudo-glioma" and subsequently suffered from epileptic fits but no sebaceous adenomata were present. Ophthalmoscopic examination of the remaining eye

showed grey-green tumours, exudative retinitis, retinitis proliferans and angiomatic nodules. Later this picture changed. The excised eye showed multiple tumours composed of nerve fibres, neural cells producing closed and open cysts, a thickened membrana limitans interna, pyramidal tumour of the optic disc, a choroidal angioma and an angiomatic growth on the surface of the retina. The authors state that the condition is a primary ectodermal malformation and the disturbance of balance of ectoderm and mesoderm leads secondarily to a mesodermal tumour.

H. B. STALLARD.

- (2) **M'Glone, Joseph (Glasgow).**—A case of chloroma. *Glas. Med. J.*, August, 1942.

(2) **M'Glone** reports a case of chloroma in a boy of ten years, involving both orbits and the left maxilla, having probably originated in the sphenoid. The course was rapid, lasting 18 weeks from the onset of proptosis until severe haemorrhage from the naso-pharynx caused death.

The general post-mortem appearance of the tumour is well illustrated. Secondary deposits were found in the lateral sinuses, paravertebral tissues in the upper thorax, and in the heart; histologically these deposits were composed of round cell infiltrations and myelocytes. Green urine was present in the bladder but the colour faded on exposure. It re-appeared with the addition of sodium hydrosulphite and it is suggested that this may prove to be a useful test.

B. W. RYCROFT

- (3) **St. Clair Roberts, B. H. (Worcester).**—Remarks on some types of industrial cataract: *Med. Press and Circ.*, November 4, 1942.

(3) To a special number of the *Medical Press and Circular* on industrial accidents, **St. Clair Roberts** contributes a very good account of some industrial cataracts. Having briefly dealt with traumatism, especially that due to perforation by metallic particles and the variety met with in country districts in those who pleach hedges, he mentions some of the chemical agents responsible; but the major part of his paper deals with physical agents and specially the type of cataract met with in chain makers, of which he is a recognized authority. The process of chain making is mostly localized in the Black Country still. Its details are set out and there is a good description of the type of cataract which occurs in the youngish middle-aged. The chain maker accepts the idea that it is customary to lose the sight in middle age. This, with the nature of their work, makes it very difficult to get them to use any form of protection in the shape of goggles. The largest chains are

made mostly by electrical machinery and the workers here are not so liable to suffer. The medium and small linked chains are still made by hand and here the workers are very likely to be affected. The author states that he has kept careful watch on a few cases who have been able to return to their work after removal of the cataract, and has never found any damage to occur at the macula.

Those of our readers who are interested in chain making might read F. Brett Young's novel "Far Forest," in which the background of the first part of the story is laid in one of the smithy hearths.

R. R. J.

- (4) **Ballantyne, A. J. (Glasgow).—Recent work on vascular disease and its significance in medical ophthalmology.**
Glas. Med. Jl., July, 1942.

(4) In his William Mackenzie Memorial Address **Ballantyne** surveys recent work on vascular disease and its association with medical ophthalmology. He classifies the retinal vascular changes as follows:—

1. *Calibre Changes.*

An increase in diameter, although uncommon, is noticed in profound anaemias with a tendency to equality in the colour and calibre of the arteries and veins, whereas a general diffuse contraction is much more common and widely recognised in such conditions as advanced atrophy of the optic nerve.

2. *Spastic Contraction of the Vessels.*

The difficulty in distinguishing between a true spasm of the vessel wall and an organic contraction is pointed out. True spasm is to be found in eclampsia and Raynaud's disease. In 38 cases who complained of a temporary loss of vision, partial or complete, 19 of them appeared to be examples of transient angiospasm, either of retinal or cortical vessels. Moreover, vascular spasm may also be the cause of permanent loss of vision if it occurs in retinal vessels previously narrowed by sclerotic changes, and the estimation of prognosis is thereby rendered difficult. A case is quoted where symptoms suggested an arterial spasm in the left eye, in which vision failed to return; identical symptoms in the right eye completely recovered.

3. *Angiospastic Retinopathy.*

The significance of arterial and arteriolar spasm in hypertension is reviewed and the element of spasm is stressed. There is reason to believe that this commences first in vessels smaller than those normally visible to the ophthalmoscope, since gross spasm, as in

quinine poisoning, does not produce similar fundus lesions. However, it is suggested that the various fundus changes may be brought about by a difference in the circulating toxin.

4. *Sheathing of the Retinal Vessels.*

Sheathing of the retinal vessels shows two distinctions, namely, that of the intima, which causes irregular or patchy thickening, and that of the media, which is more uniform. Regressive changes of a hyaline or lipoid nature render the walls visible and produce the phenomenon of sheathing. Sheathing of the veins, on the other hand, may be due to active periphlebitis, as in Eales' disease, or to "cuffing" of the vessels as in leukaemia. It is shown that sheathing is due to many causes; the blood column disappears and is replaced by a white band, but it does not necessarily mean that the vessel is completely obliterated.

5. *Arterio-Venous Crossings.*

The precise cause of Marcus Gunn's sign is disputed. It is pointed out that compression of the vein alone is not responsible for the narrowing of the lumen for in true compression expansion of the vessel wall would be expected. Ballantyne believes that Gunn's phenomenon is due to veiling of the venous blood column as it recedes into the deeper layers of the retina and passes behind the artery.

6. *New Vessel Formations.*

These may be observed in four clinical forms:—

1. In hypertension with arteriosclerotic changes where intervals between principal retinal vessels are filled by a loose network of very fine venous branches which may, or may not, be traceable to their connections with the principal veins. This type is seen in its most complete form as a sequel of central retinal vein obstruction. Such venous collaterals go by the name of "rete mirabile." As a rule this network is intraretinal, but occasionally branches may perforate the internal limiting membrane of the retina and push forward into the vitreous cavity, giving type 2.

2. Although this formation may arise at practically any part of the fundus, the internal limiting membrane does, to some extent, offer a barrier to its forward growth, and probably for this reason most of these new vascular formations arise from the vessels on the disc where the limiting membrane is absent.

3. The third type is known as retinitis proliferans and is usually at a point where a retinal haemorrhage has broken through the internal limiting membrane. Organisation of the blood clot is responsible for the opaque white bands attached to the anterior surface of the retina, and there is a preponderance of mesoblastic tissue, much more solid and substantial than the mucoid framework

of the previous type, which is sometimes so tenuous that it scarcely causes any appreciable interference with vision. The explanation for this new vessel formation is said to be mechanical and purposive in the first instance, whilst the second type is extravagant and purposeless.

4. This type deals with new vessels closely related to the main trunks of the retinal veins. They are of large size and may be accompanied by formations of tufts, fans or networks of dilated capillaries, and practically always occur along with other gross pathological changes in the retinal veins. Excellent diagrams are added to amplify the description of these larger venous formations, and micro-photographs show their relationship to the layers of the retina. It is emphasised that these changes are more related to diabetes mellitus than to hypertension. Post-mortem appearances in a woman aged 61 years, suffering from diabetes, showed details of the rete mirabile in the retina and, in addition, a pre-retinal formation of new veins, which could be traced to a large retinal vein which had perforated the internal limiting membrane. It is probable that the network of vessels lay between two layers of a divided internal limiting membrane. The occurrence of the more unusual loops and networks of large-calibred vessels in ophthalmoscopic examination of so many diabetic cases convinced the author that he was dealing with a distinct clinical entity. He points out that hypertension is not essential, and that in many diabetes cases histological examinations have shown that the vessel changes in the retina fall more emphatically on the veins (phlebosclerosis). Also that abnormal fulness of the veins is one of the earliest retinal changes in some diabetics, and that the characteristic punctate haemorrhages are apparently derived from the post-capillary venous radicles; nor indeed, are these venous changes familiar in numerous cases of hypertension.

The pathogenesis of the venous plexus and calibre changes is not clear, but it is suggested that they may be due to a retardation of the blood stream in the central retinal vein. On the other hand, Foster Moore (1924) found only one diabetic in 61 cases of thrombosis of the central retinal vein, whereas in the author's group of 16 cases 15 were diabetic, and he holds the view that gross venous changes in the retinal vascular system justifies the diagnosis obtained in the majority of cases.

7. *Oculo-Cerebral Syndromes.*

These syndromes are represented by the von Hippel-Lindau disease, with one or more haemangiomas in the retina and associated with angiomas of the cerebellum and elsewhere. Angioma of the choroid, with associated lesions of the central nervous system, is known as the Sturge-Weber Syndrome.

Five cases of subarachnoid haemorrhage are described in detail, together with post-mortem findings. They ranged in age from 39 to 62 years. In the pathological details stress is laid on the distribution of haemorrhage in relation to the brain and the sheath of the optic nerves. Of these cases all had elevation of blood pressure, and in only one was the fundus free from haemorrhage. In four cases the source of the subarachnoid haemorrhage was discovered, and in three of these it was the rupture of an aneurysm of the circle of Willis. It was shown that haemorrhage could occur not only in the dural sheath, the subdural and subarachnoid space, but also in the optic nerve and pial sheath, in the sclera near the porus opticus, in the capsule of Tenon, among the orbital fat and muscles, in the fibrous sheaths surrounding the ciliary ganglion, in the pial covering of the chiasma, and in the chiasma itself. The old explanation that blood, effused at the base of the brain, passed along the optic nerve sheaths through the optic canal is considered to be not entirely satisfactory, since there is not necessarily continuity between these various extravasations of blood.

The author is of opinion that the intra-cranial and intra-orbital haemorrhages are independent, and so also are those haemorrhages within the optic nerve sheaths and in the retina. He submits the view that the precipitating cause of these haemorrhages is the sudden rise of intra-cranial pressure which occurs at the moment of rupture of the aneurysm acting directly on the venous sinuses and causing extensive stasis and resulting extravasations.

A short bibliography completes the address.

B. W. RYCROFT.

- (5) **King, A. and Walsh, F. B. (Baltimore).—Temporary amaurosis and hemianopsia due to epilepsy.** *Amer. Jl. Ophthal.*, Vol. XXV, p. 398, 1942.

(5) **King and Walsh** describe two cases in which visual defects persisted after an epileptic attack when other functions had returned fully. One was a female child, aged 22 months, in whom blindness of both eyes persisted for 10 days after an epileptic seizure. During this period she showed no blink reflex to objects held close to her eyes and there was no evidence of nystagmus on looking down at a revolving drum. These became apparent on the tenth day. The authors comment that Ashby and Stephenson (*Lancet*, Vol. I, pp. 1294 and 1616, 1903), recorded one case of persistent blindness after an epileptic attack.

The second case was a male negro, aged 45, who had a post-epileptic left homonymous hemianopia which lasted a month and then recovered. In the blind part of the field he had visual hallucinations. In the discussion of these cases the authors support the conjecture that in cases where the pupillary reflexes are intact and

there is homonymous hemianopia there is temporary exhaustion of the visual cortex associated with the epileptic discharge.

H. B. STALLARD.

- (6) **Dunnington, John H., and Wheeler, Maynard C. (New York).—Operative results in two hundred and eleven cases of convergent strabismus.** *Arch. of Ophthalm.*, July, 1942.

(6) This paper by **Dunnington** and **Wheeler**, although covering familiar ground, contains many points of interest. The present position of squint surgery is put rather neatly by them in the aphorism that "the mathematical millenium is not in sight," and the results they quote go to prove this. In analysing the cases it was found that in as many as 28 per cent. the strabismus was present from birth though the average age of onset was 1.9 years. Amblyopia in one eye was present in roughly 50 per cent. of cases, and in 31 where constant occlusion was efficiently carried out, vision of 20/50 or better was obtained in 70 per cent. With regard to refractive error it is remarkable that only 17 per cent. of cases showed over 4D. of hypermetropia and that 68 per cent. showed less than 3D. No relationship was found to exist between the amount of deviation and the refractive error. In 71 per cent. there was a deviation of 20° or more, the average being 23°.

Estimation of the post-operative result was difficult and the authors took as an arbitrary definition of a cure a residual deviation of not more than 10 prism dioptres for either distance or near vision. Using this, they found that one operation was sufficient to effect a cure in 43 per cent. of cases. The under corrections varied in severity, but in most of them approximated to 20 prism dioptres. It is notable that abnormal retinal correspondence was more frequent in the failures than in the cures. The figures give little guidance as to the best age at which to operate, except for the observation that the highest proportion of over-corrections (20 per cent.) was obtained in the age group 3-8 years. The results obtained in alternating squint were inferior to those in the unocular variety, the tendency being towards under-correction in the former and over-correction in the latter. Although the effect of the same operation varied from case to case, it was found that the average correction obtained by recession on one eye only was 7.3 prism dioptres, while that obtained by a 4mm. recession combined with a 10mm. resection was 40 prism dioptres. The authors rightly remark, however, that much depends on the technique employed at operation.

Post-operative diplopia was frequent as a transient complaint, but in only three instances did it show any tendency to be permanent and in only one of these was it distressing.

F. A. W-N.