The Glass Bottle—with an evil genie, the glaucoma fiend

Early in the writer's career he was asked to undertake the holiday work of a friend. Everything went off all right and our only difficulty was that among the drop bottles in the consulting room there was none labelled homatropine and cocaine. Everything else one could possibly want was there, and the home of the eserine bottle was on the top of the travelling clock on the mantelpiece. Not finding any solution of "H. and C.," we searched in various receptacles and at last found a supply of tabloid ophthalmic drugs. These we made use of, and when our colleague returned we mentioned casually that we had'n't been able to find the "H. and C." bottle. "Why, that's it," said our friend, pointing to the top of the clock. We met this statement by showing that the bottle was labelled eserine. It then appeared that our friend had broken his "H. and C." bottle earlier in the year and had used the other ever since. He had forgotten to mention the fact when we took over. At the time this amused us, but had anything untoward happened the results might have been serious. We learnt a lesson we never forgot. Always have your bottles correctly labelled.

ABSTRACTS

I.—MISCELLANEOUS

- (1) **Allen, J. H. (Iowa).—Inclusion blennorrhoea.** *Amer. Jl. of Ophthalm.*, p. 833, August, 1944.

(1) Forty-three cases of inclusion blennorrhoea were observed during the last six years in the course of 9,580 deliveries, thus giving an incidence of 0·37 per cent. in the new born, or an incidence of approximately 10 per cent. of cases of ophthalmia neonatorum.

Symptoms appeared simultaneously in both eyes in 30 cases but in 12 cases one eye was involved first and in one case the infection remained confined to one eye. Three of the infants had secondary bacterial infection; treatment with 0·5 per cent. silver nitrate ointment three times a day eliminated the bacteria but had no effect on the course of the inclusion blennorrhoea.

The incubation period measured from time of birth to the first appearance of symptoms varied between 5 and 15 days with an average of 8 days.

The first sign of the disease was slight purulent discharge at the inner canthus of one or both eyes followed by oedema of the lids, subconjunctival infiltration, papillary hypertrophy and preauricular

adenopathy. Eversion of the lower lid exposed a longitudinally corrugated lower fornix. Then the purulent discharge increased with chemosis before the chronic stage developed, lasting on an average 60 days. Follicles when present are generally located in the lower lid and fornix. The total duration of the disease averaged 7 months. In several of the acute cases application of fluorescein showed a few minute staining areas.

Scrapings of the lower fornical conjunctiva stained with Giemsa showed typical inclusion bodies. The glass slides must be clean; the smears were fixed over night in absolute methyl alcohol and stained for one hour in dilute Giemsa at 37°C. Inclusion bodies were demonstrable on the average for 66 days after onset of the disease.

Occasionally differentiation between inclusion blennorrhoea and trachoma may be necessary. In trachoma, manifestations are most intense in the upper palpebral conjunctiva while infiltrates and neovascularization of normally clear corneal tissue (elementary pannus) appear early. In inclusion blennorrhoea the maximum pathological change is in the lower palpebral conjunctiva while corneal infiltrates and pannus do not occur. The apparent severity of the disease in its acute phase is out of all proportion to its good prognosis.

Cervical scrapings, made from mothers of 29 infants in this series of 43 cases, revealed inclusion bodies in 27.

The transfer of conjunctival scrapings from a case of inclusion blennorrhoea of the new-born to the normal conjunctiva of a six-year-old subject resulted in the development of a condition identical with that described for swimming-bath conjunctivitis or inclusion conjunctivitis of adults; no corneal changes such as occur in trachoma were seen. Such inoculation was carried out in 40 children in all from one eye to another by passaging; in none of the cases did the disease change its character. The disease was proved to be non-bacterial. Experimental proof of the ineffectivity of silver salts in the prevention of inclusion blennorrhoea was obtained.

The earlier work of Thygeson on inclusion blennorrhoea is referred to (*Trans. Amer. Ophthal. Soc.*, Vol. XXXIV, p. 340, 1936).

A. F. MACCALLAN.

- (2) **Somerville-Large, L. B. (Dublin).—Dublin's eye hospitals.** *Irish Jl. of Med. Sci.*, September-October, 1944.

(2) **Somerville-Large's** article on the ophthalmic hospitals of Dublin, is of great interest, and fills, what has been till now, a gap in the history of British ophthalmology. In the last century there were seven of these special hospitals in Dublin, "though only one

remained exclusively devoted to diseases of the eye during the whole of its existence." All but one of these hospitals were founded during the first half of the century. They made queer alliances: one was joined with diseases of women and children, one to maternity work, one to skins and a fourth to general medicine and surgery. The alliance between eye and ear has proved durable and still is in being today. The eye hospitals were as follows: The National Eye Hospital (1814-1904); the Charitable Institution (1817-1823); St. Mary's Hospital and Dublin Eye Infirmary (1819-1831); The Ophthalmic Hospital (1829-1834); The Coombe Lying-In Hospital and Dublin Ophthalmic Dispensary (1836-1847); St. Mark's Ophthalmic Hospital for Diseases of the Eye and Ear (1841-1904); Dublin Infirmary for Diseases of the Eye and Ear (1872-1875). At the end of the century plans for amalgamating the National Eye and Ear Infirmary and St. Mark's Ophthalmic Hospital were set afoot and resulted in the formation of the Royal Victoria Eye and Ear Hospital.

Many famous names in Irish ophthalmology and indeed of Irish medicine will be found in this careful history of part of the last century, from Isaac Ryall, Arthur Jacob, Francis White, Hugh Carmichael, Sir William Wilde, A. H. Jacob, Sir Henry Swanzy, J. B. Story, and L. Werner, to that great triumvirate, Graves, Stokes and Crampton.

The preparation of this article must have entailed a great deal of work, for many of the institutions moved their sites frequently, and the author describes those buildings which are still in being. We hope the paper will be widely read; it cannot fail to interest all English speaking ophthalmologists, and we offer hearty congratulations to the author on an admirable piece of work.

R: R. J.

II.—THERAPEUTICS

- (1) **Knight, H. C., Emory, M. and Callahan, N. (New Orleans).—Hyperpyrexia in the treatment of acute ocular inflammations.** *Amer. Jl. Ophthal.*, Vol. XXVII, p. 381, 1944.

(1) **Knight, Emory and Callahan** discuss the value of hyperpyrexia, as induced by the hypotherm cabinet, in certain inflammatory disorders of the eye such as keratitis, corneal ulceration, herpes zoster, uveitis, retinitis, optic neuritis, panophthalmitis and sympathetic ophthalmitis. They considered the lower level of therapeutic fever as 104° F. Almost all patients were able to endure 5 hours at 105° F. Gonococcal infections required 106-107° F.

Treatments were given at intervals of three to five days. A group of 52 cases were treated. The results are described as excellent in 25 cases and satisfactory in 12. There was immediate relief of pain. Visual restoration was delayed.

H. B. STALLARD.

- (2) **Hollander, L. (Pittsburgh) and Krugh, P. T. (East Point, Georgia).—Cancer of the eyelid.** *Amer. Jl. Ophthal.*, Vol. XXVII, p. 244, 1944.

(2) **Hollander and Krugh** found that carcinoma of the eyelids occurs in just over 9 per cent. of skin cancer cases. They report the results of treatment in 125 cases, 17 of these in the upper lid, 47 in the lower, 52 at the inner canthus, and 9 at the outer canthus. Sixty were examined histologically, 45 of these were basal-celled carcinoma, 13 squamous-celled, 1 mixed-celled and 1 a melanocarcinoma. Seventy-nine were men and 46 women and their ages ranged from 25 to 80 years, 82 of these were between 55 and 74.

Thirty-five neoplasms were treated by surgical excision and covering with a pedicle skin graft. Thirty of these were good results and five bad. In 12 cases the excision was done with a radio-knife, in 9 of these the results were good and in 3 unsatisfactory. Electro-desiccation was done in 13 cases, 7 of these did well and 6 badly. Seventeen cases were treated by the chaoul contact method of X-ray therapy, 12 did well and 5 badly. Twenty-one were treated by superficial X-rays, 14 of these were successful and 7 unsuccessful. Twenty-three patients had a combination of surgery and radiotherapy, in 17 the result was good and in 6 cases it was bad.

H. B. STALLARD.

BOOK NOTICE

Manual of Ophthalmology for Medical Officers. By Lieut.-Col. B. W. RYCROFT, R.A.M.C. 1944.

In the introduction to his "Manual of Ophthalmology for Medical Officers," Lieut.-Col. B. W. Rycroft, R.A.M.C. indicates that the book is for beginners and is intended in particular to emphasise the application of the speciality to service conditions.

One half of the volume is devoted to methods of examination and treatment. Routine procedures are dealt with in a practical manner and the chapter on therapeutic measures should be of value to the nursing as well as the medical profession. It seems doubtful if the