have been known to commit suicide after hearing what has practically amounted to a death sentence, and similar disastrous results have followed an unguarded remark about imminent blindness. Patients also, it is worth while remembering, often half hear, only half understand, and put a totally wrong construction on what has been said to them ; so it behoves one to be careful.

Such a thing happened a good many years ago in relation to a middle-aged female who had been admitted to hospital with double choked discs. It is true that no tragic result occurred to the patient. The tragedy was the other way. As a preliminary she was referred to a neurologist. He entered the ward accompanied by a class of students, and noticing something odd about the facial appearance of the patient, immediately made a mental diagnosis of middle fossa tumour, told his class that such a condition was always due to secondary malignant deposits and searched for a primary focus. Having found a small hard lump in one of the breasts, he went on to demonstrate the case to his class and gave a very gloomy prognosis. The patient was exceedingly angry and immediately took her discharge. She went to another hospital, and there someone remembered his first lines and had the Wassermann reaction done. It was strongly positive, and a course of anti-syphilitic treatment cured the ocular condition. To make certain, however, a surgeon was called in ; he removed the lump in the breast, which turned out to be innocent.

There is an irresistible urge, when taking a class of students, to make lightning diagnoses. Mostly these succeed, but at times, as in this instance, the maker is let down badly. It is not given to all of us to be so handy with the ophthalmoscope as the apocryphal pan-diagnostician, who was said to have diagnosed an ovarian cyst from a moment's observation of the fundus ; and better results will be obtained if we keep an open mind and stick to the known facts. So if any of our readers are smitten with a craving for spot diagnosis, let them remember the Horatian maxim *Festina Lente*.

ABSTRACTS

MEDICAL OPHTHALMOLOGY

- (1) **Burch, F. E. (St. Paul, Minnesota).—Metastatic orbital infection.** *Amer. Jl. Ophthal.*, Vol. XXVI, p. 253, 1943.

(1) **Burch** reviews some of the literature on metastatic orbital infection. He describes the case of a white man, aged 39 years, who developed a left orbital abscess 24 days after a furuncle within

the left ala nasi and basal pulmonary infiltration due to type V pneumococcus. Haemolytic staphylococcus aureus was recovered from both lungs, the furuncle and the blood.

Large doses of sulphathiazole, 32 gm. daily, produced clinical improvement, but the white cell count fell from 32,200 per c.mm. to 1,900 per c.mm.

The left orbital abscess was diagnosed by the insertion of a aspiration needle, and was drained through the lower lid and at a later recurrence through the upper lid. Ipsilateral papilloedema was evident during the illness. After eight and a half months, proptosis was only 1 mm., and vision had recovered to 20/15. Some diplopia persisted in the extreme right upper field.

H. B. STALLARD.

- (2) **Atkinson, W. S. (Waterton, New York).**—A coloured reflex from the anterior capsule of the lens which occurs in mercurialism. *Amer. Jl. Ophthal.*, Vol. XXVI, p. 685, 1943.

(2) **Atkinson** describes a brownish coloured reflex from the anterior capsule of the lens which occurs in cases of chronic mercurialism. This reflex is also found in those who have worked a long time in mercury although they may exhibit no symptoms of mercurialism. The colour of the reflex is deeper in the pupillary area and it varies in different individuals from a light brownish grey to a deep rose brown. Small, round, punched-out defects and cracks often appear in the reflex. These resemble the behaviour of metallic mercury. Glass lenses exposed to mercury vapour present a similar brown reflex. The reflex may appear before signs of mercurialism are present.

H. B. STALLARD.

- (3) **Baer, A. T. (Kansas City, Missouri).**—The vitamins in ophthalmology. *Amer. Jl. Ophthal.*, Vol. XXVI, p. 286, 1943.

(3) **Baer** comments that the modern trend is away from prescribing vitamins in a single pure form and there is a return to including the whole group or complex. The vitamin A requirement is 5,600 I.U. a day, but in disease much higher amounts may be needed. The addition of minute doses of a compound iodine solution greatly adds to the efficiency of vitamin A. The body requirement of vitamin B₁ is a minimum of 300 I.U. or 1 mg. per day for the prevention of beri-beri. The amount of thiamin chloride eliminated by the kidneys is used as an index in the diagnosis of vitamin B₁ deficiency. The parts played by vitamins C, D and K in the maintenance of ocular health and in certain pathological states of the cornea and lens, in particular, are discussed.

H. B. STALLARD.

- (4) **Harley, R. D. (Ancon, Canal Zone).**—**Ocular myiasis (ophthalmomyiasis).** *Amer. Jl. Ophthal.*, Vol. XXVI, p. 742, 1943.

(4) **Harley's** case illustrates how the larval form of a botfly masqueraded as a chalazion. There appeared a pinpoint opening in the skin over the swelling in the upper lid and through this passed minute bubbles. The swelling was incised and there was removed a white flask-shaped mass 15 mm. in length which gave evidence of life. Larva of the human botfly, *dermatobia hominis*, were identified. Removal is facilitated by anaesthetising or killing the pest by mercury bichloride soaks, for it is covered with many tiny bristles which in life make difficult its extraction from the tissues.

H. B. STALLARD.

BOOK NOTICE

Cataract and Anomalies of the Lens. By JOHN G. BELLOWS, M.D., PH.D. London: Henry Kimpton. 1944.

This book is a most valuable addition to ophthalmology, and the author is to be congratulated on the clarity of the text and the excellence of the diagrams and illustrations.

Clinicians and research students alike will find a comprehensive account of both rare and ordinary conditions of the lens, and their interest cannot fail to be stimulated by the continuity and brilliance of the subject matter.

In his account of the metabolism and chemical composition of the lens, as well as the pathogenesis of cataract, it is evident that the author has spent an immense amount of time in surveying the evidence of scientific work. He has set into order a host of recorded facts, and has not only made this subject more intelligible to the ophthalmic surgeon, but has provided a basis of reference for the research student. Wherever possible he has correlated the chemistry of the lens with that of the body as a whole.

The structure and development of the normal lens are well described, as well as the various forms of cataract. There is, in addition, a good account of certain rare forms of cataract, *e.g.*, those due to radiant energy, electricity and toxins, concerning which it was previously difficult to obtain information. Finally, there is an adequate description of operative procedures, together with the complications which may follow them.