STUDIES ON DEVELOPING RETINAL VESSELS*

VII. FLUORIDE-INDUCED VASO-OBLITERATION AND ITS RELATION TO RETINAL MATURITY

BY

CHRISTOPHER PEDLER

Department of Pathology, Institute of Ophthalmology, University of London

SINCE it was first reported that oxygen had a specific vaso-obliterative action on the vessels of the immature retina (Ashton, Ward, and Serpell, 1953), a working hypothesis has been devised and certain experiments carried out in an attempt to understand the intermediate mechanism involved (Ashton, Graymore, and Pedler, 1957). In the latter paper it was suggested that the vessels might close through external pressure from fluid imbibition into the retina as a result of an interruption in glycolysis brought about by ambient hyperoxia. Experiments were also reported wherein vessel closure analogous to that following oxygen exposure could be induced by the intravitreal administration of established inhibitors of glycolysis, such as sodium fluoride, which was found to produce complete retinal vaso-obliteration in a consistently repeatable manner. This finding alone is not sufficient to support the idea of cell swelling as a cause of vaso-obliteration in the immature retina, and many other facets of this hypothesis still require experimental investigation; however, support has recently been provided by experiments (Graymore, 1958) which showed that, under defined conditions *in vitro*, excised retinæ incubated in solutions containing known amounts of sodium fluoride underwent a significant increase in Wet Weight : Dry Weight ratio. Similarly, other experiments have demonstrated that the intravitreal injection of sodium fluoride into the intact living eye can result in a histologically measurable swelling of the inner cell layers of the immature but not of the mature retina (Pedler, 1959). Hence, if retinal swelling is responsible for the vaso-obliteration observed in the living eye, then the vessel closure should also show correlation with the degree of retinal maturity. This paper reports the results of some experiments designed first to investigate this possibility, and secondly to determine whether there is any similarity between the age at which fluoride ceases to affect the retinal vessels and that at which ambient hyperoxia becomes ineffective.

Materials and Methods

(1) **Intravitreal Injection.**—A previous paper on this subject (Ashton and others, 1957) gives details of the method used for intravitreal injection, but one small though important detail has been changed:

* Received for publication January 16, 1959.

It is now considered essential for the tip of the needle to pass right through the vitreous to impinge upon the surface of the retina. If this is not done, it is common for some of the injected fluid to leak back along the needle track and to escape from the eye. When the retina is touched, this appears to rupture the vitreous surface so that flow takes place into the subhyaloid space.

(2) **Direct Observation.**—The results shown in the Table were all obtained by a method which allows both cinephotography and medium-power microscopic examination of the living retina using a stereoscopic optical technique. This method is fully described elsewhere (Ashton and Cook, 1954; Pedler, 1957) and no significant changes in it have been made.

TABLE
RELATIONSHIP BETWEEN AGE AND VASO-OBLITERATIVE ACTION OF INTRA-VITREALLY INJECTED SODIUM FLUORIDE

Exp. No.	Age (days)	Weight (g.)	Dose (ml.)	Concentration (per cent.)	Degree of Closure	Time at which Maximum Closure was Seen (min.)
192	6	210	0.2	0.61	++	22
194	7	215	0.2	0.61	++	23
228		160	0.2	0.01M	+	22
188	9	230	0.3	0.61	++	12
221		250	0.2	0.01M	++	26
293	10	245	0.2	0.61	++	6
177	11	300	0.2	0.61	++	18
302	12	230	0.2	0.61	++	20
197	14	230	0.2	0.61	++	5
229		235	0.2	0.61	++	24
237		300	0.2	0.61	+	24
281		260	0.2	0.61	++	7
297	15	295	0.2	0.61	++	26
230	18	270	0.2	0.61	++	18
274	19	370	0.2	0.61	0	—
278		420	0.18R 0.2L	0.5 0.5	++ +	24 Not recorded
236	20	340	0.2	0.61	+	23

contin.

contin.

Exp. No.	Age (days)	Weight (g.)	Dose (ml.)	Concentration (per cent.)	Degree of Closure	Time at which Maximum Closure was Seen (min.)
231	21	300	0.2	0.61	+ +	23
284		360	0.2	0.61	+	10
279	22	390	0.2	0.61	0	—
241	25	330	0.2	0.61	+ +	28
270	30	390	0.2	0.61	+ +	18
245		440	0.2	0.61	0	—
271	34		0.2R 0.2L	0.61 0.61	0 + +	— 28
235	42	510	0.2	0.61	+	18
250	50	700	0.2R 0.2L	0.01M 0.61	+ +	24 20
251	62	850	0.2R 0.2L	0.61 0.61	0 0	— —
291	73	1,050	0.25	0.61	0	—
180	Adult	—	2.6	0.61	0	—
198		3,400	0.2	0.61	0	—
204		2,900	0.5	0.61	0	—
226		3,300	0.25	0.61	0	—
249		3,600	0.2R	0.61	0	—
			0.2L	0.61	0	—

Results and Interpretation

The Table shows the effect of intravitreally injected sodium fluoride on the retinal vessels of kittens of different ages, starting at 6 days and ending with the results of five experiments performed on fully adult animals. In all cases, unless otherwise stated, the injection consisted of 0.2 ml. of an 0.61 per cent. solution of Analar sodium fluoride. The sixth column from the left indicates whether the resultant vaso-obliteration was complete (++), incomplete (+), or absent (0), and the right-hand column shows the time after injection at which maximum closure was seen. The degree of vaso-obliteration produced in any single case was assessed only at the site of injection, the reason being that the extent of the closure depended in some measure upon the spread of the injected sodium fluoride within the subhyaloid space and the spread of the fluoride in turn depended upon the exact position of the needle tip and was thus subject to slight variations. Where complete

vaso-obliteration is recorded, however, this indicates local obliteration of all vessels with the occasional exception of the main veins. Where incomplete closure is recorded this indicates that some part of the arterial circulation was not completely closed by the injection. The zero sign has only been used where the sodium fluoride failed to produce any significant vaso-obliteration. The closure produced by sodium fluoride occurs in a specific sequence (Ashton and others, 1957), and can easily be distinguished from other changes in the calibre of the vessels. For example, in a few experiments on older animals (not included in the Table), some obliteration was noted within 2 or 3 minutes of injection. It could be seen that this took the form of a localized and uneven change in thickness of the vessel wall and was not due to changes in the surrounding retinal tissue. Furthermore, it was rapidly reversed by intravitreal injection of 1 per cent. Amethocaine. The true fluoride-induced closure was unaffected by this substance, and would in fact proceed to completion in its presence, suggesting that vascular spasm was involved as a separate entity in these experiments. Despite these factors, it still remains apparent that the reactivity of the retinal vessels to local sodium fluoride starts to become inconstant between the ages of 14 and 21 days and becomes gradually less with increasing age until after approximately 50 days the vessels are apparently unaffected. It has previously been reported (Ashton and others, 1954) that a raised ambient oxygen concentration will produce total or severe vaso-obliteration from 1 to 14 days of age, and mild vaso-obliteration from 15 to 21 days of age, and it is doubtful or absent after this period. Joint consideration of these two groups of experiments indicates that the developing retinal vessels gradually cease to react to both hyperoxia and sodium fluoride within age limits which are comparable if it is accepted that an intravitreal injection of an enzyme poison is probably a more potent procedure than raising the ambient oxygen concentration.

A correlation of this nature does not of course suggest that the mechanisms involved are identical, and considerable experimental work is still necessary before the degree of similarity between the vaso-oblitative properties of the two agents can be finally established.

Summary

(1) Experiments are described which show that the vaso-oblitative effect of sodium fluoride on the vessels of the immature retina starts to become inconstant between the ages of 14 and 21 days, and becomes gradually less in increasingly older animals, until after approximately 50 days the calibre of the vessels is no longer affected.

(2) This finding is related to the well-established correlation between age and the vaso-oblitative effect of ambient hyperoxia.

It is a pleasure to record the encouragement and helpful criticism I have received from Prof. Norman Ashton during the performance of these experiments.

I should also like to thank Mr. G. Knight, Miss E. Robins, and Miss E. FitzGerald for their assistance. The Medical Research Council kindly made a grant towards the expenses entailed in this work.

REFERENCES

- ASHTON, N., and COOK, C. (1954). *Brit. J. Ophthalm.*, **38**, 433.
 ———, GRAYMORE, C., and PEDLER, C. (1957). *Ibid.*, **41**, 449.
 ———, WARD, B., and SERPELL, G. (1953). *Ibid.*, **37**, 513.
 ———, ———, ——— (1954). *Ibid.*, **38**, 397.
 GRAYMORE, C. (1958). *Ibid.*, **42**, 348.
 PEDLER, C. (1957). *Ibid.*, **41**, 174.
 ——— (1959). *Ibid.*, **43**, 559.