


NEW
PRODUCT
FOR GLAUCOMA

Sno-Pilo[®]

pilocarpine you can always
be sure of.




Sno-Pilo is presented in a specially designed, easy-to-use, plastic dropper bottle containing 10ml pilocarpine hydrochloride eye drops. There are five strengths available: 0.5% w/v, 1.0% w/v, 2.0% w/v, 3.0% w/v and 4.0% w/v. PL 0033/0064 - 0068. Full prescribing information is available. Smith & Nephew Pharmaceuticals Ltd., Bessemer Road, Welwyn Garden City, Hertfordshire, United Kingdom.




Opulets

eye drops

Sterile, single-dose, individually wrapped and labelled one-piece dropper presentation designed to eliminate the risk of cross infection.



No other ophthalmic dropper presentation conforms so closely to DHSS recommendations.



Opulets presentations

Drops

Fluorescein Sodium B.P. 1.0%
Sulphacetamide Sodium B.P. 10.0%
Chloramphenicol B.P. 0.5%

Each in cartons of 20 sterile, sealed droppers.
Basic NHS Price £2.00

Ointments

There is also a range of Opulets single dose, sterile ointment capsules for treating eye or ear infections.

- 1 Outer film sachet maintains sterility of dropper. Contents, batch and expiry date clearly marked on each sachet.
- 2 Strong, simple-to-use disposable plastic dropper, moulded as one unit.
- 3 Integrally moulded non-replaceable twist-off top. Cannot become loose or be resealed.
- 4 Formula identification moulded into plastic, cannot be erased.
- 5 Clearly visible fill.
- 6 Dropper neck design ensures easily controlled, uniform drop.
- 7 Finger pressure points for optimum expulsion and control of drops.
- 8 Easily opened sachet.



Further information available on request

Pharmax Limited Bourne Road Bexley Kent DA5 1NX Telephone Crayford 526551

NEW
PRODUCT
FOR GLAUCOMA

Sno-Pilo

pilocarpine you can always
be sure of.




Sno-Pilo is presented in a specially designed, easy-to-use, plastic dropper bottle containing 10ml pilocarpine hydrochloride eye drops. There are five strengths available: 0.5% w/v, 1.0% w/v, 2.0% w/v, 3.0% w/v and 4.0% w/v PL 0033/0064 - 0068. Full prescribing information is available. Smith & Nephew Pharmaceuticals Ltd., Bessemer Road, Welwyn Garden City, Hertfordshire, United Kingdom.



Alcon

Pneumatograph



THE "PTG"

THE FIRST tonograph which measures intraocular pressure by applanating the eye ... the Alcon Applanation Pneumatograph. It also brings a new dimension to the rapidly developing field of ocular pulse studies.

The Alcon Applanation Pneumatograph encompasses the following unique features:

- Soft tip, self-aligning applanating sensor
- Accurate recorder reading directly in mm Hg
- Remotely positionable magnetic digital display
- Modular Design ... allowing for expansion of system
- Two or four minute tonography
- Heat stylus ... no ink required


The Alcon Applanation Pneumatograph is ... exciting ... versatile ... and opens a new horizon for glaucoma management.

"The Pneumatograph has the advantage of being reproducible in different postures: the influence of posture on intraocular pressure can therefore be more exactly determined. Applanation pneumatonography is more versatile as it is portable and does not require a slit lamp or fluorescein, and a permanent written record is available."

*"M. R. JAIN and V. J. MARMION
BRIT. J. OPHTHAL. (1976) 60, 107"*

For further information apply:—

Alcon Laboratories (U.K.) Ltd.,
Ophthalmic Division,
Imperial Way, Watford,
Herts WD2 4YR Tel: 46133



TWENTY-FOUR GOOD REASONS FOR USING MINIMS™

MINIMS-STERILE, SINGLE-DOSE EYE DROPS

There are twenty-four presentations in the Minims single-dose range. Each one selected for its usefulness in day to day clinical situations.

Minims solutions are sterile and, because of the over-wrap and special cap closure, remain so right up to the moment of application. No preservatives are included in the formulations, so Minims may be used in open eye surgery.

Full prescribing information is available from
Smith & Nephew Pharmaceuticals Limited
Welwyn Garden City, Herts, England



PL: 0033/5000-17, 0033/5019-22, 0033/0048, 0033/0050, 0033/0055.



So how long should cycloplegia persist?

Ideally, as most ophthalmologists would agree, cycloplegia should last long enough to complete refraction – and as little as possible afterwards.

Mydriacyl is the most effective way to meet these criteria. Two drops, 1 to 5 minutes apart permits you to start refracting within 20 minutes. If you are not ready to attend to the patient at that time, instillation of a further drop (repeatable at 30 minute intervals) maintains maximum cycloplegia.

Mydriacyl makes life easier for you – easier for your patients.

Even without use of miotics fair accommodation returns within 3 to 4 hours, and complete recovery within 6 hours. Add a miotic, and patients can resume normal life even sooner.

Which is a vast improvement on conventional techniques often giving patients blurred vision for up to 48 hours.

Supplied: For cycloplegia – Mydriacyl 1.0%. For mydriasis (with moderate, short-acting cycloplegia) – Mydriacyl 0.5%. Both in 15cc Drop-tainer dispenser. Rx only.

Caution: as with any mydriatic, caution should be used when intraocular pressure is high or unknown, or when anterior chamber is shallow. Contraindicated in narrow angle glaucoma.

“Recovery of accommodation is rapid and the cycloplegic effect is gone in 2 to 4 hours, with complete return of accommodation within 6 hours”.

B. C. Gettes, Philadelphia Archives of Ophthalmology May 1961.

MYDRIACYL
MYDRIACYL
(TROPICAMIDE)

for timed mydriasis and cycloplegia

Ophthalmic Division, Alcon Laboratories (UK) Ltd, Imperial Way, Watford, Herts, England WD2 4YR
Telephone Watford 46133. Telex Number 923709

Alcon

World leaders in ophthalmic therapy.