

THE NEW REDUCED EYE ADAPTED TO ACCOMMODATION

BY

V. C. VERBITZKY,

ASSISTANT IN THE OPHTHALMIC CLINIC, UNIVERSITY OF ODESSA.

IN my recent publications ¹ and ², I have attempted to present a reduced eye, differing essentially in principle of construction from all others hitherto offered. I have composed the following system of equations, according to the exact schematic eye of Gullstrand³:—

$$\text{Refractive power } \frac{n-1}{R} = 58.64D \quad . \quad . \quad \text{I and}$$

$$\text{Spherical aberration } \frac{3.2^2}{8n(n-1)R} = 0.387 \text{ mm.} \quad \text{II.}$$


From these the constants of the reduced eye are determined: R = 6.8 mm., and n = 1.4. The values are much higher than the constants of the generally used reduced eyes of Listing⁴, Donders⁵ and Gullstrand³, but they are less than the constants of von Hasner⁶. Thus my reduced eye occupies the middle place. The following table shows the advantages of my reduced eye:—

	Schematic eye of Gullstrand.	Reduced eye of Donders.		Reduced eye of Hasner.		Reduced eye of Gullstrand.		Reduced eye of Verbitzky.	
			Error in %		Error in %		Error in %		Error in %
Refractive power...	58.64D	66.67	13.7	66.67	13.7	58.48	0.3	58.82	0.3
Astigmatism of oblique incidence (at $\angle a = 5^\circ$)	0.086D	0.104	20.9	0.083	3.5	0.092	7.0	0.084	2.3
Spherical aberration (at r = 2 mm.)	1.0D	2.33	133.0	0.80	20.0	1.56	56.0	0.99	1.0
Length of the eye...	24.0mm.	19.3	19.6	22.2	7.5	22.2	7.5	23.4	2.5
Radius of curvature of the corneal surface	7.7mm.	5.0	35.1	7.5	2.6	5.7	26.0	6.8	11.7
Radius of curvature of the retinal surface	10.5mm.	8.4	20.0	9.7	7.6	9.7	7.6	10.2	2.9

In the present work I shall try to adapt the reduced eye to accommodation. In the drawing is represented the contour (optical) of the reduced eye. In order that such an eye shall closely correspond to the schematic eye it must possess coincidence of its contours and equality of its angles. For the exact schematic eye of Gullstrand⁷ it is known that DC = 4.757 mm. at rest and = 5.136 mm. at maximal accommodation. Then, taking into consideration that the radius of curvature of cornea R remains without change in the

schematic eye in accommodation, we deduce from the formula $\sin \phi = \frac{DC \sin \alpha}{R}$ for the middle $\angle \alpha$ at 5° two values of the $\angle \phi$:

3.045° at rest and 3.288° at maximal accommodation. Further, applying the known coefficient of augmentation in diaphragms for the schematic eye of Gullstrand⁷ $K = 0.909$ at rest and $K = 0.941$ at maximal accommodation from the formula $K = \frac{n \tan \alpha}{n' \tan \pi}$ [where $\alpha = 5^\circ$, n (air) = 1, n' (vitreous humour) = 1.336] we shall



obtain for the angle ω two values: 4.121° and 3.981° . From the correlation $\alpha - \phi = \omega - \psi$ we shall have the $\angle \psi$ the two values: 2.166° and 2.269° . But, since $n = \frac{\sin \phi}{\sin \psi}$ we determine two values for the index of refraction: 1.4057 and 1.4487 at rest and at maximal accommodation. In the schematic eye of Gullstrand³ the maximal accommodation = 10.6D. Then the change of n by 1 dioptré of accommodation = $\frac{1.4487 - 1.4507}{10.6} = 0.00406$, or, taking the round figure, $\Delta n = 0.004$.

The refractive power of the reduced eye proposed by me = 58.82D, at maximal accommodation = $58.82 + 10.6 = 69.42$. Hence, by

the formula $D = \frac{n - 1}{R} = \frac{0.4 + 0.004 - 10.6 - 1}{R} = 69.42$ we shall find $R = 6.373$ mm. The radius of curvature changes from 6.8 to 6.373. On each one dioptré the change will be $\frac{0.427}{10.6} = 0.04028$, or $\Delta R = 0.04$ mm.

Thus we obtain the following simple rule:—*For every dioptré of accommodation of the reduced eye the index of refraction n increases by 0.004 and the radius of curvature R diminishes by 0.04 mm.* How satisfactory is the effect of accommodation in this reduced eye will be seen in the following table:—

Accommodation.	R	n.	D it must be.	D in reality.	Difference.	Error of the accommodation in %
	mm.					
0D	6.80	1.400	58.8	58.8	0	0
1	6.76	1.404	59.8	59.8	0	0
2	6.72	1.408	60.8	60.7	-0.1	5
3	6.68	1.412	61.8	61.7	-0.1	3
4	6.64	1.416	62.8	62.6	-0.2	5
5	6.60	1.420	63.8	63.6	-0.2	4
6	6.56	1.424	64.8	64.6	-0.2	3
7	6.52	1.428	65.8	65.6	-0.2	3
8	6.48	1.432	66.8	66.7	-0.1	1
9	6.44	1.436	67.8	67.7	-0.1	1
10	6.40	1.440	68.8	68.8	0	0
11	6.36	1.444	69.8	69.8	0	0
12	6.32	1.448	70.8	70.9	+0.1	1

Whether my rule for the accommodated reduced eye is the most suitable one, or some other correlation would be better, will be seen in further works. But the numberless problems of accommodation which require solution denote persistently, I think, the necessity of the construction of an accommodated reduced eye.

LITERATURE

1. Verbitzky, V.—“The New Attempt of the Simplification of the Optical System of the Eye.” *The Review of the Pure and Applied Science*, Odessa, 1921. (Russian.)
2. Verbitzky, V.—“Zur Frage von den schematischen und reduzierten Augen.” *Klin. Monatsbl. f. Augenheilk.*, Vol. LXVIII, 1922.
3. Gullstrand, A.—“Zusätze im Handb. d. physiol. Optik. v. Helmholtz's.” 3 Aufl., 1909.
4. Helmholtz, H. v.—“Handbuch d. physiolog. Optik.” 2 Aufl., 1896.
5. Donders, F.—“On the Anomalies of Accommodation and Refraction of the Eye,” 1864.
6. Hasner, v.—“Über das reduzierte Auge.” *Arch. f. Augen. u. Ohrenheilkunde*. Vol. VII, 1878.
7. Rohr, M.—“Zur Dioptrik des Auges.” Asher u. Spiro. *Ergebnisse d. Physiologie*, Vol. VIII, 1909.