

BJO

British Journal of Ophthalmology

Editorials

Proliferative vitreoretinopathy

The term proliferative vitreoretinopathy (PVR) is the latest and most enduring of a series of terms used since the early 1960s to describe the proliferation and migration of cells to form fibrocellular membranes on and around the retina. The clinical picture of fibrous contraction, leading to inoperable retinal detachment, was initially thought to be a process originating within the vitreous, the contracted gel leading to retinal shortening and irreversible fibrotic contraction, hence the term massive vitreous retraction (MVR). Both Taylor-Smith¹ and Cibis,² however, recognised and reported the significance of epiretinal membranes in the development of complicated retinal detachments.

With the introduction of closed intraocular microsurgery by Machemer,³ direct access to the preretinal and subretinal spaces enabled him and others to confirm that PVR was in fact a process of periretinal cellular proliferation, the more clinically significant and damaging element of which was epiretinal. The term massive periretinal proliferation (MPP) was therefore coined by Machemer,⁴ but subsequently changed to PVR, at the behest of the Retina Society.⁵

Proliferative vitreoretinopathy describes a fibrocellular cascade, in which both glial and retinal pigment epithelial cells participate⁶⁻¹² and, encouraged to proliferate, migrate, and undergo metaplasia by fibronectin, growth factors, and other stimulants,¹³⁻²² form contractile membranes on the surface of the retina and beneath it. Contrary to previous thinking, the role of the vitreous gel in the development of PVR is comparatively small. Posterior vitreous detachment generally initiates the chain of events leading to PVR, by creating dehiscences in the internal limiting lamina and/or full thickness retinal breaks, but cells released into the vitreous gel proliferate only locally, do not form membranes, and do not cause contraction of the gel. Proliferation of cells forming contractile membranes on the posterior surface of the detached gel and at the vitreous base, nevertheless, contributes to the forces acting on the anterior retina (anterior PVR).

The simplest membranes are those that derive from glial cells and commonly develop in the absence of full thickness retinal breaks. They sometimes result in areas of localised retinal shortening, as in macular pucker, but are often found in the eyes of asymptomatic individuals, in response to small dehiscences in the internal limiting lamina.^{23,24} Such membranes have only limited contractile properties and rarely lead to extensive retinal detachment.

In more complex membranes, usually resulting from the release of cells into the retrohyaloid space (through

full thickness retinal breaks following posterior vitreous detachment), RPE cells together with glial cells proliferate, migrate, and form strongly contractile fibrocellular sheets.^{4,6-11} These commonly lead to gross retinal shortening and may distort, enlarge, and hold open retinal breaks, leading to persistent, recurrent, and/or progressive retinal detachment. Proliferation of similar cells in the subretinal space, however, uncommonly gives rise to cohesive membrane formation and adhesion to the undersurface of the retina is usually sparse and weak, so that gross retinal shortening is uncommon.²⁵⁻²⁷

The paper by Mietz and Heimann, on page 874 of this issue of *BJO*, indicates that the time course during which the process of fibrocellular proliferation, migration, and contraction occurs, is remarkably consistent, adding weight to the view that the cascade of events leading up to development of clinical PVR is a consistent one and, hence, that its interruption by therapeutic agents at one or more points in the chain, may be feasible.

Treatment of PVR can be preventive or curative. Prevention is better than cure and, as Scott²⁸ and others have suggested, the most promising solution to the prevention of PVR lies in the success of primary retinal reattachment surgery. Nevertheless, PVR can, and often does, supervene even after initially appropriate and successful surgery for retinal detachment, or even before any attempt at repair has been made. Work has been undertaken in the past 15 years by Binder, Blumenkranz, Tano, Wiedemann and others,²⁹⁻⁴¹ to test a wide variety of agents, targeted at various points along the PVR cascade, most of them more or less successful in the laboratory, but with only limited success in the clinical arena. Why this should be so is a matter for conjecture, but it is probably because the timing of delivery in the animal model is very different from that which presents as an opportunity in the clinical context. This is what makes the paper by Mietz and Heimann so interesting and apposite. The tight time course of PVR development should allow us to target treatment at a specific stage, identifiable by the time at which each part of the process occurs, not only in primary cases, but in recurrences.

Prevention of PVR, rather than its cure, is particularly urgent at the present time because surgical treatment of complex retinal detachments is costly, in terms of a skilled workforce and expensive equipment, while the quality of visual outcomes, even in successful cases, is open to question. Although recent studies by some workers have indicated that patients with retinal detachments

complicated by PVR in one eye, but with a healthy fellow eye, are mostly pleased to have undergone surgery to the affected eye (even when the results seem to be of very limited value in visual terms),^{42 43} work by Andenmatten and Gonvers⁴⁴ suggests that surgery for PVR in the eye of a patient with a healthy fellow eye, is helpful only if binocular single vision can be restored and this was found to be possible in less than one third of cases.

The use of a 'cocktail' of agents aimed at arresting the progress of PVR at an early stage in its development, each targeted at a specific point in the fibrocellular cascade, could be more effective than any single agent alone. A multicentre, randomised, clinical trial of just such a 'cocktail' of agents is about to be undertaken in the UK. The ultimate goal of the study is to find an agent or combination of agents which can be given safely to all patients who develop retinal breaks with posterior vitreous detachment, whether they require surgical intervention or just retinopexy. Such a compound, given locally, systemically, or both, would ideally prevent the migration and proliferation of cells which form membranes, without hindering the reparative processes in the retina necessary to seal retinal breaks. The problems concerned with developing such an agent are, in some ways, common to those experienced in the fight against neoplastic disease, the treatment sometimes being worse than the cure.

Advances in pharmacology, and other therapeutic modalities, combined with improvements in methods of retinopexy engendered by advances in laser technology, are most likely to result in the development of new treatment methods which will rid us of the scourge of PVR.

PETER K LEAVER

Moorfields Eye Hospital,
City Road,
London EC1V 2PD

- 1 Smith TR. Pathological findings after retinal surgery. In: Schepens S, ed. *Importance of vitreal body in retinal surgery with special emphasis on reoperation*. St Louis: C V Mosby, 1960:61.
- 2 Cibis PA. *Vitreoretinal pathology and surgery in retinal detachment*. St Louis: C V Mosby, 1965.
- 3 Machemer R, Buettner H, Norton EWD, Parel JM. Vitrectomy: a pars plana approach. *Trans Am Acad Ophthalmol Otolaryngol* 1971; 75: 813.
- 4 Machemer R, Laqua H. Pigment epithelium proliferation in retinal detachment (massive periretinal proliferation). *Am J Ophthalmol* 1975; 80: 1.
- 5 The Retina Society Terminology Committee. The classification of retinal detachment with proliferative vitreoretinopathy. *Ophthalmology* 1983; 90: 121.
- 6 Laqua H, Machemer R. Glial cell proliferation in retinal detachment (massive periretinal proliferation). *Am J Ophthalmol* 1975; 80: 602.
- 7 Van Horn DL, Aaberg TM, Machemer R, Fenzl R. Glial cell proliferation in human retinal detachment with massive periretinal proliferation. *Am J Ophthalmol* 1977; 84: 383.
- 8 Smith RS, van Heuven WAJ, Streeten B. Vitreous membranes: a light and electron microscopic study. *Arch Ophthalmol* 1976; 94: 1556.
- 9 Clarkson JG, Green WR, Massof D. A histopathologic review of 168 cases of preretinal membrane. *Am J Ophthalmol* 1977; 84: 1.
- 10 Kampik A, Kenyon KR, Michels RG, Green WR, de la Cruz ZC. Epiretinal and vitreous membranes: comparative study of 56 cases. *Arch Ophthalmol* 1981; 99: 1445.
- 11 Grierson I, Rahi AHS. Structural basis of contraction in vitreal fibrous membranes. *Br J Ophthalmol* 1981; 65: 737.
- 12 Roth AM, Foos RY. Surface wrinkling retinopathy in eyes enucleated at autopsy. *Trans Am Acad Ophthalmol Otolaryngol* 1971; 75: 1047.
- 13 Weller M, Heimann K, Wiedemann P. Immunohistochemical findings in proliferative retinal diseases: on the significance of fibronectin, macrophages and transferrin. *Fortschr Ophthalmol* 1989; 86: 121.
- 14 Weller M, Wiedemann P, Breggen M, Heimann K. Vitronectin: mediator of cell adhesion in proliferative vitreoretinopathy? *Fortschr Ophthalmol* 1990; 87: 221.
- 15 Hiscott P, Waller HA, Grierson I, Butler MG, Scott D. Local production of fibronectin by ectopic human retinal cells. *Cell Tissue Res* 1992; 267: 185.
- 16 Wilson-Holt N, Khaw P, Savage F, Grierson I. The chemoattractant activity of the vitreous to human scleral fibroblasts following retinal detachment and proliferative vitreoretinopathy. *Br J Ophthalmol* 1992; 76: 159.
- 17 Campochiaro PA, Jerdan JA, Glaser BM. Serum contains chemoattractants for human retinal pigment epithelial cells. *Arch Ophthalmol* 1984; 102: 1830.
- 18 Campochiaro PA, Glaser BM. Platelet-derived growth factor is chemotactic for human retinal pigment epithelial cells. *Arch Ophthalmol* 1985; 103: 576.
- 19 Connor TB, Roberts AB, Sporn MB, Danielpour D, Dart LL, Michels RG, et al. Correlation of fibrosis and transforming growth factor-beta type 2 levels in the eye. *J Clin Invest* 1989; 83: 1661.
- 20 Gaudric A, Glacet-Bernard A, Clement G, Falquerho L, Barrिताult D, Coscas G, et al. Transforming growth factor beta in the vitreous of patients with epiretinal proliferation. *Ophthalmologie* 1990; 4: 51.
- 21 Wiedemann P. Growth factors in retinal disease: proliferative vitreoretinopathy, proliferative diabetic retinopathy, and retinal degeneration. *Surv Ophthalmol* 1992; 35: 373.
- 22 Campochiaro PA, Jerdan JA, Glaser BM. The extracellular matrix of human retinal pigment epithelial cells *in vivo* and its synthesis *in vitro*. *Invest Ophthalmol Vis Sci* 1986; 27: 1615.
- 23 Foos RY. The spectrum of nonvascular proliferative extra-retinopathies. In: Nicholson, ed. *Ocular pathology update*. Los Angeles, 1980: 107.
- 24 Foos RY. Surface wrinkling retinopathy. In: Freeman HM, Hirose T, Schepens CL, eds. *Vitreous surgery and advances in fundus diagnosis and treatment*. New York: Appleton Century Crofts, 1977: 23.
- 25 Sternberg P, Machemer R. Subretinal proliferation. *Am J Ophthalmol* 1984; 98: 456.
- 26 Schwartz D, de la Cruz Z, Green WR, Michels RG. Proliferative vitreoretinopathy: ultrastructural study of 20 retorectinal membranes removed by vitreous surgery. *Retina* 1988; 8: 275.
- 27 Lewis L, Aaberg M, Abrams GW, McDonald HR, Williams GA, Mieler WF. Subretinal membranes in proliferative vitreoretinopathy. *Ophthalmology* 1989; 96: 1403.
- 28 Scott John D. Duke-Elder Lecture. Prevention and perspective in retinal detachment. *Eye* 1989; 3: 491.
- 29 Binder S, Skorpice CH, Paroussis P, Kulnig W, Menapace R, Prokesch E, et al. High-energy electrons used to inhibit experimental intraocular proliferation and detachment. *Ophthalmologica* 1987; 195: 128.
- 30 Binder S, Bonnet M, Velikay M, Gerard JP, Stolba U, Wedrich A, et al. Radiation therapy. A prospective randomised study. *Graefes Arch Clin Exp Ophthalmol* 1994; 232: 211.
- 31 Blumenkranz MS, Ophir A, Clafin AJ, Hajek A. Fluorouracil for the treatment of massive periretinal proliferation. *Am J Ophthalmol* 1982; 94: 458.
- 32 Blumenkranz MS, Hernandez E, Ophir A, Norton EW. 5-Fluorouracil: new applications in complicated retinal detachment for an established antimetabolite. *Ophthalmology* 1984; 91: 122.
- 33 Blumenkranz MS, Clafin A, Hajek AS. Selection of therapeutic agents for intraocular proliferative disease: cell culture evaluation. *Arch Ophthalmol* 1984; 102: 598.
- 34 Tano Y, Chandler D, Machemer R. Treatment of intraocular proliferation with intravitreal injection of triamcinolone acetonide. *Am J Ophthalmol* 1980; 90: 810.
- 35 Wiedemann P, Lemmen K, Schmiedl R, Heimann K. Intraocular daunorubicin for the treatment and prophylaxis of traumatic proliferative vitreoretinopathy. *Am J Ophthalmol* 1987; 104: 10.
- 36 Wiedemann P, Leinung C, Hilgers RD, Heimann K. Daunomycin and silicone oil for the treatment of proliferative vitreoretinopathy. *Graefes Arch Clin Exp Ophthalmol* 1991; 229: 150.
- 37 Chandler DB, Hida T, Sheta, Proia AD, Machemer R. Improvement in efficacy of corticosteroid therapy in an animal model of proliferative vitreoretinopathy by pretreatment. *Graefes Arch Clin Exp Ophthalmol* 1987; 225: 259.
- 38 Hui YN, Liang HC, Cai YS, Kirchoff B, Heimann K. Corticosteroids and daunomycin in the prevention of experimental proliferative vitreoretinopathy induced by macrophages. *Graefes Arch Clin Exp Ophthalmol* 1993; 231: 109.
- 39 Kuriyama S, Ohuchi T, Yoshimura N, Honda Y, Hiraoka M, Abe M, et al. Evaluation of radiation therapy for experimental proliferative vitreoretinopathy in rabbits. *Graefes Arch Clin Exp Ophthalmol* 1990; 228: 552.
- 40 Alvira G, Hartzler M, Blumenkranz MS. Heparin inhibits cell-mediated collagen gel contraction. *Invest Ophthalmol Vis Sci* 1986; 27: 305.
- 41 Radtke ND, Weinsieder AD, Ballou RJ. Pharmacological therapy for proliferative vitreoretinopathy. *Graefes Arch Clin Exp Ophthalmol* 1986; 224: 230.
- 42 Sullivan PM, Luff AJ, Jubois SA, Canning CR. Patient satisfaction following vitreoretinal surgery. *Eye* 1993; 7: 433.
- 43 McCormack P, Simcock PR, Charteris D, Lavin MJ. Is surgery for proliferative vitreoretinopathy justifiable? *Eye* 1994; 8: 75.
- 44 Andenmatten R, Gonvers M. Sophisticated vitreoretinal surgery in patients with a healthy fellow eye. An 11 year retrospective study. *Graefes Arch Clin Exp Ophthalmol* 1993; 231: 495.