

BRIEF REVIEWS ON ASPECTS OF AGING AND THE EYE

Aging and the cornea

R G A Faragher, B Mulholland, S J Tuft, S Sandeman, P T Khaw

Aging, the persistent decline in age specific fitness of an organism as a result of internal physiological deterioration, is a common process among multicellular organisms.¹ In humans, aging is usually monitored in relation to time, which renders it difficult to differentiate between time dependent biological changes and damage from environmental insults. There are essentially three types of aging at work in any adult tissue; the aging of long lived proteins, the aging of dividing cells, and the aging of non-dividing cells.² Dividing cells may be derived from renewing populations in which the rate of cell loss and division is great. An example is the corneal epithelium in which complete turnover occurs within 5-7 days after terminal differentiation.^{3,4} Conditional renewal populations, which normally have an extremely low proliferation rate, can also produce dividing cells in response to extrinsic stimuli. Stromal keratocytes are a prime example of a conditional renewing population.⁵ Corneal endothelial cells retain the capacity to undergo mitosis and conditional renewal in humans although they very seldom do so.⁶⁻⁹ Non-dividing cells are those from static cell populations (exemplified by cerebral neurons) which never divide during adult life.³

Corneal aging produces both structural and functional changes. These changes in turn can affect the ability of the organ to refract light, to repair itself, and to protect itself and the internal structures of the eye.¹⁰ A variety of corneal aging changes have been reported. However, as it is difficult to distinguish age specific deterioration from degenerations modified by environmental and genetic factors, we think it is helpful to consider these alterations within the broader framework of the aging process. The study over the past 30 years of isolated cells in culture as a model system for aging changes has greatly advanced our understanding of these concepts.

Cell aging

Normal adult cell populations do not divide indefinitely either in the culture dish or in the body.^{3,11} Cellular senescence or replicative failure is the process that imposes a limit on their replicative lifespan, and it is thought that cell senescence acts as a powerful tumour suppression mechanism which thus lengthens the healthy reproductive lifespan of the organism.¹² However, the emergence of senescent cells also contributes to the aging process in mitotically competent tissues. This theory, the cell hypothesis of aging, proposes that the gradual accumulation of senescent cells is the primary event that leads to the development of age linked degenerative changes in tissue.^{13,14} A key feature of this hypothesis is the presence of senescent cells, but it says nothing about the mechanism that causes the cells to become senescent in the first place. These mechanisms are considered below.

What are senescent cells?

A concept of cell senescence can perhaps best be appreciated after a consideration of what it is not. Senescence is distinct from quiescence, a transient growth arrest state,

also known as contact inhibition. Rather confusingly, both senescence and quiescence are referred to as the G0 phase of the cell cycle (sometimes more helpfully distinguished as G0Q and G0S).¹⁵ Senescence is also distinct from cell death, occurring either by apoptosis or necrosis, and it is not a form of terminal differentiation.^{16,17} The phenotypes of growth and senescence are totally distinct cell cycle compartments; there is no such thing as a half senescent cell. Cells that enter replicative senescence acquire two phenotypes: they leave the cell cycle with a G1 DNA content,¹⁸ and they undergo a characteristic series of changes in biology and gene expression that alters the function of the cell.^{13,19} In this latter situation some genes are transcriptionally repressed, some gene expression is upregulated, and some totally senescent specific genes are turned on.²⁰ These changes cover practically every aspect of cell physiology and occur in a highly selective manner.² As many of the changes occur in genes coding for secreted products the senescent cell can potentially affect the surrounding microenvironment. This altered function of senescent cells may thus be the critical phenotype that compromises tissue function and integrity. As these changes have been largely studied *in vitro* it is important to examine the means by which cells become senescent and to question their significance within aged tissue.

Where could senescent cells come from within the cornea?

There are two main routes by which a cell may become senescent; they are explained below.

CONSTITUTIVE CELL SENESCENCE

Replicative failure is often visualised as the cell counting a fixed number of divisions and then entering senescence but, while conceptually straightforward, this is a misleading oversimplification. Rather than simply counting its way to senescence, each time a cell divides it has a given chance of never dividing again, a chance that increases each time the cell replicates until senescence becomes a certainty.^{21,22} However, since the process is controlled by chance, a cell that has divided only once can still be unlucky and enter senescence. The constitutive process is thus rather like playing Russian roulette, the chance of the fatal bullet is fixed, but the outcome is uncertain. Unlike Russian roulette, each time the cell divides extra bullets are loaded into the revolver. In tissue culture the end result of tens of thousands of chance decisions are cell populations which have a mixture of growing and senescent cells, the proportions of which shift in the direction of total senescence as the culture is passaged and the cells divide.²³ In tissue, even very limited division can thus begin to produce senescent cells. The kinetics of constitutive senescence can be explained in terms of the inheritance of chromosomes with progressively shortened telomeric DNA sequences.²⁴

REACTIVE CELL SENESCENCE

This pathway to the senescent state was demonstrated recently and provides a fascinating alternative to the

constitutive route. Current data show that this stage is essentially identical to constitutive senescence but happens in a matter of hours. It has been shown that the induction of the activated H-ras oncogene into growing fibroblasts can trigger senescence.²⁵ This suggests that, rather like apoptosis, senescence can be induced by mutation or mitogenic overload; this may implicate topical treatment with anticancer drugs in the induction of senescence. In contrast with the constitutive route, this pathway appears to require little if any cell division. Thus, senescent cells may appear much more frequently in quiescent tissue than previously thought, particularly if that tissue is in a mutagenic environment. This may be clinically relevant with the increasing use of potentially mutagenic agents such as 5-fluorouracil and mitomycin C to prevent scarring after pterygium excision or glaucoma filtration surgery; particularly since mitomycin C has been shown to rapidly induce senescent-like changes in cultured fibroblasts.²⁶ These drugs soak into sclera, conjunctiva, and cornea, particularly after subconjunctival injection, but probably also after sponge application.²⁷ We have seen prolonged effects on the tissue fibroblasts in the drug treated area that seem unable to divide further despite maximal serum stimulation.²⁸ It is possible that some of this growth arrest is reactive cell senescence although this remains to be proved. This is an important distinction as senescence is irreversible, unlike prolonged growth arrest seen in vitro that may recover.^{29,30} The clinical importance of these observation is that accelerated senescence may thus cause disease within the affected tissue that may not become apparent for many years.

Particularly interesting from the perspective of senescence is the recent observation that removal of the corneal epithelium can trigger apoptosis in the underlying anterior keratocytes.³¹ These cells are then replaced after re-epithelialisation by division and migration from the posterior stroma. This apoptosis repopulation process is believed to form a line of defence against invading viruses,³² it also provides a mechanism by which cell division, and hence the constitutive senescence pathway, may be activated. These processes may be manifest as a decline in the density of keratocytes with advancing age.

The cornea is saturated in light that is potentially mutagenic and reactive senescence can potentially occur in any of the cell layers. Cell turnover within the epithelium is continuous, and thus constitutive senescence has the potential to occur in both the transient amplifying population or, more seriously, in the dividing stem cells. Corneal epithelium has not been examined for the presence of senescent cells but studies of skin strongly suggest they will be present.^{33,34} The behaviour of cultured keratocytes from

old donors is also consistent with an elevated fraction of senescent cells in the biopsy rather than a reduced number of starting cells.³⁵ An age related increase in the number of senescent cells in human endothelium has been observed.^{36,37}

How does cell senescence affect the cornea?

Structural and functional alterations documented in the aging cornea are listed in Table 1. Although insufficient information exists on the senescence of epithelial cells to draw more than speculative observations, senescence is associated with a decreased ability to resist a wide range of physiological stresses. Changes in the ocular surface render the aging cornea more susceptible to infection for various reasons. There is an increase in epithelial permeability with age that may either represent a breakdown of epithelial barrier function⁴⁷ or an increased tear contact time.⁴⁶ Changes in the distribution of integrin subunits in the epithelium could also reduce the epithelial barrier function. The $\alpha 6$ subunit and the $\beta 4$ subunit, components of hemidesmosomes, become discontinuous with age. However, the number and distribution of hemidesmosomes along the basal lamina do not appear to change with age.⁴⁸ A reduced ability of corneal cells to upregulate adhesion molecules and a reduced phagocytic ability of reactive polymorphonucleocytes in response to infection also occur with aging,^{43,44} and this could impair the ability to eliminate a bacterial infection. Epithelial disease, in turn, has the potential to contribute to cell loss within the endothelial layer.⁴⁹

The major cellular component of the corneal stroma is the keratocyte. Few studies have been conducted on these cells, but in vitro studies of senescent dermal and lung fibroblast-like cells have demonstrated constitutive over-expression of collagenase, stromolysin, and elastase.^{50,51} Simultaneously, the expression of tissue inhibitors of metalloproteinases (TIMP 1 and TIMP 2)⁵² are greatly reduced, as is collagen mRNA.⁵³ Fibronectin is produced in an altered form which is a less efficient substrate for cell adhesion,⁵⁴ proteoglycan synthesis falls,⁴⁹ and the migration rate⁵⁵ and the ability of fibroblasts to contract a collagen lattice in vitro also decline.⁵⁶ Lipofuscin and endogenous ceramide levels increase.^{57,58} Gap junction assembly times increase by an order of magnitude and membrane permeability increases sharply.⁵⁹ Specific inhibitors of calcium dependent membrane currents are induced.⁶⁰ In addition, the glycation of corneal collagen produces an increase in intramolecular spacing.⁴² The overall result of these changes is a radical shift of the senescent cell into a highly catabolic phenotype² and, in aging skin, senescent cells have been demonstrated in close proximity to degen-

Table 1 Alterations seen in the aging cornea

Characteristic	Result of change
(1) Changes in shape and optical properties	(i) Steepening of keratometry and a shift from with the rule to against the rule astigmatism ^{38,39} (ii) Transparency is unaffected in central cornea in absence of scar or degeneration ⁴⁰ (iii) Collagen intramolecular and interfibrillar spacing increases—possibly via increased protein glycation ^{41,42} (iv) Increased thickness of Descemet's membrane
(2) Corneal degenerations (influenced by environmental and genetic factors)	(i) Cornea farinata (ii) White limbus girdle (iii) Mosaic degeneration (iv) Deep crocodile shagreen (v) Hassall-Henle bodies (vi) Arcus senilis
(3) Physical properties	(i) Resistance to infection reduced (ii) Failure to upregulate ICAM-1 and reduced inflammatory cell infiltration ⁴³ (iii) Reduced phagocytically active cells after infection ⁴⁴ (iv) Decline in high energy metabolism ⁴⁵ (v) Increased tear contact time ⁴⁶ (vi) Increased epithelial permeability to fluorescein ⁴⁷

erative and disorganised collagen fibrils.³³ The reduced keratocyte density within the aging cornea, the breakdown of collagen fibres, and the appearance of collagen-free spaces may reflect similar changes within this tissue.⁶¹⁻⁶² The increase in lipofuscin granules seen in the aged stroma (corneal farinata) may represent deposition of products from senescent cells.

Some of these factors could adversely affect wound repair. A reduced number of fibroblasts, an inability of many of those cells to divide, the decreased ability for migration, a reduced ability to contract a wound lattice, and the depressed synthesis of collagen are believed to contribute strongly to impaired wound healing in the aging dermis. While no direct evidence for this currently exists in the normal cornea, patients with Werner's syndrome—a hereditary disease characterised by premature fibroblast senescence⁶³—show severely impaired corneal wound healing following cataract surgery.⁶⁴ The implication that senescence may have a detrimental effect on the outcome of corneal surgery is thus strong. Decreased healing of wounds in the aged may be advantageous in some circumstances such as glaucoma filtration surgery, but age has also been identified as the most important individual variable affecting the outcome after refractive surgery; the amount of effective aggression decreasing proportional to increasing age.⁶⁵⁻⁶⁷

Age related changes in the corneal endothelium have been examined clinically and experimentally. It has been estimated that between the ages of 20 and 80 years the annual reduction in cell density averages approximately 0.6%, with concomitant increases in polymethism and pleomorphism.⁶⁸⁻⁷¹ However, as the mean age of the population sample increases, there is an increased spread in the range of the endothelial cell density counts.⁷² This means that the measurement of endothelial cell density is not a reliable index of the chronological age of the cornea, and suggests an environmental influence. Changes in cell density and shape with age have been observed to occur in the human,⁷³ monkey,⁷⁴ rat,⁷⁵ cat,⁷⁴⁻⁷⁶ dog,⁷⁷ and rabbit,⁷⁸ but in each of these species the adult mean cell density (about 2500 cells/mm²) is remarkably constant. Interestingly, it has been noted in the rat, which has the ability for endothelial cell division, that the total reduction in endothelial cell numbers is of the same order of magnitude as occurs in humans, but the cell loss is compressed into the shorter life span of this species.⁷⁵ The biological mechanisms behind the gradual endothelial cell loss during aging remain to be elucidated, but might involve hormonal changes or environmental influences such as ultraviolet irradiation and chemical toxicity. In particular, the degradation of enzymes in the anterior segment that normally metabolise and detoxify hydrogen peroxide and other free radicals may lead to progressive damage to the endothelial layer.⁷⁹ Reduction in endothelial cell numbers and the increased variability in cell size and shape that accompany normal aging may adversely affect endothelial function,⁸⁰ although this reduced function may also be the result of a decline in high energy metabolism with age.⁴⁵ The aged cornea is slower to recover from hypoxic stress,⁸¹ and grafts from older donors usually require a longer post-operative period to attain their final thickness.⁸²⁻⁸³ Although advanced donor age does not preclude the use of a cornea for grafting, the life span of a transplanted endothelial cell is, as yet, unknown. In rabbits, where the endothelial cell layer is able to regenerate, the pattern of corneal endothelial wound healing after transcorneal freezing is slower and less extensive in corneas from adult animals than from young animals.⁸⁴

Conclusions

A wide range of changes occur in the aging cornea, some of which can be linked to the changes seen in aging cells in culture. The lack of uniform culture systems for corneal epithelium and endothelium has limited the study of senescence phenotype in these cell types. Although the growth arrest phenotype of senescence is universal among different cell types the changes in function that accompany it are not, with many growth arrest genes showing high tissue specificity. Only studies of these cell types will allow firm conclusions to be drawn. Simple inactivation of the mechanism of replicative failure is intrinsically undesirable since it apparently functions as an antitumour mechanism. A more sensible strategy appears to be to define in greater detail the functional phenotype of senescence and then to attempt to modify this through therapy, an intriguing clinical possibility for the future.

This work was supported by the Royal College of Ophthalmologists/BUPA fellowship, Peel Medical Research Trust, and Moorfields Special Trustees Locally Organised Research Scheme.

R G A FARAGHER

Ocular Research Group, Department of Pharmacy,
University of Brighton

B MULHOLLAND
S J TUFT

Wound Healing Research Unit,
Institute of Ophthalmology and Moorfields Eye Hospital,
Bath Street, London EC1V 9EL

S SANDEMAN

Ocular Research Group, Department of Pharmacy,
University of Brighton

P T KHAW

Wound Healing Research Unit,
Institute of Ophthalmology and Moorfields Eye Hospital,
Bath Street, London EC1V 9EL

- Rose MR. *Evolutionary biology of aging*. Oxford: Oxford University Press, 1991.
- Campisi J. The biology of replicative senescence. *Eur J Cancer* 1997;33:703-10.
- Leblond CP. Classification of cell populations on the basis of their proliferative behaviour. *Natl Cancer Inst Monogr* 1964;14:119-49.
- Marshall J. The susceptible visual apparatus. In: Marshall J, ed. *Vision and visual dysfunction*. Vol 16. London: Macmillan Press, 1991.
- Tuft SJ, Gartry DS, Rawe IM, Meek KM. Photorefractive keratectomy: implications of corneal wound healing. *Br J Ophthalmol* 1993;77:243-7.
- Treffers WF. Corneal endothelial wound repair in vivo and in vitro. *Ophthalmology* 1982;89:605-13.
- Van Horne DL, Hyndiuk RA. Endothelial wound repair in primate cornea. *Exp Eye Res* 1975; 21:113-24.
- Schultz G, Cipolla L, Whitehouse A, Eiferman R, Woost P, Jumblatt M. Growth factors and corneal endothelial cells. III: Stimulation of adult human corneal endothelial cell mitosis in vitro by defined mitogenic agents. *Cornea* 1992;11:20-7.
- Mishima S. Clinical investigations on the corneal endothelium. *Am J Ophthalmol* 1982;93:1-29.
- Greiner JV, Kenyon KR. In: Albert DM, Jakobiec FA, eds. *Principles and practice of ophthalmology—basic sciences*. Chapter 52. Philadelphia: WB Saunders, 1994.
- Shall S. Mortalisation or reproductive sterility of animal cells in culture. In: Potten CS, ed. *Perspectives on mammalian cell death*. Oxford: Oxford University Press, 1987:184-201.
- Smith JR, Periera-Smith OM. Replicative senescence: implications for in vitro aging and tumor suppression. *Science* 1996;273:63-7.
- Hayflick L. The cell biology of aging. *J Invest Dermatol* 1979;73:8-14.
- Chang E, Harley CB. Telomere length and replicative aging in human vascular tissue. *Proc Natl Acad Sci* 1995;92:11190-4.
- Norwood TH, Smith JR, Stein GH. Aging at the cellular level: the human fibroblast like cell model. In: *Handbook of the biology of aging*. London: Academic Press, 1990:131-54.
- Norsgaard H, Clark BFC, Rattan SIS. Distinction between differentiation and senescence and the absence of increased apoptosis in human keratinocytes undergoing cellular ageing in vitro. *Exp Gerontol* 1996;31:655-68.
- Maciera-Coelho A, Ponten J, Philipson L. The division cycle and RNA synthesis in diploid human cells at different passage levels in vitro. *Exp Cell Res* 1966;42:673-84.
- Gorman SD, Cristofalo VJ. Analysis of the G1 arrest position of senescent W138 cells by quinacrine dihydrochloride nuclear fluorescence. *Exp Cell Res* 1981;167:87-94.
- Sheshadri T, Campisi J. Repression of c-fos and an altered genetic programme in senescent human fibroblasts. *Science* 1990;247:205-9.
- Wistrom C, Villeponteau B. Cloning and expression of SAG: a novel marker of cellular senescence. *Exp Cell Res* 1992;199:355-62.
- Smith JR, Whitney RG. Intracloonal variation in proliferative potential of human diploid fibroblasts: stochastic mechanism for cellular ageing. *Science* 1980;207:82-4.

- 22 Ponten J, Stein WD, Shall S. A quantitative analysis of the ageing of human glial cells in culture. *J Cell Physiol* 1983;117:342-52.
- 23 Cristofalo VJ, Scharf BB. Cellular senescence and DNA synthesis. *Exp Cell Res* 1973;76:419-27.
- 24 Allsopp RC, Vaziri H, Patterson C, Goldstein S, Younglai EV, Futcher AB, et al. Telomere length predicts replicative capacity of human fibroblasts. *Proc Natl Acad Sci* 1992;89:10114-8.
- 25 Serrano M, Lin AW, McCurrach ME, Beach D, Lowe SW. Oncogenic ras provokes premature cell senescence associated with accumulation of p53 and p16INK4a. *Cell* 1997;88:593-602.
- 26 Brenneisen P, Gogo J, Bayreuther K. Regulation of DNA synthesis in mitotic and postmitotic WI38 fibroblasts in the fibroblast stem cell system. *J Cell Biochem* 1993;17D:152.
- 27 Kondo M, Araie M. Concentration change of fluorouracil in the external segment of the eye after subconjunctival injection. *Arch Ophthalmol* 1988;106:1718-21.
- 28 Khaw PT, Doyle JW, Sherwood MB, Grierson I, Schultz G, McGorray S. Prolonged localised tissue effects from 5-minute exposures to fluorouracil and mitomycin C. *Arch Ophthalmol* 1993;111:263-7.
- 29 Khaw PT, Ward S, Porter A, Grierson I, Hitchings RA, Rice NS. The long term effects of 5-fluorouracil and sodium butyrate on human Tenon's fibroblasts. *Invest Ophthalmol Vis Sci* 1992;33:2042-52.
- 30 Khaw PT, Sherwood MB, Mackay SL, Rossi MJ, Schultz G. Five-minute treatments with fluorouracil, fluoridine, and mitomycin have long-term effects on human Tenon's capsule fibroblasts. *Arch Ophthalmol* 1992;110:1150-4.
- 31 Wilson SE, He YG, Weng J, Li Q, McDowall AW, Vital M, et al. Epithelial injury induces keratocyte apoptosis: hypothesised role for the interleukin 1 system in the modulation of corneal tissue organisation and wound healing. *Exp Eye Res* 1996;62:325-8.
- 32 Wilson, SE. Molecular cell biology for the refractive corneal surgeon: programmed cell death and wound healing. *J Refract Surg* 1997;13:171-6.
- 33 Dimri, G, Lee, X, Basile G, Acosta M, Scott G, Roskelley C, et al. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci* 1995;92:9362-7.
- 34 Gilchrist BA. Relationship between actinic damage and chronologic aging in keratinocyte cultures in human skin. *J Invest Dermatol* 1983;81:184s-9s.
- 35 Salla S, Redbrake C, Franz A, Reim M. Changes of human donor corneas preserved for longer than 4 weeks. *Vision Res* 1996;36:81.
- 36 Blake DA, Yu H, Young DL, Caldwell DR. Matrix stimulates the proliferation of human corneal endothelial cells in culture. *Invest Ophthalmol Vis Sci* 1997;38:1119-29.
- 37 Hoppenreijts VPT, Pels E, Vrensen GFJM, Treffers WF. Effects of platelet derived growth factor on endothelial wound healing of human corneas. *Invest Ophthalmol Vis Sci* 1994;35:150-61.
- 38 Hayashi K, Masumoto M, Fujino S, Hayashi F. Changes in corneal astigmatism with age. *Nippon Ganka Gakkai Zasshi* 1993;97:1193-6.
- 39 Hayashi K, Hayashi H, Hayashi F. Topographic analysis of the changes in corneal shape due to ageing. *Cornea* 1995;14:527-32.
- 40 Van den Berg TJ, Tan KE. Light transmittance of the human cornea from 320 to 700 nm for different ages. *Vision Res* 1994;34:1453-6.
- 41 Malik NS, Moss SJ, Ahmed N, Furth AJ, Wall RS, Meek KM. Ageing of the human corneal stroma: structural and biochemical changes. *Biochim Biophys Acta* 1992;1138:222-8.
- 42 Malik NS, Meek KM. Vitamins and analgesics in the prevention of collagen ageing. *Age Ageing* 1996;25:279-84.
- 43 Hobden JA, Masinick SA, Barrett RP, Hazlett LD. Aged mice fail to upregulate ICAM-1 after *Pseudomonas aeruginosa* corneal infection. *Invest Ophthalmol Vis Sci* 1995;36:1107-14.
- 44 Hazlett LD, Kreindler FB, Berk RS, Barrett R. Aging alters the phagocytic capability of inflammatory cells induced into cornea. *Curr Eye Res* 1990;9:129-38.
- 45 Lass JH, Greiner JV, Merchant TE, Glonek T. The effects of age on phosphatic metabolites of the human cornea. *Cornea* 1995;14:89-94.
- 46 Nzekwe EU, Maurice DM. The effect of age on the penetration of fluorescein into the human eye. *J Ocular Pharm* 1994;10:521-3.
- 47 Chang SW, Hu FR. Changes in corneal autofluorescence and corneal epithelial barrier function with aging. *Cornea* 1993;12:493-9.
- 48 Trinkaus-Randall V, Tong M, Thomas P, Cornell-Bell A. Confocal imaging of the alpha 6 and beta 4 integrin subunits in the human cornea with aging. *Invest Ophthalmol Vis Sci* 1993;34:3103-9.
- 49 Hubbard K, Ozer, HL. Senescence and immortalisation of human cells. In: Studzinski GP, ed. *Cell growth and apoptosis: a practical approach*. Oxford: IRL Press, 1995:229-49.
- 50 Gray, MD, Norwood TH. Cellular aging in vitro. *Rev Clin Gerontol* 1995;5:369-81.
- 51 Zeng G, Millis AJT. Differential regulation of collagenase and stromelysin mRNA in late passage cultures of human fibroblasts. *Exp Cell Res* 1996;222:150-6.
- 52 Millis AJ, Hoyle M, McCue HM, Martini H. Differential expression of metalloproteinase and tissue inhibitor of metalloproteinase genes in aged human fibroblasts. *Exp Cell Res* 1992;201:373-9.
- 53 Millis AT, Sottile J, Hoyle M, Mann DM, Diemer V. Collagenase production by early and late passage cultures of human fibroblasts. *Exp Gerontol* 1989;24:559-75.
- 54 Sorrentino JA, Millis AJT. Structural comparisons of fibronectin isolated from early and late passage cells. *Mech Age Dev* 1984;28:83-97.
- 55 Schneider EL, Mitsui Y. The relationship between in vitro cellular aging and in vivo human age. *Proc Natl Acad Sci* 1976;73:3584-8.
- 56 Bell E, Ivarsson B, Merrill C. Production of tissue-like structure by contraction of collagen lattices by human fibroblasts of different proliferative potential in vitro. *Proc Natl Acad Sci* 1979;76:1274-8.
- 57 Poot M, Visser WJ, Verkerk A, Jonkind JF. Autofluorescence of human skin fibroblasts during growth inhibition and in vitro ageing. *Gerontology* 1985;31:158-65.
- 58 Venable ME, Lee JY, Smyth MJ, Bielawska A, Obeid LM. Role of ceramide in cell senescence. *J Biol Chem* 1995;270:30701-8.
- 59 Macieira-Coelho A. Changes in membrane properties associated with cellular aging. *Int Rev Cytol* 1983;83:183-220.
- 60 Liu S, Thweatt R, Lumpkin CK, Goldstein S. Suppression of calcium dependent membrane currents in human fibroblasts by replicative senescence and forced expression of a gene sequence encoding a putative calcium-binding protein. *Proc Natl Acad Sci* 1994;91:2186-90.
- 61 Moller-Pederson T. A comparative study of human corneal keratocyte and endothelial cell density during aging. *Cornea* 1997;16:333-8.
- 62 Kanai A, Kaufman HE. Electron microscopic studies of corneal stroma: aging changes of collagen fibers. *Ann Ophthalmol* 1973;5:285-92.
- 63 Herd RM, Faragher RGA, Shall S, Hunter JAA. Werner's syndrome: a review of the clinical and pathological features and pathogenesis. *Eur J Dermatol* 1993;3:425-32.
- 64 Jonas JB, Rupprecht KW, Schmitz-Valckenberg P, Brambring D, Platt D, Gebhart E, et al. Ophthalmic surgical complications in Werner's syndrome: report of 18 eyes of nine patients. *Ophthalmol* 1987;18:760-4.
- 65 Dutt S, Steinert RF, Raizman MB, Puliafito CA. One year results of excimer laser photorefractive keratectomy for low to moderate myopia. *Arch Ophthalmol* 1994;112:1427-36.
- 66 Waring GO 3d, Lynn MJ, Nizam A, Kutner MH, Cowden JW, Culbertson W, et al. Results of the Prospective Evaluation of Radial Keratotomy (PERK) study five years after surgery: the PERK study group. *Ophthalmology* 1991;98:1164-76.
- 67 Chatterjee A, Shah S, Doyle SJ. Effect of age on final refractive outcome for 2342 patients following refractive keratectomy. *Invest Ophthalmol Vis Sci* 1996;37:S57.
- 68 Blatt HL. Endothelial cell density in relation to morphology. *Invest Ophthalmol Vis Sci* 1979;18:856-9.
- 69 Bourne WM, Nelson LR, Hodge DO. Central corneal endothelial changes over a ten year period. *Invest Ophthalmol Vis Sci* 1997;38:779-82.
- 70 Laule A, Cable MK, Hoffman CE, Hanna C. Endothelial cell population changes of human cornea during life. *Arch Ophthalmol* 1978;96:2031-5.
- 71 Murphy C, Alvarado J, Juster R, Maglio M. Prenatal and postnatal cellularity of the human corneal endothelium: a quantitative histologic study. *Invest Ophthalmol Vis Sci* 1984;25:312-22.
- 72 Laing RA, Sandstrom MM, Berrosipi AR, Leibowitz HM. Changes in the corneal endothelial cell function as a function of age. *Exp Eye Res* 1976;22:587-94.
- 73 Murphy C, Alvarado J, Juster R, Maglio M. Prenatal and postnatal cellularity of the human corneal endothelium. *Invest Ophthalmol Vis Sci* 1984;25:312-22.
- 74 Baroody RA, Bito LZ, Deroousseau CJ, Kaufman P. Ocular development of ageing. I Corneal endothelial changes in cats and in free-ranging and caged rhesus monkeys. *Exp Eye Res* 1987;45:607-22.
- 75 Fitch KL, Nadakavukaren MJ, Richardson A. Age-related changes in the corneal endothelium in the rat. *Exp Gerontol* 1982;17:179-83.
- 76 MacCallum DK, Bahn CF, Lillie JH, Meyer RF, Martonyi CL. Evidence for corneal endothelial cell hypertrophy during postnatal growth of the cat cornea. *Invest Ophthalmol Vis Sci* 1983;24:247-50.
- 77 Gwin RM, Lerner I, Warren JK, Gurn G. Decrease in canine corneal endothelial cell density and increase in corneal thickness as functions of age. *Invest Ophthalmol Vis Sci* 1982;22:267-71.
- 78 Oh JO. Changes with age in the cornea of normal rabbits. *Acta Ophthalmol* 1963;41:568-73.
- 79 Green K. Free radicals and ageing of anterior segment tissues of the eye: a hypothesis. *Ophthalmic Res* 1995;27(Suppl):143-9.
- 80 O'Neal MR, Polse KA. Decreased endothelial pump function with aging. *Invest Ophthalmol Vis Sci* 1986;27:457-63.
- 81 Polse KA, Brand R, Mandell R, Vastine D, Demartini D, Flom R, et al. Age differences in corneal hydration control. *Invest Ophthalmol Vis Sci* 1989;30:392-9.
- 82 Ehlers N. Graft thickness after penetrating keratoplasty. *Acta Ophthalmol* 1974;52:893-903.
- 83 Kimura C, Tanishima T. Thickness of the corneal graft after penetrating keratoplasty. *Jpn J Ophthalmol* 1975;19:348-53.
- 84 Staaz WD, Van-Horn DL. The effects of aging and inflammation on corneal endothelial wound healing in rabbits. *Invest Ophthalmol Vis Sci* 1980;19:983-6.