Postscript

LETTERS

If you have a burning desire to respond to a paper published in *BJO*, why not make use of our "rapid response" option?

Log onto our website (www.bjophthalmol.com), find the paper that interests you, and send your response via email by clicking on the "eLetters" option in the box at the top right hand corner.

Providing it isn't libellous or obscene, it will be posted within seven days. You can retrieve it by clicking on "read eLetters" on our homepage.

The editors will decide as before whether to also publish it in a future paper issue.

Orbital varices and orbital wall defects

Orbital varices are a vascular hamartoma typified by a plexus of low pressure, low flow, thin walled and distensible vessels that intermingle with the normal orbital vessels.¹⁻⁴ If freely communicating with the orbital circulation, engorgement of varices can occur by increasing venous pressure through the Valsalva manoeuvre,⁵ bending posture,⁶ coughing or straining and these, in turn, lead to the clinical characteristics of variable proptosis, intermittent pain, and orbital haemorrhage.^{7,8}

Observation is usually warranted for small lesions, but surgical intervention may be necessary in advanced cases: indications for surgical intervention include non-resolving episodes of thrombosis, severe disfiguring proptosis or displacement of the globe, and optic nerve compression.¹⁻³ Surgery can be extremely difficult, as varices are very friable and intimately intermixed with normal orbital structures; there is also a significant risk of visual loss as a result of haemorrhage or optic nerve damage, the latter being generally caused by vascular compromise.^{9,10} The association of orbital venous anomalies with orbital wall defects provides a further source of surgical difficulty because of the close proximity of intracranial structures and the continuity with extraorbital or intracranial venous anomalies.

Case series

The orbital database, at Moorfields Eye Hospital, was used to identify patients with a clinical diagnosis of low pressure orbital varices and their orbital imaging (computed tomography and/or magnetic resonance image) was reviewed. Images were examined for evidence of orbital expansion, osseous defects of the orbit, nose or sinuses, and anomalies of the frontal lobes. Patients who had either orbital or intracranial surgery before the date of imaging were excluded from the investigation.

The clinical diagnosis of orbital varices was identified in 310 patients, and imaging was available for 223 patients (72%). Six patients with previous orbital or intracranial surgery were excluded and nine cases (4%) had

Figure 1 (A) (Patient 1) Extensive left orbital varices (white arrows) causing orbital expansion, globe displacement, and "pitting" of the orbital roof and lamina papyracea (black arrows). (B) (Patient 5) Bilateral orbital varices associated with multiple defects, rather than pitting, of the orbital walls. (C) (Patient 6) Right orbital varices, with phleboliths, extending through the orbital apex into the middle cranial fossa (white arrows) and (D) associated with intracranial bone "pitting" and "defects" (black arrows). (E) Coronal and (F) axial CT scans of patient 2 with superonasal varices (white arrows) of right orbit in 1981. (G-J) Repeat coronal and axial CT scans in 2001 show significant enlargement of the bone defect with complete loss of mineralisation, and expansion of the frontal lobe meninges into the orbital wall defect (black arrow).

associated anomalies of the neighbouring orbital walls (table 1).

Four cases (patients 1-4) were associated with "pitting" of the orbital wall secondary to orbital varices (fig 1A). Another three cases (patients 6-8) were associated with enlarged

superior orbital fissure and two cases (patients 5 and 9) with multiple orbital roof "defects" (fig 1B). Orbital varices were present up to the dural space in two cases (patients 4 and 5), and involved the frontal lobe parenchyma in one case (patient 6; fig 1C, D).

Table 1 Characteristics of nine patients with orbital wall defects in association with orbital varices

No	Side	Age (years) at referral	Sex	Main location of orbital varix	Expansion of orbit	Absent walls	Ethmoid	Cribriform	Frontal
1	Left	6	M	Medial and extensive superomedial	Present	Small roof defect	Pitted bone and smaller ethmoid	L-low R-normal	Dips low at cribriform
2	Right	21	F	Extraconal-medial	Present	Tiny thin area SNQ	Pitted bone and smaller ethmoid	R-low L-mild	Low frontal lobe over cribriform
3	Left	62	M	Superomedial	Present	Pitted roof and small defects of veins	Compressed	Normal	Hint of varix but otherwise normal
4	Right	58	F	Panorbit intraconal and extraconal	Present	Post superior wall and pitted bone	Normal	Normal	Varices up to frontal lobe and intraconal
5	Right	47	M	Panorbit intraconal and extraconal	Absent	Posterior orbital roof	Normal	Normal	Varices up to dural space
6	Right	14	F	Posterior intraconal, superior extraconal	Present	Enlarged SOF	Normal	Normal	Varices into frontal lobe
7	Left	40	F	Posterior intraconal	Present	Enlarged SOF and small lateral wall	Slightly smaller	Unknown	Normal
8	Left	37	F	Posterior intraconal and extraconal	Present	Very enlarged SOF, patchy SNQ defects posteriorly	Normal	Normal	Normal
9	Left	66	M	Extraconal-superior (large)	Present	Posterior orbital roof	Slightly smaller	Normal	Normal

SNQ = superonasal quadrant; SOF = superior orbital fissure.

One patient (case 2) had thinning of the superonasal quadrant of the orbital wall, nasal orbital wall pitting, and a low ipsilateral cribriform plate, when first seen at age 21 in 1981 (fig 1E, F). On repeat imaging 20 years later (2001), this patient was noted to have developed proptosis, a defect in the superonasal wall of the orbit, and a new mid-line nasal encephalocele (fig 1I, J).

Comment

Fine cut (3 mm) orbital CT scans easily delineate varices and diagnostic phleboliths, which occur from thrombus formation,⁷ and provide an excellent natural contrast between brain, bone, and varix. The typical findings for varices include an ill defined multiloculated mass, with some patchy contrast enhancement, in communication with the neighbouring orbital circulation⁴⁻⁷; diffuse expansion of the orbital walls is well recognised in some cases, especially in childhood lesions.

Several factors may have biased the study population: many are symptomatic patients, having been referred from other ophthalmic units in consideration for surgical intervention. The apparent incidence of orbital wall defects (4%) in our series may, therefore, be a slight overestimate. In a minority of patients, orbital varices may be associated with orbital wall defects, and such defects may, eventually, lead to an encephalocele formation. Clinicians should be aware of these, apparently unreported, associations before embarking on surgical intervention for orbital varices.

N Islam, K Mireskandari, G E Rose
Moorfields Eye Hospital, London EC1V 2PD, UK

Correspondence to: Mr G E Rose, Orbital Service, Moorfields Eye Hospital, City Road, London EC1V 2PD, UK;

doi: 10.1136/bjo.2003.024547

Accepted for publication 10 July 2003

References

1 Rootman J. *Diseases of the orbit. A multidisciplinary approach*, 2nd ed. Philadelphia: Lippincott Williams and Wilkins, 2002:517-22.

2 Beyer R, Levine MR, Sternberg I. Orbital varices: a surgical approach. *Ophthalm Plast Reconstr Surg* 1985;1:205-10.
3 Wright JE, Sullivan TJ, Garner A, et al. Orbital venous anomalies. *Ophthalmology* 1997;104:905-13.
4 Rubin PAD, Remulla HD. Orbital venous anomalies demonstrated by spiral computed tomography. *Ophthalmology* 1997;104:1463-70.
5 Shields JA, Dolinskas C, Augsburger JJ, et al. Demonstration of orbital varix with computed tomography and Valsalva manoeuvre. *Am J Ophthalmol* 1984;97:108-10.
6 Cohen JA, Char DH, Norman D. Bilateral orbital varices associated with habitual bending. *Arch Ophthalmol* 1995;113:1360-2.
7 Bullock JD, Goldberg SH, Connelly PJ. Orbital varix thrombosis. *Ophthalmology* 1990;97:251-6.
8 Foroosan R, Shields CL, Shields JA, et al. Congenital orbital varices causing extreme neonatal proptosis. *Am J Ophthalmol* 2000;129:693-4.
9 Harris GJ, Sakol PJ, Bonavolonta G, et al. An analysis of thirty cases of orbital lymphangioma. Pathophysiologic considerations and management recommendations. *Ophthalmology* 1990;97:1583-92.
10 Weill A, Cognard C, Castaings L, et al. Embolization of an orbital varix after surgical exposure. *Am J Neuroradiol* 1998;19:921-3.

Exenteration of invasive conjunctival squamous cell carcinoma

Ocular surface squamous neoplasia (OSSN) includes conjunctival intraepithelial neoplasia with dysplasia, carcinoma in situ and conjunctival squamous cell carcinoma (SCC). Beside ultraviolet B irradiation identified as an risk factor, OSSN is associated with human papillomavirus type 16 and 18 (HPV-16, HPV-18).¹⁻⁵ The exact role and possible prognostic value of p53 overexpression is unclear and little is known about its expression during the development of conjunctival SCC.

Case reports

Patient 1

A 75 year old man was referred with a 10 year history of a conjunctival mass of the left eye with visual acuity of hand movement.

Previous biopsies had revealed conjunctival dysplasia. On examination, the tumour of the ocular and tarsal conjunctiva of the lower lid covered the entire corneal surface (fig 1A).

Patient 2

A 90 year old patient presented with a 2 year history of an extensive conjunctival papillomatous tumour of the left eye covering three quarters of the cornea with visual acuity of light perception. A full thickness biopsy was performed.

Both patients underwent orbital exenteration including removal of the eyelids. Histopathologically the focal invasive, completely removed tumour of patient 1 grew in a papillomatous manner. The tumour cells of the conjunctival neoplasm showed strongly enlarged nuclei with prominent nucleoli, and formed cohesive units with intercellular bridges (fig 1B).

The exophytic tumour of patient 2 was predominantly intraepithelial with foci of subepithelial invasion. Focal tumour anaplasia was observed in the otherwise moderately differentiated tumour with squamous cell differentiation.

Immunostaining of both specimens revealed strong p53 (monoclonal mouse-anti-human p53-protein DO-7, Dako) over-expression (>26% of epithelium cells) and low expression of p21 (<6% of epithelium cells) of the invasive region of the tumour indicating an inactivating p53 mutation (fig 1C). While in patient 1 expression for p53 was found in all epithelial layers, in patient 2 it was expressed suprabasally. In contrast, both p53 and p21 showed moderate reactivity in the dysplastic region up to the middle layer of the tumour (fig 1D). In the apical layer epithelium cells were occasionally p21 positive.

Immunostaining for HPV (HPV screening antibody, Virofem Diagnostica, Germany) was positive in patient 1.

Comment

The high recurrence rate of OSSN of 9-64% after resection seems to depend on the histopathological grade and status of surgical margins.⁵ HPV-16 and HPV-18 are considered

Figure 1 Patient 1. (A) Extensive papillomatous tumour, subtotally covering the corneal surface of the left eye. Nodular thickening of the lower eyelid indicates eyelid involvement. (B) Histological appearance. Papillomatous pattern of the large epithelial lesion with focal invasion above the cornea. Subepithelially, inflammatory cells and some dilated vessels (haematoxylin and eosin, original magnification, $\times 2.5$). (C) p53, showing strong diffuse reactivity in invasive region indicating an inactivating p53 mutation (original magnification, $\times 10$). (D) p53, showing moderate expression predominantly in the suprabasal layers in dysplastic conjunctiva of the same specimen (original magnification, $\times 10$).

to be possible cofactors involved in initiation and early progression of OSSN.^{1-4,7} Though both of the presented tumours were clinically papillomatous, immunostaining for HPV was positive only in patient 1.

The tumour suppressor gene p53 has been found to be inactivated in over 50% of human cancers. In OSSN, overexpression of p53 has been previously reported in some SCC of conjunctiva.^{6,7} In the SCC of our patients, p53 overexpression indicating inactivating p53 mutations were observed only in the invasive part of the tumour, but not in the carcinoma in situ. While Dushku and coworkers assumed that p53 mutations could be an early event in tumour development consistent with ultraviolet radiation, our findings clearly indicate that mutations of p53 are a late event that occur with disease progression, as observed with other solid tumours.^{7,8} Karcioğlu and associates found a correlation between p53 overexpression and unfavourable clinical course.⁹ In contrast, Aoki and colleagues found no expression for p53 in SCC but positive staining in dysplasia.¹⁰

Our results of two exenterated advanced stages of SCC emphasise the necessity to remove dysplastic OSSN completely to prevent progression to invasive carcinomas. Identification of inactivating p53 mutations may indicate an increased risk for invasiveness. Therefore immunohistochemical analysis of biopsy specimen may help in the management of these tumours.

R Guthoff, W E Lieb

Department of Ophthalmology, Julius Maximilians University Würzburg, Germany

P Ströbel, A Zettl

Institute of Pathology, Julius Maximilians University Würzburg, Germany

Correspondence to: Dr R Guthoff, Universitäts-Augenklinik, Josef-Schneider-Strasse 11, 97080 Würzburg, Germany; r_guthoff@yahoo.de

doi: 10.1136/bjo.2003.038588

Accepted for publication 7 December 2003

References

- McDonnell JM, Mayr AJ, Martin WJ. DNA of human papillomavirus type 16 in dysplastic and malignant lesions of the conjunctiva and cornea. *N Engl J Med* 1989;**320**:1442-6.
- McDonnell JM, McDonnell PJ, Sun YY. Human papillomavirus DNA in tissues and ocular surface swabs of patients with conjunctival epithelial neoplasia. *Invest Ophthalmol Vis Sci* 1992;**33**:184-9.
- Nakamura Y, Mashima Y, Kameyama K, et al. Detection of human papillomavirus infection in squamous tumours of the conjunctiva and lacrimal sac by immunohistochemistry, in situ hybridisation, and polymerase chain reaction. *Br J Ophthalmol* 1997;**81**:308-13.
- Karcioğlu ZA, Issa TM. Human papilloma virus in neoplastic and non-neoplastic conditions of the external eye. *Br J Ophthalmol* 1997;**81**:595-8.
- Lee GA, Hirst LW. Ocular surface squamous neoplasia. (Review) *Surv Ophthalmol* 1995;**39**:429-50.
- Toth J, Karcioğlu ZA, Moshfeghi AA, et al. The relationship between human papillomavirus and p53 gene in conjunctival squamous cell carcinoma. *Cornea* 2000;**19**:159-62.
- Dushku N, Hatcher SL, Albert DM, et al. p53 expression and relation to human papillomavirus infection in pingueculae, pterygia, and limbal tumors. *Arch Ophthalmol* 1999;**117**:1593-9.
- Gryfe R, Swallow C, Bapat B, et al. Molecular biology of colorectal cancer. *Curr Probl Cancer* 1997;**21**:233-300.
- Karcioğlu ZA, Toth J. Relation between p53 overexpression and clinical behavior of ocular/orbital invasion of conjunctival squamous cell carcinoma. *Ophthalmol Plast Reconstr Surg* 2000;**16**:443-9.
- Aoki S, Kubo E, Nakamura S, et al. Possible prognostic markers in conjunctival dysplasia and squamous cell carcinoma. *Jpn J Ophthalmol* 1998;**42**:256-61.

Familial pseudotumoral sclerochoroidal calcification associated with chondrocalcinosis

Sclerochoroidal calcification is the deposition of calcium at the level of the sclera and choroid. Two entities have been described: metastatic calcifications resulting from deposition of calcium in normal tissues caused by phosphocalcic metabolism abnormality such as primary and secondary hyperparathyroidism,¹ pseudohypoparathyroidism, hypervitaminose D, vitamin D intoxication, hypophosphataemia, sarcoidosis, Bartter syndrome, and Gitelman syndrome²³; and dystrophic calcifications caused by secondary deposition of calcium in abnormal tissues despite normal serum levels of calcium and phosphate.

Sclerochoroidal calcifications can also be idiopathic.⁴ We describe the first case of hereditary form of sclerochoroidal calcifications associated with familial articular chondrocalcinosis.

Case report

A 69 year old man was admitted to the department of ophthalmology in November 1999 with gradual deterioration of vision in both eyes. He had a medical history of articular chondrocalcinosis. His father, brother, and son were treated for the same disease.

On examination, best corrected visual acuity was 20/120 in the right eye and finger counting in the left eye. Slit lamp examination and ocular tension were normal. The funduscopy revealed multiple bilateral pseudotumoral white choroidal masses in both eyes (fig 1). Ultrasound examination of the eyes confirmed the calcific nature of the lesions.

On fluorescein angiography in March 1979 the pseudotumoral lesions were smaller and did not involve the macular area in left eye (fig 1, left).

He had an extensive metabolic evaluation in the Broussais department of physiology that was normal. The systemic diagnosis was familial articular chondrocalcinosis.

We decided to examine the whole family to search for ophthalmic abnormalities linked with familial chondrocalcinosis.

The 74 year old brother and the 40 year old son also suffered from chronic articular chondrocalcinosis. Their best visual acuity was 20/20 in both eyes.

Figure 1 Fluorescein angiography of the 69 year old patient in 1979 (left) and 2000 (right) showing evolution of the sclerochoroidal calcifications with time.

The brother's funduscopy revealed multiple bilateral, extrafoveal pseudotumoral white choroidal masses (fig 2, top). The son's funduscopy revealed plaque-like and only slightly elevated lesions seen in the mid-periphery (fig 2, bottom). Ultrasonograms confirmed the calcific nature of these lesions.

Comment

In 1997, Shields *et al*⁵ described a case of sclerochoroidal calcifications in a normocalcaemic patient who had chondrocalcinosis.

We first describe a familial case of sclerochoroidal calcifications associated with calcium pyrophosphate dihydrate (CPPD). In this family, autosomal dominant inheritance is highly likely because there are affected individuals in each generation, there is male to male transmission, and every affected member has an affected parent.

Inheritance in sclerochoroidal calcifications has never been described; however, hereditary forms of chondrocalcinosis have already been described.

In our report, a patient had a 24 year follow up showing a progressive involvement of the macular area, suggesting a growth of the calcifications. Two types of calcifications have been described previously, the plaque-like and the pseudotumoral type.

To our knowledge, it has never been determined if the plaque-like lesions evolve into tumour-like lesions. In 1992, Schachat and associates⁶ reported 10 cases with follow up ranging from 7 months to 10 years, for whom no change in the lesion was seen. This is the first observation with 24 years of follow up suggesting a possible evolution of plaque-like lesions to pseudotumoral lesions.

We suggest that every patient affected by familial chondrocalcinosis should have an ophthalmic examination to detect sclerochoroidal calcification. These lesions seem to be evolving in time with possible involvement of the macula. Choroidal neovascularisation is also a vision threatening complication of sclerochoroidal calcifications.⁷ Our case suggests the need to perform ophthalmological

examination in patients and family members of patients affected by chondrocalcinosis.

S Boutboul, T Bourcier, J-P Heligon, P Houiller, M Ullern, M Abitbol, V Borderie, L Laroche
CHNO des XV-XX, 28, Rue de Charenton 75571 Paris
Cedex 12 Paris, France

Correspondence to: Dr Sandrine Boutboul, CHNO des XV-XX, 28, Rue de Charenton 75571 Paris Cedex 12 Paris, France; skowronek@free.fr

doi: 10.1136/bjo.2003.039925

Accepted for publication 15 December 2003

References

- 1 Suzuki JJ, Takeda M, Sekine N, *et al*. Bilateral metastatic sclerochoroidal calcification in a patient with hyperparathyroidism. *Ophthalmologica* 1992;**205**:10-14.
- 2 Bourcier T, Blain P, Massin P, *et al*. Sclerochoroidal calcification associated with Gitelman syndrome. *Am J Ophthalmol* 1999;**128**:767-8.
- 3 Vezzoli G, Soldati L, Jansen A, *et al*. Choroidal calcifications in patients with Gitelman's syndrome. *Am J Kidney Dis* 2000;**6**:855.
- 4 Lim JJ, Goldberg MF. Idiopathic sclerochoroidal calcification. *Arch Ophthalmol* 1989;**107**:1122-3.
- 5 Shields JA. Sclerochoroidal calcification in calcium pyrophosphate dihydrate deposition disease (pseudogout). *Arch Ophthalmol* 1997;**115**:1077-9.
- 6 Schachat AP, Robertson DM, Mieler WF, *et al*. Sclerochoroidal calcification. *Arch Ophthalmol* 1992;**110**:196-9.
- 7 Cohen SY, Guyot-Sionnest M, Puech M. Choroidal neovascularization as a late complication of hyperparathyroidism. *Am J Ophthalmol* 1998;**126**:320-2.

Whole body PET/CT imaging for detection of metastatic choroidal melanoma

Metastatic choroidal melanoma typically presents in the liver. Therefore, liver enzyme assays are the most common haematological evaluation performed after treatment.¹

In 1985, The Collaborative Ocular Melanoma Study required periodic medical evaluations including a physical examination, liver functions studies, a complete blood count, and a chest x ray. If liver enzymes exceeded 1.5 times normal, computed tomography (CT) of the abdomen was required. If low attenuation hepatic nodules suggested metastatic disease, fine needle aspiration biopsy of the liver tumours provided cytopathological confirmation.¹

Positron emission tomography (PET) is a molecular imaging technique that uses radiolabelled molecules to image metabolic activity in vivo.²⁻³ When whole body PET was combined with computed radiographic tomography (CT), PET/CT put anatomy and function on the same page making practical a new era of physiological imaging.²⁻⁶

This study examines the ability of positron emission tomography combined with computed tomography (PET/CT) to allow for detection of previously occult metastatic melanoma.

Case report

A 77 year old woman presented with a 15.4×15 mm width and 13.2 mm high collar button shaped choroidal melanoma with a large secondary retinal detachment in her right eye. Her preoperative medical evaluation

Figure 2 Funduscopy of the patient's brother (top) and son (bottom) revealing tumour-like and plaque-like calcifications.

Table 1 Patient characteristics						
Physical	Examination	Blood examination	X ray	CT scan abdomen with contrast	Whole body PET/CT	
					CT	PET
General	Icterus					
Subcutaneous tissue	2 nodules in the anterior abdominal wall			No lesions noted	1.4 cm subcutaneous nodule in the anterior abdominal wall	Hypermetabolic focus in the subcutaneous tissue of the anterior abdominal wall
Nodes	None			No lesions noted	Enlarged para aortic lymph nodes on the left side	Hypermetabolic focus in the upper abdomen
Lungs	No abnormalities noted		Pleural effusion of right lung base. No nodules	Pleural effusion right lung base and calcified hilar nodes	Two small 3 mm nodules in the right upper lobe. Right pleural effusion	No foci noted
Liver	Enlarged	High bilirubin, AST, ALT, alkaline phosphatase, GGT		Enlarged. 2 areas of low attenuation in the anterior aspect of the right lobe	Low attenuation lesion >20 cm in greatest diameter in the right lobe with calcification seen posteriorly. Numerous low attenuation lesions throughout both lobes of the liver	Enlarged. Large hypermetabolic focus in the right lobe with mass effect. Numerous hypermetabolic foci throughout remainder of liver.
Kidney				8 mm cyst midpole and 2 cm cyst upper pole of left kidney	Large right renal cyst. Additional smaller cysts	Photopenic defect due to large right renal cyst
Bone				No lesions noted	No lesions noted	Hypermetabolic foci in the right skull base, left scapula, left humerus, sternum, multiple bilateral ribs, thoracic and lumbosacral spine, pelvis, and bilateral femurs

(including CT imaging of the abdomen) proved negative. She was treated by enucleation.

Two years later a follow up medical evaluation revealed elevated liver function studies (table 1) and a chest x ray showed a

pleural effusion. CT of the abdomen with contrast revealed multiple low attenuation hepatic foci consistent with metastatic melanoma.

A PET/CT was requested. Fifty minutes after intravenous administration of 15.2 mCi of fluorodeoxyglucose, whole body PET/CT imaging revealed enlarged para-aortic lymph nodes and a subcutaneous nodule in the abdominal wall (fig 1). The CT portion of the PET/CT also revealed two 3 mm nodules in the upper lobe of the right lung (too small to be visualised by PET imaging). PET imaging was able to reveal multiple bony metastasis that were not seen on the CT portion (fig 1). Both CT and PET showed a large liver metastasis, but CT was better at defining tumour size. Since it is a physiological assay, PET also demonstrated the metabolic activity of the metastatic tumours (fig 1).

Comment

In this case, whole body PET/CT was found to be capable of uncovering metastases not seen with abdominal CT alone. This led us away from considering regional perfusion of the liver, hepatic resection, and towards systemic treatment.⁷⁻⁸ Therefore, when PET/CT identifies extrahepatic involvement, it can have a significant impact on the management of patients with metastatic choroidal melanoma.

PET/CT could also be used for initial staging. Early detection of occult metastases

Figure 1 On the left, the CT demonstrates the anatomy; on the right the PET shows areas of hypermetabolism (glucose uptake); in the middle the two images are fused. PET/CT revealed enlarged para-aortic lymph nodes and a subcutaneous nodule in the anterior abdominal wall. The PET imaging portion of the PET/CT was able to reveal multiple bony metastases that were not seen on the CT portion of the examination. Both CT and PET showed a large liver metastasis.

offers the potential to avoid ineffective and expensive enucleations, radioactive plaques, proton irradiation, eye wall resections, or laser treatment.⁹⁻¹⁰ Local therapies would be abandoned in favour of systemic treatments.

Another issue related to PET/CT is cost. Up to five times more than CT of the abdomen, PET/CT is only covered (Medicare) for melanoma staging/restaging when the stage of the cancer remains in doubt after completion of conventional imaging (or if the clinical management would differ depending on the PET findings). Since PET/CT revealed extrahepatic foci in this case, it changed our clinical approach. There is little doubt about the improved ability of PET/CT to detect lesions; the real issue is cost and if the results will change outcomes.

This study goes one step further than CT, MRI, or PET alone. By combining whole body PET and CT, this examination joins anatomy and function in one examination (fig 1). The relative efficacy of PET/CT to locate metastases should be evaluated within the framework of a prospective study.

P T Finger, M Kurli

The New York Eye Cancer Center, New York, USA

P T Finger, M Kurli, P Wesley, L Tena

The New York Eye and Ear Infirmary, New York, USA

P T Finger, L Tena

St Vincent's Comprehensive Cancer Center, New York, USA

P T Finger, K R Kerr

Beth Israel Medical Center

P T Finger, A Pavlick

New York University School of Medicine, New York, USA

Correspondence to: Paul T Finger, MD, The New York Eye Cancer Center, 115 East 61st Street, New York, NY 10021, USA; pfinger@eyecancer.com

doi: 10.1136/bjo.2003.039289

Accepted for publication 1 January 2004

This work was supported by The EyeCare Foundation and Research to Prevent Blindness, New York, USA.

References

- Diener-West M, Hawkins BS, Markowitz JA, *et al.* A review of mortality from choroidal melanoma. II A meta-analysis of 5-year mortality rates following enucleation, 1966 through 1988. *Arch Ophthalmol* 1992;110:245-50.
- Townsend DW, Cherry SR. Combined anatomy and function: the path to true image fusion. *Eur Radiol* 2001;11:1968-74.
- Mijnhout GS, Hoekstra OS, van Tulder MW, *et al.* Systematic review of the diagnostic accuracy of 18F-fluorodeoxyglucose positron emission tomography in melanoma patients. *Cancer* 2001;91:1530-42.
- Stas M, Stroobants S, Dupont P, *et al.* 18-FDG PET scan in the staging of recurrent melanoma: additional value and therapeutic impact. *Melanoma Res* 2002;12:479-90.
- Swetter SM, Carroll LA, Johnson DL, *et al.* Positron emission tomography is superior to computed tomography for metastatic detection in melanoma patients. *Ann Surg Oncol* 2002;9:646-53.
- Finger PT, Czechonska G, D'Arienzo P, *et al.* Positron emission tomography of choroidal melanoma. *Probl Med Nucl* 1996;10:235-9.
- Mavligit GM, Charnsangavej C, Carrasco CH, *et al.* Regression of ocular melanoma metastatic to the liver after hepatic arterial chemoembolization with cisplatin and polyvinyl sponge. *JAMA* 1988;260:974-6.
- Albert DM, Niffenegger AS, Willson JK. Treatment of metastatic uveal melanoma: review and recommendations. *Surv Ophthalmol* 1992;36:429-38.
- Moshfeghi DM, Moshfeghi AA, Finger PT. Enucleation. *Surv Ophthalmol* 2000;44:277-301.
- Finger PT. Radiation therapy for choroidal melanoma. *Surv Ophthalmol* 1997;42:215-32.

Trans-Tenon's retrobulbar triamcinolone infusion for small choroidal neovascularisation

Intravitreal and sub-Tenon's corticosteroids are being evaluated for the treatment of choroidal neovascularisation (CNV).¹⁻³ We reported on the efficacy of triamcinolone acetonide administered as a trans-Tenon's retrobulbar infusion (triamcinolone infusion) in reducing inflammation in uveitic eyes.⁴ Here we evaluated the same treatment in eyes with small subfoveal CNV.

Case reports

Triamcinolone infusion was performed in 22 eyes of 22 patients with subfoveal CNV of greatest diameter less than or equal to 1 disc diameter (DD). The diagnoses were age related macular degeneration (AMD) in 14 eyes, idiopathic CNV in four eyes, polypoidal choroidal vasculopathy (PCV) in three eyes, and punctate inner choroidopathy (PIC) in one eye. One AMD eye had undergone ablative argon laser photocoagulation for CNV previously, but no other eyes had received previous treatment. The median post-triamcinolone infusion follow up was 7.5 months (range 4-27 months). Pretreatment fluorescein angiography (FA)

revealed predominantly classic CNV in 12 eyes and predominantly occult CNV in 10 eyes. Records were reviewed retrospectively and did not require institutional review board approval. Informed consent was obtained before each procedure.

The patient's eye was prepared with povidone-iodine and sterile drapes applied. After topical anaesthesia, conjunctiva and Tenon's capsule were incised in the infero-temporal quadrant. A 23 gauge curved blunt cannula approximately 2.1 cm in length (#HS-2764, Handaya Co, Ltd, Tokyo, Japan) was inserted to the hub into sub-Tenon's space and 20 mg (0.5 ml) triamcinolone acetonide (Bristol Pharmaceutical, KK, Tokyo, Japan) was infused. The wound was left unsutured and 0.5% levofloxacin was instilled topically three times a day for 1 week.

Onset of CNV fibrosis was observed in 14 eyes (64%) by 3 months post-triamcinolone infusion (fig 1). Rates of fibrosis were 50% (7/14 eyes) for AMD, 100% (4/4 eyes) for idiopathic CNV, 67% (2/3 eyes) for PCV, and 100% (1/1 eye) for PIC. Fibrosis did not correlate with CNV size or lesion composition. FA performed at 3 months showed decreased lesion leakage in 12 eyes (55%), no change in five eyes (23%), and increased leakage in five eyes (23%). Best corrected visual acuity (VA) at 3 months for all eyes improved by ≥ 0.2 logarithm of the minimum angle of resolution (logMAR) in four eyes (18%), remained unchanged in 13 eyes (59%), and worsened by ≥ 0.2 logMAR in five eyes (23%). The median decimal VA for all eyes was 0.30 before treatment (range 0.08-1.0) and 0.24 at 3 months after treatment (range 0.05-1.2). Of the 14 eyes with AMD, the VA at 3 months improved by ≥ 0.2 logMAR in one eye (7%), remained

Figure 1 Colour fundus photographs and fluorescein angiography late images of an eye in a 65 year old patient with age related macular degeneration and small subfoveal choroidal neovascularisation before (A, B; best corrected visual acuity 0.3) and 3 months after triamcinolone infusion (C, D; best corrected visual acuity 0.7).

unchanged in 10 eyes (71%), and worsened by ≥ 0.2 logMAR in three eyes (21%). In these AMD eyes, the median decimal VA was 0.30 before treatment (range 0.08–1.0) and 0.20 at 3 months after treatment (range 0.05–0.7). Complications such as intraocular pressure elevation, infection, or cataract progression were not noted in any eyes.

Comment

This interventional case series shows that trans-Tenon's retrobulbar infusion of triamcinolone acetonide resulted in lesion fibrosis in the majority of eyes with small CNV, efficacy being best for idiopathic CNV or CNV related to PIC. The mechanism of action of triamcinolone acetonide in inhibiting CNV growth probably involves several pathways. The effect of corticosteroids in inhibiting inflammatory cells that participate in the neovascular response probably has a prominent role.^{3,6} Triamcinolone acetonide has specifically been shown to inhibit basic fibroblast growth factor induced migration and tube formation of choroidal microvascular endothelial cells.⁷ Furthermore, triamcinolone acetonide inhibits choroidal neovascularisation induced by laser trauma in a rat model.⁸ Finally, triamcinolone acetonide may decrease vascular permeability, thereby decreasing influx of serum proteins that may contribute to an angiogenic micro-environment.⁹ Longer follow up and greater numbers of cases in a randomised clinical trial are needed to confirm these results.

A A Okada, T Wakabayashi, E Kojima, Y Asano, T Hida

Kyorin Eye Center, Kyorin University School of Medicine, 6-20-2 Shinkawa, Mitaka, Tokyo, Japan

Correspondence to: A A Okada, MD, Department of Ophthalmology, Kyorin University School of Medicine, 6-20-2, Shinkawa, Mitaka, Tokyo 181-8611 Japan; aokada@po.ijnet.or.jp

doi: 10.1136/bjo.2003.039719

Accepted for publication 6 January 2004

References

- Challa JK, Gillies MC, Penfold PL, *et al.* Exudative macular degeneration and intravitreal triamcinolone: 18 month follow-up. *Aust NZ J Ophthalmol* 1998;26:277–81.
- Danis RP, Ciulla TA, Pratt LM, *et al.* Intravitreal triamcinolone acetonide in exudative age-related macular degeneration. *Retina* 2000;20:244–50.
- Gillies MC, Simpson JM, Luo W, *et al.* A randomized clinical trial of a single dose of intravitreal triamcinolone acetonide for neovascular age-related macular degeneration: one year results. *Arch Ophthalmol* 2003;121:667–73.
- Okada AA, Wakabayashi T, Morimura Y, *et al.* Trans-Tenon's retrobulbar triamcinolone infusion for the treatment of uveitis. *Br J Ophthalmol* 2003;87:968–71.
- Proia A, Hiraoka A, McInnes J, *et al.* The effect of angiostatic steroids and B-cyclodextrin tetradecasulfate on corneal neovascularization in the rat. *Exp Eye Res* 1993;57:693–8.
- Oh H, Takagi H, Takagi C, *et al.* The potential angiogenic role of macrophages in the formation of choroidal neovascular membranes. *Invest Ophthalmol Vis Sci* 1999;40:1891–8.
- Wang Y, Friedrichs U, Eichler W, *et al.* Inhibitory effects of triamcinolone acetonide on bFGF-induced migration and tube formation in choroidal microvascular endothelial cells. *Graefes Arch Clin Exp Ophthalmol* 2002;240:42–8.
- Ciulla TA, Criswell MH, Danis RP, *et al.* Intravitreal triamcinolone acetonide inhibits

choroidal neovascularization in a laser-treated rat model. *Arch Ophthalmol* 2001;119:399–404.

- Carnuccio R, Di Rosa M, Guerrasio B, *et al.* Vasocortin: a novel glucocorticoid-induced anti-inflammatory protein. *Br J Pharmacol* 1987;90:443–5.

Retinoblastoma in a child conceived by in vitro fertilisation

As the number of infants born through in vitro fertilisation (IVF) grows, there is increased interest regarding the long term effects of IVF and other assisted reproduction techniques on such offspring. Recent reports have noted cancer in children born after IVF or fertility drug use (table 1).¹

In 2001, retinoblastoma was reported to occur in a child born through IVF in Israel.² Since then, an additional report documented four cases from the Netherlands.³ Here we add a sixth case and the first from the United States. Of these children four had unilateral retinoblastoma and two bilateral disease (table 2).

During 2002, a 16 month old child was referred to The New York Eye Cancer Center with no known family history of eye disease. She had a blind painful right eye with neovascular glaucoma. Intraocular pressures were 35 mm Hg in the right eye, and 14 in the left. Examination of the anterior chamber of the right eye revealed cells on the corneal endothelium and iris neovascularisation. While ophthalmoscopic examination of the right eye was not possible because of opaque media, ultrasonography revealed a densely calcified mass in the posterior pole (fig 1). Computed radiographic tomography demonstrated tumour calcification with no evidence of extrascleral or optic nerve extension.⁴ Post-enucleation histopathology confirmed the diagnosis of retinoblastoma.⁵ The parents did not approve genetic studies of the child.

Further history revealed that this child was born through IVF with a donor egg and the father's sperm. In order to carry the child, the postmenopausal mother received oestrogen and progesterone before and during gestation.

Comment

Several theories of IVF related carcinogenesis exist. Prenatal exposure to fertility drugs may initiate cancer in the embryo or parental germ cells. However, an evidence based association between IVF treatments and

Table 1 Anomalies and cancers reported in offspring of IVF

Condition	Reference
Neuroectodermal tumours	White <i>et al</i> ¹⁰ Kobayashi <i>et al</i> ¹¹
Neuroblastoma (Note: associated with the use of fertility drugs only.)	Kramer <i>et al</i> ¹² Michalek <i>et al</i> ¹³
Retinoblastoma	Moll <i>et al</i> ⁶ Cruysberg <i>et al</i> ¹⁴ Anteby <i>et al</i> ¹⁵
Hepatoblastoma	Melamed <i>et al</i> ¹⁵ Toren <i>et al</i> ¹⁶
Clear cell kidney sarcoma	Toren <i>et al</i> ¹⁶
Lymphoma	Kobayashi <i>et al</i> ¹¹
Transposition of the great arteries	Lancaster ¹⁷ Berg <i>et al</i> ¹⁸
Neural tube defects	Lancaster ¹⁷

Several types of congenital anomalies and cancers have been reported in the literature, primarily in the form of case reports. The relation between IVF and these conditions is not definitively established.

cancer development (in the women taking the drugs) has not been established.⁶

Another possible mechanism has been associated with the culture medium used in the IVF test tube. Experiments on mice suggest that subtle changes in the ingredients of the culture media may alter the activity of imprinted genes.⁷ Like fertility drugs, it remains to be seen whether altered gene imprinting can lead to cancers such as retinoblastoma.

A third possible mechanism of carcinogenesis includes inheritance of genetic defects from gametes and embryo trauma performed during routine intracytoplasmic sperm injection (ICSI) type IVF.

The most likely explanation for an increased risk of secondary carcinogenesis is that the population of patients seeking IVF is dissimilar to the general population. Patients seeking IVF are older or have medical problems that interfere with fertility. This poses a problem with any study other than a randomised prospective trial that compares babies from untreated infertility patients who become pregnant versus babies from infertile patients who become pregnant using IVF. To date all studies published are observations of clusters of disease with the suggestion that there is a cause and effect relation.⁸

Table 2 Retinoblastoma in children born through IVF

Reference	Sex and age at diagnosis (months)	Eye	Cause of subfertility	DNA 13q14 analysis	Assisted reproductive technique
Anteby <i>et al</i> ¹⁵	NR, 30	Unilateral	NR	NR	IVF with donor sperm
Moll <i>et al</i> ⁶	F, 38	Both	Unexplained	Exon 8 mutated	IVF with 6 courses of clomid
Moll <i>et al</i> ⁶	M, 15	Left	Maternal cause	Normal	IVF
Moll <i>et al</i> ⁶	F, 34	Right	Unexplained	Normal	IVF
Moll <i>et al</i> ⁶	M, 8.5	Both	Unexplained	Intron 3 mutated	IVF with 8 AI attempts
Moll <i>et al</i> ⁶	F, 32	Left	Paternal cause	Normal	IVF with ICSI
This report	F, 16	Right	Maternal cause	NR	IVF

NR, not reported; AI, artificial insemination; ICSI, intracytoplasmic sperm injection.

Current literature on retinoblastoma in children born through IVF includes three case reports that describe a total of six cases of retinoblastoma in these children. The characteristics of these patients and the patient discussed in this report are listed below.

Figure 1 B-scan ultrasonography. (A) A longitudinal view of the right eye demonstrates a tumour posterior to a closed funnel retinal detachment. (B) A transverse view of the eye demonstrates a rounded, densely calcified tumour. R, retina; S, shadow in orbit; C, calcification; T, tumour; ON, optic nerve.

Retinoblastoma is the most common intraocular cancer of childhood and affects approximately 300 children each year in the United States. Retinoblastoma is a manifestation of a *de novo* deletion or mutation of the q14 band of chromosome 13, occurring as a "second hit" during embryogenesis or the result of two hit deletions in retinal cells. In that it could be the result of chromosomal breakage and deletions in IVF born children, surveillance of retinoblastoma incidence in children born through IVF is warranted.⁷⁻⁹

With the advent of assisted reproductive technology (ART) in 1977, American couples have increasingly turned to such treatments to overcome fertility problems. Nationwide, 99 629 procedures were performed in 2000 by ART. In that year, fertility treatments in which the egg and sperm are handled in the laboratory resulted in 25 228 live births and 35 025 infants. This report expands information on geography and determinants of both ART success and multiple birth risks (beyond that which appears in the 2000 Assisted Reproductive Technology Success Rates¹⁹). Therefore, it seems reasonable to maintain a registry of post-IVF children, to support large epidemiological studies with long term follow up, and prospective randomised studies of infertile couples in order to determine if there is a relation between IVF and cancers such as retinoblastoma.

I Lee, P T Finger

The New York Eye Cancer Center, New York City, NY, USA

I Lee, P T Finger, J A Grifo, A R Rausen,

A Rebarber

New York University School of Medicine, New York City, NY, USA

D H Barad

Albert Einstein College of Medicine, New York City, New York, USA

Correspondence to: P T Finger, The New York Eye Cancer Center, 115 East 61st Street, New York City, NY 10021, USA; pfinger@eyecancer.com

doi: 10.1136/bjo.2003.041160

Accepted for publication 8 January 2004

Supported by The EyeCare Foundation, Inc and Research to Prevent Blindness, New York City, USA.

The authors have no proprietary interest in any material mentioned in this study.

References

- 1 Klip H, Burger CW, deKraker J, *et al*. Risk of cancer in the offspring of women who underwent ovarian stimulation for IVF. *Hum Reprod* 2001;**16**:2451-8.
- 2 Anteby I, Cohen E, Anteby E, *et al*. Ocular manifestations in children born after in vitro fertilization. *Arch Ophthalmol* 2001;**119**:1525-9.
- 3 Moll AC, Imhof SM, Cruysberg JR, *et al*. Incidence of retinoblastoma in children born after in-vitro fertilisation. *Lancet* 2003;**361**:309-10.
- 4 Finger PT, Harbour JW, Karcioglu Z. Risk factors for metastasis in retinoblastoma. *Surv Ophthalmol* 2002;**47**:1-16.
- 5 Moshfeghi DM, Moshfeghi AA, Finger PT. Enucleation. *Surv Ophthalmol* 2000;**44**:277-301.
- 6 Lerner-Geva L, Geva E, Lessing JB, *et al*. The possible association between in vitro fertilization treatments and cancer development. *Int J Gynecol Cancer* 2003;**13**:23-7.
- 7 Doherty AS, Mann MR, Tremblay KD, *et al*. Differential effects of culture on imprinted H19 expression in the preimplantation mouse embryo. *Biol Reprod* 2000;**62**:1526-35.
- 8 Bruinsma F, Venn A, Lancaster P, *et al*. Incidence of cancer in children born after in-vitro fertilization. *Human Reprod* 2000;**15**:604-7.
- 9 Lerner-Geva L, Toren A, Chetrit A, *et al*. The risk for cancer among children of women who underwent in vitro fertilization. *Cancer* 2000;**88**:2845-7.
- 10 White L, Giri N, Vowels MR, *et al*. Neuroectodermal tumours in children born after assisted conception. *Lancet* 1990;**336**:1577.
- 11 Kobayashi N, Matsui I, Tanimura M, *et al*. Childhood neuroectodermal tumours and malignant lymphoma after maternal ovulation induction. *Lancet* 1991;**338**:955.
- 12 Kramer S, Ward E, Meadows AT, *et al*. Medical and drug risk factors associated with neuroblastoma: a case-control study. *J Natl Cancer Inst* 1987;**78**:797-804.
- 13 Michalek AM, Buck GM, Nasca PC, *et al*. Gravid health status, medication use, and risk of neuroblastoma. *Am J Epidemiol* 1996;**143**:996-1001.
- 14 Cruysberg JR, Moll AC, Imhof SM. Bilateral sporadic retinoblastoma in a child born after in vitro fertilization. *Arch Ophthalmol* 2002;**120**:1773, author reply 1773-4.
- 15 Melamed O, Bujanover Y, Hammer J, *et al*. Hepatoblastoma in an infant born to a mother after hormonal treatment for sterility. *N Engl J Med* 1982;**307**:820.
- 16 Toren A, Sharon N, Mandel M, *et al*. Two embryonal cancers after in vitro fertilization. *Cancer* 1995;**76**:2372-4.
- 17 Lancaster PA. Congenital malformations after in vitro fertilization. *Lancet* 1987;**2**:1392-3.
- 18 Berg T, Ericson A, Hillensjö T, *et al*. Deliveries and children born after in-vitro fertilization in Sweden

1982-95: a retrospective cohort study. *Lancet* 1999;**354**:1579-85.

19 National Summary and Fertility Clinic Report. Assisted reproductive technology success rates. *MMWR* 2000;**52**:No SS-9.

Angle closure glaucoma after laser photocoagulation for retinopathy of prematurity

Infantile angle closure glaucoma (ACG) is a rare consequence of retinopathy of prematurity (ROP) and usually occurs a few years after laser treatment for ROP.¹⁻³ A Medline search for ACG following laser photocoagulation extracted only one case. In the case, ACG occurred in 2 weeks after laser photocoagulation and although occurrence of iris bombe in both eyes was described, the mechanism for the ACG was not fully clarified.⁴

We present a case of bilateral ACG that occurred within a several weeks after the laser photocoagulation for ROP. We shall discuss the importance of ultrasound biomicroscopy (UBM) in the diagnosis.

Case report

A baby girl, born at 25 weeks gestation weighing 796 g, was diagnosed with stage 2 plus, zone 2 ROP bilaterally at 33 weeks. Diode laser photocoagulation, 986 applications right eye and 629 left eye, with 200-240 mW, 0.4 second duration, was performed by a paediatric ophthalmologist. On the following day, severe hyphaema was observed bilaterally but there was no evidence of choroidal detachment by B-mode ultrasound sonography. Topical atropine and corticosteroid were started and she was

Figure 1 Photographs of the anterior segment of left eye (A) and right eye (B) obtained with an operative microscope after peripheral iridectomy. The photographs show shallow anterior chambers and persistent pupillary membranes and iridohyaloid vessels (A), and corneal opacity (B).

Figure 2 Composite images of the anterior segment scanned with UBM (top) and schematic representations (bottom). (A) Left eye. Posterior synechia and iris bombe can be seen. C, cornea; L, lens; AC, anterior chamber; S, posterior synechia; PC, posterior chamber. (B) Right eye. The high echo lines (arrowheads) along the corneal endothelium represent the iris adherent to the corneal endothelium and the iris bombe. The lens is displaced into the anterior chamber and the anterior chamber is completely obliterated. The arrow represents the posterior capsule of the lens. Note that the cornea is distended and is thin as a result of the high intraocular pressure. The thickness of the cornea at the centre is 394 μm , and 613 μm in the left eye as measured with a caliper installed in the UBM model 840 Humphrey.

followed up conservatively. During the follow up period, total posterior synechia was formed in the both eyes and the anterior chamber became shallow.

At 39 weeks, the corneal diameter had increased, and the anterior chamber was extremely shallow bilaterally. The intraocular pressure (IOP) in the right eye was elevated to 28 mm Hg, and she was referred to our hospital.

Our examination showed that the corneal diameter was increased to 11 mm bilaterally. Slit lamp examination showed corneal oedema and shallow anterior chamber depth bilaterally, especially in the right eye. The corneal oedema made the funduscopy difficult bilaterally. The IOP under the general anaesthesia was 33 mm Hg right eye and 17 mm Hg left eye. Persistent pupillary membranes and iridohyaloid vessels were observed but rubeosis iridis was not observed (fig 1).

UBM images of anterior segments demonstrated iris bombe bilaterally, and the entire right iris surface was adherent to the corneal endothelium. As a result, the anterior capsule

of the lens was also attached to the corneal endothelium (fig 2). Choroidal detachment and a retrolental mass were not observed by B-mode ultrasound sonography (fig 2).

Peripheral iridectomy was performed bilaterally (fig 1). Postoperatively, her peripheral anterior chamber deepened bilaterally although the lens in the right eye was still adherent to the corneal endothelium. Indirect ophthalmoscopy revealed normal cup to disc ratio. The IOP fell to normal levels bilaterally.

Comment

Shallow anterior chambers in ROP patients are known to be caused by various factors—for example, choroidal detachment after excessive photocoagulation, development of retrolental mass, or relative increment in lens thickness,³ but usually the cause of shallow anterior chamber cannot be determined. In our case, the development of hyphaema after photocoagulation induced posterior synechia, and the iris bombe followed. The displacement of the anterior chamber structures was induced by the forward movement of the iris-lens diaphragm in the right eye, and the ocular fragility in premature baby may explain this deformity.

Vitreous haemorrhage is known to occur in 7.9% of ROP cases after photocoagulation.⁶ In our case, there is a possibility that the hyphaema was derived from vitreous haemorrhage. Another possibility is an accidental photocoagulation of persistent pupillary membranes and/or iridocorneal vessels caused the hyphaema. We are not aware of such morphological changes after photocoagulation for ROP.

ACG that occurs immediately after retinal photocoagulation in ROP patients is rare, but is still an important complication. In ROP patients, the lens and its ligament are weak, and therefore not only ACG but also lens displacement occurred. It is important that we be aware of the possible development of ACG following retinal photocoagulation for ROP.

A Uehara, T Kurokawa, N Gotoh,
N Yoshimura

Department of Ophthalmology, Shinshu University
School of Medicine, Matsumoto, Japan

T Tokushima

Department of Ophthalmology, Nagano Children's
Hospital, Nagano, Japan

Correspondence to: N Yoshimura, MD, PhD,
Department of Ophthalmology and Visual Sciences
Kyoto University Graduate School of Medicine, Kyoto
606-8507, Japan; nagaeye@kuhp.kyoto-u.ac.jp

doi: 10.1136/bjo.2003.037994

Accepted for publication 22 January 2004

References

- 1 **Blodi FC.** Retrolental fibroplasias (retinopathy of prematurity). *Trans Am Acad Ophthalmol Otolaryngol* 1955;**59**:35–8.
- 2 **McCormic AQ, Pratt-Johnson JA.** Angle closure glaucoma in infancy. *Ophthalmic Surg* 1971;**2**:91–3.
- 3 **Smith J, Shvitz I.** Angle-closure glaucoma in adult with cicatricial retinopathy of prematurity. *Arch Ophthalmol* 1984;**102**:371–2.
- 4 **Lee AL, Lee LR, Gole GA.** Angle-closure glaucoma after laser treatment for retinopathy of prematurity. *J AAPOS* 1998;**2**:383–4.
- 5 **Hibino Y, Takahashi M, Majima A.** Studies on ocular functions of cicatricial retinopathy of

prematurity. *Managements of refractive elements.* *Jpn J Clin Ophthalmol* 1978;**32**:655–62.

- 6 **Fallaha N, Lynn MJ, Aaberg TM Jr, et al.** Clinical outcome of confluent laser photocoagulation for retinopathy of prematurity. *J AAPOS* 2002;**6**:81–5.

Sequential treatment of central retinal vein occlusion with intravitreal tissue plasminogen activator and intravitreal triamcinolone

Treatment for central retinal vein occlusion (CRVO) remains disappointing despite recently proposed intraocular surgical techniques.^{1–2} We previously introduced the use of intravitreal tissue plasminogen activator (TPA) for acute central retinal vein occlusion in 1999.³ Numerous investigators have confirmed its safety and suggested that it may have a beneficial role in the treatment of acute central retinal vein occlusion.^{4–7} Although some studies in rabbits suggest the rabbit retina is not permeable to TPA,

Figure 1 (A) Fundus photograph shows an acute CRVO with dilated retinal veins, retinal haemorrhages, and retinal oedema. (B) Fundus appearance 1 month after intravitreal injection of tissue plasminogen activator. (C) Photograph of the same eye 5 months later.

Figure 2 (A) Fluorescein angiogram reveals persistent macular oedema and hyperfluorescence 6 months after intravitreal TPA injection. (B) Late frames show resolution of macular oedema 6 weeks after intravitreal triamcinolone.

investigators found the porcine retina was, in fact, permeable to TPA.^{8,9} Our clinical experience with intravitreal TPA in humans with CRVO and large submacular haemorrhages strongly suggests that intravitreal TPA does cross the human retina. Greenberg *et al* were the first to report the possible beneficial effect of intravitreal steroids on patients with chronic CRVO.¹⁰ This report describes a sequential treatment strategy for patients with CRVO who present early in the course of their disease and can be performed in the office while avoiding the risks of vitrectomy. It utilises intravitreal TPA in the acute phase of the vein occlusion to attempt clot lysis, and then treats any remaining vascular leakage with intravitreal triamcinolone.

Case report

A 59 year old obese, hypertensive flight instructor presented with a sudden decrease in vision for 7 days in the right eye. Vision was 20/400 right eye and 20/20 left eye. The patient was diagnosed with an acute CRVO in the right eye (fig 1A). The left eye was normal. After being advised of the risks and benefits, the patient elected to undergo intravitreal injection of TPA (75 µg). Thirteen days later, the patient noted marked improvement in vision with 20/60 vision. Thirty four days after the injection, the patient's vision was 20/30 (Fig 1B).

Six months after intravitreal TPA injection, the vision remained 20/30, but the patient still complained of metamorphopsia and blurry vision despite resolution of other findings of CRVO (fig 1C). Fluorescein angiogram (FA) revealed persistent macular oedema (fig 2A). Optical coherence tomography (OCT) showed the foveal thickness to be 331 µm with mild intraretinal oedema. After

being advised of the risks and the benefits, the patient then underwent injection of intravitreal triamcinolone (4 mg).

Six weeks after the intravitreal triamcinolone, the FA returned to normal and OCT showed decreased foveal thickness from 331 µm to 291 µm. The patient reported a significant improvement in vision with decreased metamorphopsia. Vision was 20/25 with no late leakage on the fluorescein angiogram (fig 2B).

Comment

To our knowledge, this represents the first published case of CRVO treated sequentially with intravitreal TPA for the acute phase and intravitreal triamcinolone for the chronic phase. TPA is a drug that must be used early in the course of thrombus formation to be effective. We do not recommend its use for patients with chronic symptoms. Intravitreal steroids appear to decrease the blood-retinal barrier breakdown and macular oedema, but recurrent oedema may occur since the steroids do not appear to affect the thrombus itself.

J M L, J J K, M C Cheung

Ophthalmology, University of California at San Francisco, San Francisco, CA, USA

J M Lahey, J J K

Ophthalmology, Kaiser Permanente, Hayward, CA, USA

Correspondence to: J M Lahey, MD, Ophthalmology (Retina), Kaiser Permanente Medical Center, 27400 Hesperian Boulevard, Hayward, CA 94545-4299, USA; Mike.Lahey@kp.org

doi: 10.1136/bjo.2004.043406

Accepted for publication 3 February 2004

References

- 1 Opremac EM, Bruce RA, Lomeo MD, *et al*. Radial optic neurotomy for central retinal vein occlusion: a retrospective pilot study of 11 consecutive cases. *Retina* 2001;21:408–15.
- 2 Weiss JN, Byone LA. Injection of tissue plasminogen activator into a branch retinal vein in eyes with central retinal vein occlusion. *Ophthalmology* 2001;108:2249–57.
- 3 Lahey JM, Fong DS, Kearney J. Intravitreal tissue plasminogen activator for acute central retinal vein occlusion. *Ophthalmic Surg Lasers* 1999;30:427–34.
- 4 Weizer JS, Fekrat S. Intravitreal tissue plasminogen activator for the treatment of central retinal vein occlusion. *Ophthalmic Surg Lasers Imaging* 2003;34:350–2.
- 5 Glacet-Bernard A, Kuhn D, Vine AK, *et al*. Treatment of recent onset central retinal vein occlusion with intravitreal tissue plasminogen activator: a pilot study. *Br J Ophthalmol* 2000;84:609–13.
- 6 Ghazi NG, Noureddine B, Haddad RS, *et al*. Intravitreal tissue plasminogen activator in the management of central retinal vein occlusion. *Retina* 2003;23:780–4.
- 7 Elman MJ, Raden RZ, Carrigan A. Intravitreal injection of tissue plasminogen activator for central retinal vein occlusion. *Trans Am Ophthalmol Soc*. 2001;99: 219–21; discussion 222–3.
- 8 Kamei M, Misono K, Lewis H. A study of the ability of tissue plasminogen activator to diffuse into the subretinal space after intravitreal injection in rabbits. *Am J Ophthalmol* 1999;128:739–46.
- 9 Mahmoud TH, Peng Y-W, Proia A, *et al*. Intravitreal tissue plasminogen activator penetrates the retinal veins in a porcine model of vascular occlusion. *Invest Ophthalmol Vis Sci*.

2002;43: E-abstract 3533, Presented at ARVO, 2002.

- 10 Greenberg PB, Martidis A, Rogers AH, *et al*. Intravitreal triamcinolone acetate for macular oedema due to central retinal vein occlusion. *Br J Ophthalmol* 2002;86:247–8.

Severe post-laser suprachoroidal haemorrhaging in a diabetic patient receiving anticoagulants

Although the aetiology is not well understood, expulsive suprachoroidal haemorrhaging (ESH) is the most severe complication associated with intraocular surgery. Anticoagulants are considered a risk factor for spontaneous suprachoroidal haemorrhaging in cases with high myopia, age related macular degeneration, and diabetic retinopathy.^{1–5} However, ESH post photocoagulation is extremely rare regardless of anticoagulant therapy. We have experienced a severe case of post-laser ESH correlated with anticoagulant therapy, which resulted in irreversible visual disturbance.

Case report

A 70 year old woman was diagnosed with pre-proliferative diabetic retinopathy based on fluorescein angiographic examinations. Two months before diagnosis, she had right eye cataract surgery. During the past 6 years, the patient received warfarin (4 mg/day) and aspirin (81 mg/day) because of atrial fibrillation after myocardial infarction. Laser photocoagulation was performed in her right eye with a Nidek MC-7000, yellow-green laser. Operating conditions were 200–280 burns per session with a spot size 200 µm, exposure 0.2 seconds, power 100–120 mW using a QuadraSpheric contact lens (Volk, Tokyo, Japan). Treatment was separated into three partitions with a minimum 2 week interval between sessions. Three days after final photocoagulation, the patient had a sudden visual loss to hand movements. In slit lamp examinations, the retina seemed to be attached to the posterior surface of the implanted intraocular lens. Severe choroidal detachment was found by fundus examination (fig 1). The B-mode ultrasonography showed massive haemorrhaging in the choroidal space (fig 2). In systemic examinations, multiple purple spots were observed in both her arms. Microhaematuria was also found. Blood examination revealed blood sugar 167 mg/dl; platelet number 179 000 × 10⁹/l; PT% 19% (control 70–120); PTs 28.7 seconds; PT INR 4.72 (control 1); APTT 80.2 seconds (control 24.0–38.0); and bleeding time 5 minutes. Although surgery was planned to proceed as soon as the anticoagulant was washed out, her right eye lost all light perception before treatment.

Comment

We have described a case of ESH after laser photocoagulation in a patient receiving anticoagulant therapy. Laser photocoagulation is known as an effective treatment for various ocular diseases and is a widely used, non-incision surgical procedure. However, a number of complications have been reported, with some citing an irreversible visual disturbance.⁶ On the other hand, anticoagulant therapy is prevalent after cardiac/brain infarctions, which necessitate long term coagulation system management. In the present case, the PT INR was extremely prolonged (a respected value of 2–3 is

Figure 1 A fundus photograph of the patient after laser photocoagulation. A severe choroidal detachment associated with secondary retinal detachment was found.

Figure 2 The findings of B-mode ultrasound examination. A massive haemorrhage in the choroids is present.

appropriate for post-cardiac infarction). Presumably, choroidal microbleeding initiated by photocoagulation persisted because of an overly suppressed coagulation system; blood pooled in the choroidal space, which assumed an ESH. To our knowledge there is only one other similar case reported by Khairallah *et al* that showed post-laser choroidal haematoma in a diabetic patient treated with anticoagulant.⁷ Even though ESH incidence is low, extreme caution must be exercised when performing laser therapy in patients using anticoagulants, because of potentially serious outcomes. An age of 65 years or more, history of stroke, history of gastrointestinal bleeding, a serious morbid condition (recent myocardial infarction, renal insufficiency, or severe anaemia), and atrial fibrillation are five high risk factors for major bleeding in outpatients treated with warfarin.⁸ If possible, preoperative coagulation system examinations are recommended for high risk patients receiving anticoagulant treatments.

A Mikawa, S Honda, I Sugita, N Okamoto, H Toda

Department of Ophthalmology, Kitano Hospital, Tazukekafukai Medical Research Institute, Osaka, Japan

Correspondence to: A Mikawa, Department of Ophthalmology, Kitano Hospital, Tazukekafukai Medical Research Institute, 2-4-20 Ougimachi, Kita-ku, Osaka 530-8480, Japan; a-matsuo@kitano-hp.or.jp

doi: 10.1136/bjo.2004.043489

Accepted for publication 14 February 2004

References

- 1 Chak M, Williamson TH. Spontaneous suprachoroidal haemorrhage associated with high myopia and aspirin. *Eye* 2003;17:525-7.
- 2 Fadi EB, William H, Jarrett H, *et al*. Massive hemorrhage complicating age-related macular degeneration. *Ophthalmology* 1986;93:1581-92.
- 3 Edwards P. Massive choroidal hemorrhage in age-related macular degeneration: a complication of anticoagulant therapy. *J Am Optom Assoc* 1996;67:223-6.
- 4 Yang SS, Arthur D, McDonald R. Massive spontaneous choroidal hemorrhage. *Retina* 2003;23:139-44.
- 5 Raj A, Sekhri R, Salam A, *et al*. Massive subretinal bleed in a patient with background diabetic retinopathy and on treatment with warfarin. *Eye* 2003;17:649-652.
- 6 Zweng HC, Little HL, Hammond AH. Complications of argon laser photocoagulation. *Trans Am Acad Ophthalmol Otolaryngol* 1974;78:OP195-204.
- 7 Khairallah M, Chachia N. Post-laser choroidal hematoma in a diabetic patient treated with an oral anticoagulant. *J Fr Ophthalmol* 1994;17:138-140.
- 8 Landefeld CS, Goldman L. Major bleeding in outpatients treated with warfarin: incidence and prediction by factors known at the start of outpatient therapy. *Am J Med* 1989;87:144-52.

MAILBOX

Authors' reply

We appreciate the interest and many comments we have received regarding our recent article.¹ In reply to the comments by Dr Vedantham, we acknowledge the paucity of experimental data to prove that accurate placement of corticosteroids into the sub-Tenon's space provides good drug penetration into the eye. However, the studies to the contrary cited by Vedantham have all used needles to make such "accurate placement," including the study by Jennings *et al*,² which utilised the technique described by Tessler.³ Use of needles represents not only a potential hazard to the eye in terms of accidental globe penetration, but also makes it much more difficult to place any sub-Tenon's injection under the posterior Tenon's capsule near the macula and/or around the optic nerve. It has been shown that many injections intended for the sub-Tenon's space merely end up somewhere in the orbit outside of Tenon's capsule.⁴ We believe that our method using a 23 gauge blunt, curved, long cannula (the one we used was No HS-2764 by Handaya Co, Ltd, Tokyo, Japan) assures accurate placement into the target space, thereby increasing therapeutic efficacy and obviating the need for globe invasive procedures such as intravitreal injection of corticosteroids, corticosteroid intravitreal implants, and/or therapeutic vitrectomy.

However, we are in agreement with Vedantham, in that ultimately corticosteroid placed outside of the eye may be no match for the efficacy that may be obtained by corticosteroid placed inside the eye. Yet we have found such a high efficacy rate for the trans-Tenon's retrobulbar infusion of triamcinolone in uveitis that we can conceive of no reason why this treatment should not be tried *before* procedures such as intravitreal injections that carry risks of severe complications are considered. For example, as also pointed out by Vedantham, the risks of intravitreal corticosteroid injections even include development of a rare form of mycobacterial endophthalmitis.⁵ Therefore, the risk to the eye of intravitreal procedures, especially when involving corticosteroid administration, cannot be taken lightly. Furthermore, we believe that the reason why sub-Tenon's injections of corticosteroids have not become popular among retina specialists who for example treat diabetic macular oedema, is more likely related to the lower efficacy rate when using needles as opposed to the technique using an infusion cannula that we advocate. Lastly, obtaining the infusion cannula seems like a small inconvenience (and an even smaller cost) to the physician compared to the risk of doing intravitreal injections of corticosteroids as a treatment of first choice as advocated by Vedantham. We strongly encourage all uveitis and retina specialists who have up until now been disappointed with the efficacy of their sub-Tenon's corticosteroid injections, to make the effort to obtain an appropriate cannula and revise their technique before jumping to intravitreal procedures.

In reply to the first comment by Dr Mehta,⁶ we acknowledge the current WHO guidelines, revised for 2003, that include recommendations for extrapulmonary tuberculosis.⁷ However, we would also like to amend Mehta's comment, in that the WHO admits in those guidelines that there are many

regimens with reported efficacy including a 6 month regimen of rifampicin (with streptomycin also given in the initial phase only) for meningeal tuberculosis. Furthermore, the WHO recommendations are for *active* extrapulmonary tuberculosis that has been diagnosed by specimen examination or strong clinical evidence, and give no recommendations for *latent* infection. As we have previously reported in a series on intraocular tuberculosis, systemic examination failed to identify a focus of active tuberculosis in the majority of our patients,⁸ and we have come to suspect that the uveitis we observed may be an immune response to latent tuberculosis antigen sequestered elsewhere. Therefore, the patients we described were given a diagnosis of "presumed intraocular tuberculosis," that is with uveitis presumed to be related to the *Mycobacterium tuberculosis* organism. Furthermore, we would like to clarify that in the cases of presumed ocular tuberculosis that received trans-Tenon's retrobulbar triamcinolone infusion,¹ this treatment was judged to be effective in two of three eyes. Regardless, since the focus of active or latent tuberculosis was never identified in our patients, a two drug regimen of isoniazid and rifampicin was used as a therapeutic trial for antituberculosis therapy. A similar therapeutic trial for ocular tuberculosis, albeit with isoniazid alone, has been previously advocated in Japan by Ishihara and Ohno.⁹

With regard to the second comment, among the 16 patients who were receiving some form of systemic immunosuppressive therapy, we did not notice any difference in outcome when compared to patients who were not on immunosuppressive therapy. In other words, the efficacy of trans-Tenon's retrobulbar triamcinolone infusion was the same. However, we suspect that the recurrence rate after triamcinolone infusion may be different, and we are currently investigating this possibility.

A A Okada, T Wakabayashi

Kyoin Eye Center, Kyoin University School of Medicine, Tokyo, Japan

Correspondence to: Annabelle A Okada, Kyoin Eye Center, Kyoin University School of Medicine, Tokyo, Japan; aokada@po.iiijnet.or.jp

doi: 10.1136/bjo.2003.038851

Accepted for publication 20 November 2003

References

- Okada AA, Wakabayashi T, Morimura Y, et al. Trans-Tenon's retrobulbar triamcinolone infusion for the treatment of uveitis. *Br J Ophthalmol* 2003;**87**:968–71.
- Jennings T, Rusin MM, Tessler HH, et al. Posterior sub-Tenon's injections of corticosteroids in uveitis patients with cystoid macular edema. *Jpn J Ophthalmol* 1988;**32**:385–91.
- Tessler H. Uveitis. In: Peyman GA, Sanders DR, Goldberg MF, eds. *Principles and practice of ophthalmology*, Vol 2. 1st ed. Philadelphia: Saunders, 1980:1554–629.
- Freeman WR, Green RL, Smith RE. Echographic localization of corticosteroids after periorbital injection. *Am J Ophthalmol* 1987;**103**:281–8.
- Benz MS, Murray TG, Dubovy SR, et al. Endophthalmitis caused by *Mycobacterium chelonae* abscessus after intravitreal injection of triamcinolone. *Arch Ophthalmol* 2003;**121**:271–3.
- Mehta SA. Trans-Tenon's triamcinolone and pretreatment. *Br J Ophthalmol* 2003. www.bjophthalmol.com/cgi/eletters/87/8/968#183, 1 Aug 2003.
- World Health Organization. Case definitions (p 24) and standardised treatment regimens (p 35–36). In: *Treatment of tuberculosis: guidelines for national programmes*, 3rd ed. (WHO/CDS/TB/2003.313). Geneva: WHO, 2003 (accessed on 13 October 2003).
- Morimura Y, Okada AA, Kawahara S, et al. Tuberculin skin testing in uveitis patients and treatment of presumed intraocular tuberculosis in Japan. *Ophthalmology* 2002;**109**:851–7.
- Ishihara M, Ohno S. [Ocular tuberculosis]. *Nippon Rinsho* 1998;**56**:3157–61.

Owls' eyes move

"Double crossed," the cover illustration and article by Schwab on the barn owl refers to the alleged immobility of the owl's eyes.¹ This is a myth which should not be perpetuated in the *BJO*. The owl's eyes do in fact move, and while the amount is not large, it is just enough for two papers on the subject.^{2–3} The phrase "nearly immobile" is preferable.

M J Steinbach

Correspondence to: Professor Martin J Steinbach, University of Toronto, Department of Ophthalmology and Vision Sciences, Toronto Western Hospital, 399 Bathurst Street, Toronto M5T 2S8, Canada; mjs@yorku.ca

doi: 10.1136/bjo.2004.042291

Accepted for publication 16 January 2004

References

- Schwab IR. Double crossed. *Br J Ophthalmol* 2003;**87**:1442.
- Steinbach MJ, Money KE. Eye movements of the owl. *Vis Res* 1973;**13**:889–91.
- Steinbach MJ, Angus RG, Money KE. Torsional eye movements of the owl. *Vis Res* 1974;**14**:745–6.

NOTICES

Low vision care

The latest issue of *Community Eye Health* (No 49) deals with the problems and management of low vision. For further information please contact: Journal of Community Eye Health, International Resource Centre, International Centre for Eye Health, Department of Infectious and Tropical Diseases, London School of Hygiene and

Tropical Medicine, Keppel Street, London WC1E 7HT, UK (tel: +44 (0)20 7612 7964; email: Anita.Shah@lshtm.ac.uk; online edition: www.jceh.co.uk). Annual subscription (4 issues) UK £28/US\$45. Free to developing country applicants.

Elimination of avoidable blindness

The 56th World Health Assembly (WHA) considered the report on the elimination of avoidable blindness (doc A56/26) and urged Member States to: (1) Commit themselves to supporting the Global Initiative for the Elimination of Avoidable Blindness by setting up a national Vision 2020 plan by 2005; (2) Establish a national coordinating committee for Vision 2020, or a national blindness prevention committee to help implement the plan; (3) Implement the plan by 2007; (4) Include effective monitoring and evaluation of the plan with the aim of showing a reduction in the magnitude of avoidable blindness by 2010; (5) To support the mobilisation of resources for eliminating avoidable blindness. The WHA also urged the Director-General to maintain and strengthen WHO's collaboration with Member States and the partners of the Global Initiative for the Elimination of Avoidable Blindness as well as aid in the coordination and support of national capability.

4th International Congress on Autoimmunity

The 4th International Congress on Autoimmunity will take place 3–7 November 2004 in Budapest, Hungary. The deadline for the receipt of abstracts is 20 June 2004. Further details: Kenes International Global Congress Organisers and Association Management Services, 17 Rue du Cendrier, PO Box 1726, CH-1211 Geneva 1, Switzerland (tel: +41 22 908 0488; fax: +41 22 732 2850; email: autoim04@kenes.com; website: www.kenes.com/autoim2004).

XVI International Congress for Eye Research

The XVI International Congress for Eye Research will be held on 29 August – 3 September 2004 in Sydney, Australia. For further information, please contact: icer2004@tourhosts.com.au (website: www.tourhosts.com.au/icer2004).

Glaucoma Society Silver Jubilee Meeting 2004

The Silver Jubilee Meeting and Dinner for the Glaucoma Society will be held on 3 December 2004 at the Royal College of Physicians in Regents Park, London. The meeting will take place between 8.30am and 5pm and the dinner will be held between 6.30pm and 10pm. For further information, please contact: Janet Flowers, Administrator, 29 Quarry Hill, Grays, Essex, RM17 5BT (tel: 01375 383172; e-mail: glausoc@ukeire.freeseerve.co.uk).

PostScript

LETTERS

If you have a burning desire to respond to a paper published in *BJO*, why not make use of our "rapid response" option?

Log onto our website (www.bjophthalmol.com), find the paper that interests you, and send your response via email by clicking on the "eletters" option in the box at the top right hand corner.

Providing it isn't libellous or obscene, it will be posted within seven days. You can retrieve it by clicking on "read eletters" on our homepage.

The editors will decide as before whether to also publish it in a future paper issue.

Cystoid macular oedema with trypan blue use

We present a large comparative series of trypan blue use in cataract surgery. This series of trypan blue used in all eyes regardless of cataract severity may be unique. We found an apparent increased rate of cystoid macular oedema (CMO) associated with trypan blue use.

Melles *et al*'s¹ report on the use of trypan blue in cataract extraction in 1999 combined with Apple *et al*'s²⁻⁴ series on dye enhanced cataract surgery facilitated widespread acceptance of this technique. The dye has been shown to stain basement membrane of lens capsule.⁵ Trypan blue is now widely used to assist in cataract extraction when visualisation of the anterior capsule is poor because of loss of red reflex. Trypan blue has also been used to improve contrast during cataract extraction in eyes with corneal opacities⁶ and to stain internal limiting membrane and epiretinal membrane during vitreoretinal surgery.^{7,8} The safety profile of trypan blue appears good with no adverse effects reported in several large series.⁹⁻¹¹

Patients and methods

In this retrospective, comparative study we identified a consecutive series of 75 patients (group A) in whom trypan blue had been used "routinely" regardless of cataract type or density. A consecutive series of 94 patients (group B) who had routine phacoemulsification by the same surgeon were used as a control group.

Apart from the use of trypan blue to facilitate capsulorhexis, standard phacoemulsification techniques were used in both groups.

The data from the two cohorts were compared using mean and standard deviations for continuous variables such as age, and proportions for categorical variables such as sex. For acuity a numeric ordinal score was created from 1 to 10 by placing all the recorded acuities in order. This numeric ordinal score allowed us to plot the data using box plots, and to analyse the data using non-parametric methods to produce p values where necessary.

Figure 1 (A) Preoperative best corrected acuity (group A = 75, group B = 94); (B) preoperative acuity without co-morbidity (group A = 66, group B = 93); (C) postoperative acuity without co-morbidity. Acuity key: 6/5 = 1; 6/6 = 2; 6/9 = 3; 6/12 = 4; 6/18 = 5; 6/24 = 6; 6/36 = 7; 6/60 = 8; 2/60 = 9; <2/60 = 10.

Results

The two groups compared favourably with regard to age and sex (table 1) but the preoperative best corrected acuity was worse

in group A (fig 1A). Group A also had a higher proportion of patients with ocular co-morbidity (table 1).

If all the cases with co-morbidity are removed from both groups, the preoperative best corrected acuity remains worse in group A, suggesting that the cataracts in this group were visually of greater significance (fig 1B).

There were four patients with clinically apparent CMO in group A and no cases in group B. The CMO was confirmed in all cases by the vitreoretinal specialist (PRS) using standard biomicroscopy techniques with a fundal contact lens. The incidence of clinically apparent CMO was 5.3% in group A and 0% in group B ($p = 0.037$).

All cases of CMO were treated with a combination of topical steroid and non-steroidal anti-inflammatory drops and resolved completely with good visual outcomes.

The postoperative acuities compared favourably in both groups after patients with co-morbidity were removed from both groups (fig 1C).

Comment

Cystoid macular oedema resulting in visual loss occurs in up to 3.8% of cases following cataract extraction by phacoemulsification without posterior capsule rupture^{12,13} and up to 13% of cases with posterior capsule rupture.^{14,15} Ursell *et al*¹⁶ demonstrated an incidence of angiographic CMO of 19% following phacoemulsification (none of the patients in this group had clinically apparent CMO).

CMO has not been reported as a result of trypan blue use, but trypan blue has been shown to cause damaged photoreceptors in rabbit eyes after intravitreal injection.¹⁷ Trypan blue has also been shown to be carcinogenic and teratogenic in animal studies.^{18,19} The clinical significance or relevance to a human population of the animal studies is uncertain.

Trypan blue has been shown to inadvertently stain posterior capsule²⁰ and intra-ocular lens implants.^{21,22}

None of the patients in our group who developed CMO had associated posterior capsule rupture or ocular co-morbidity and all of the cases were treated successfully with good visual outcomes.

Dada *et al*²³ and Sharma *et al*²⁴ suggested the use of trypan blue in routine cases to aid in training of junior surgeons. Our study would suggest some caution with this approach in view of the apparent increase in the incidence of CMO with trypan blue use.

Table 1 Age and sex distribution and co-morbidity

Variable	Group A (n = 75)	Group B (n = 94)
Mean age (SD)	79.4 (9.8)	78.4 (8.5)
Sex:		
Male	25 (33.3%)	31 (33%)
Female	50 (66.7%)	63 (67%)
ARMD	7	1
CVA	1	0
ERM	1	0

ARMD, age related macular degeneration; CVA, cerebrovascular accident; ERM, epiretinal membrane.

The preoperative best corrected acuity was decreased in the group in which trypan blue was used. This suggests that the cataracts in this group were of greater density, possibly requiring more energy to remove using phacoemulsification. The energy used during surgery however was not recorded. The CMO may therefore be a reflection of higher energy used in denser cataract.

A prospective trial with matched cohorts is required to prove the suggested higher incidence of CMO with trypan blue use. OCT scanning of the maculas in both groups would give non-invasive objective evidence of CMO.

We suggest the following steps to limit the apparent complication of CMO with trypan blue use:

- Use the smallest amount and lowest concentration of trypan blue possible (trypan blue in concentrations as low as 0.0125% has been shown to effectively stain the anterior capsule²⁵)
- Increase postoperative steroid or anti-inflammatory drops prophylactically
- Use only in appropriate cases—that is, with poor visualisation of the anterior capsule.

**P Gouws, M Merriman, S Goethals,
P R Simcock, R J Greenwood, G Wright**
Bristol Eye Hospital, Lower Maudlin Street,
Bristol BS1 2LX, UK

Correspondence to: P Gouws, Bristol Eye Hospital,
Bristol, UK; pieter@gouws.freeserve.co.uk

doi: 10.1136/bjo.2004.041756

Accepted for publication 12 February 2004

References

- Melles GR, de Waard PW, Pameyer JH, *et al*. Trypan blue capsule staining to visualize the capsulorhexis in cataract surgery. *J Cataract Refract Surg* 1999;**25**:7–9.
- Pandey SK, Werner L, Escobar-Gomez M, *et al*. Dye-enhanced cataract surgery. Part 1: Anterior capsule staining for capsulorhexis in advanced/white cataract. *J Cataract Refract Surg* 2000;**26**:1052–9.
- Werner L, Pandey SK, Escobar-Gomez M, *et al*. Dye-enhanced cataract surgery. Part 2: Learning critical steps of phacoemulsification. *J Cataract Refract Surg* 2000;**26**:1060–5.
- Pandey SK, Werner L, Escobar-Gomez M, *et al*. Dye-enhanced cataract surgery. Part 3: Posterior capsule staining to learn posterior continuous curvilinear capsulorhexis. *J Cataract Refract Surg* 2000;**26**:1066–71.
- Singh AJ, Sarodia UA, Brown L, *et al*. A histological analysis of lens capsules stained with trypan blue for capsulorhexis in phacoemulsification cataract surgery. *Eye* 2003;**17**:567–70.
- Bhartiya P, Sharma N, Ray M, *et al*. Trypan blue assisted phacoemulsification in corneal opacities. *Br J Ophthalmol* 2002;**86**(8):857–9.
- Perrier M, Sebag M. Trypan blue-assisted peeling of the internal limiting membrane during macular hole surgery. *Am J Ophthalmol* 2003;**135**:903–5.
- Li K, Wong D, Hiscott P, *et al*. Trypan blue staining of internal limiting membrane and epiretinal membrane during vitrectomy: visual results and histopathological findings. *Br J Ophthalmol* 2003;**87**:216–9.
- Jacob S, Agarwal A, Agarwal A, *et al*. Trypan blue as an adjunct for safe phacoemulsification in eyes with white cataract. *J Cataract Refract Surg* 2002;**28**:1819–25.
- Melles GR, de Waard PW, Pameijer JH, *et al*. [Staining the lens capsule with trypan blue for

visualizing capsulorhexis in surgery of mature cataracts]. *Klin Monatsbl Augenheilkd* 1999;**215**:342–4. (In German).

- Van Dooren BT, de Waard PW, Poort-van Nieuhuys H, *et al*. Corneal endothelial cell density after trypan blue capsule staining in cataract surgery. *J Cataract Refract Surg* 2002;**28**:574–5.
- Malik AR, Qazi ZA, Gilbert C. Visual outcome after high volume cataract surgery in Pakistan. *Br J Ophthalmol* 2003;**87**:937–40.
- Riley AF, Malik TY, Grupcheva CN, *et al*. The Auckland Cataract Study: co-morbidity, surgical techniques, and clinical outcomes in a public hospital service. *Br J Ophthalmol* 2002;**86**:185–90.
- Chan FA, Mathur R, Ku JJK, *et al*. Short-term outcomes in eyes with posterior capsule rupture during cataract surgery. *J Cataract Refract Surg* 2003;**29**(3):537–41.
- Blomquist PH, Rugwani RM. Visual outcomes after vitreous loss during cataract surgery performed by residents. *J Cataract Refract Surg* 2002;**28**:847–52.
- Ursell PG, Spalton DJ, Whitcup SM, *et al*. Cystoid macular edema after phacoemulsification: relationship to blood-ocular barrier damage and visual acuity. *J Cataract Refract Surg* 1999;**25**:1492–7.
- Veckeneer M, van Overdam K, Monzer J, *et al*. Ocular toxicity study of trypan blue injected into the vitreous cavity of rabbit eyes. *Graefes Arch Clin Exp Ophthalmol* 2001;**39**:698–704.
- Chung KT. The significance of azo-reduction in the mutagenesis and carcinogenesis of azo dyes. *Mutat Res* 1983;**114**:269–81.
- Ema M, Kanoh S. Studies on the pharmacological bases of fetal toxicity of drugs. Effect of trypan blue on pregnant rats and their offspring. *Nippon Yakurigaku Zasshi* 1982;**79**:369–81.
- Birchall W, Raynor MK, Turner GS. Inadvertent staining of the posterior lens capsule with trypan blue dye during phacoemulsification. *Arch Ophthalmol* 2001;**119**:1082–3.
- Werner L, Apple DJ, Crema AS, *et al*. Permanent blue discoloration of a hydrogel intraocular lens by intraoperative trypan blue. *J Cataract Refract Surg* 2002;**28**:1279–86.
- Fritz WL. Digital image analysis of trypan blue and fluorescein staining of anterior lens capsules and intraocular lenses. *J Cataract Refract Surg* 2002;**28**:1034–8.
- Dada T, Ray M, Bhartiya P, *et al*. Trypan-blue-assisted capsulorhexis for trainee phacoemulsification surgeons. *J Cataract Refract Surg* 2002;**28**:575–6.
- Sharma N, Bhartiya P, Sinha R, *et al*. Trypan blue assisted phacoemulsification by residents in training. *Clin Experiment Ophthalmol* 2002;**30**:386–7.
- Yetik H, Devranoglu K, Ozkan S. Determining the lowest trypan blue concentration that satisfactorily stains the anterior capsule. *J Cataract Refract Surg* 2002;**28**:988–91.

Familial intraocular cysts in association with anisometropia

Vitreous cysts are rare. Their origin is postulated to be a congenital remnant of the primary hyaloidal system¹ or ciliary body pigment epithelium.² Although most vitreal cysts are asymptomatic, some may cause intermittent visual field defects. Treatment is seldom indicated, though laser photocoagulation or vitrectomy have been suggested.^{1,3}

Iris cysts include iris pigment epithelial cysts and stromal cysts.⁴ The former may get dislodged into either the anterior chamber or into the vitreous chamber. They become symptomatic when they enlarge and occlude the visual axis. Treatment includes aspiration or surgical excision of the cyst.

Both vitreous and iris cysts have been previously reported as sporadic findings. In this report, we present the clinical and echographic manifestations of intraocular cysts in two siblings.

Case report

Two sisters, 11 and 3 years old, were referred for evaluation because of intraocular cysts and amblyopic fellow eyes. They were the products of a full term normal pregnancy with an uneventful perinatal history. Their parents were not relatives. Past medical history was unremarkable. The children in the family were reportedly healthy with no ocular pathology but were inaccessible for examination.

The older sister was known to have worn glasses since the age of 7 years. She complained of intermittent obscuration of vision in her right eye. Her vision was 6/12, J1 right eye and counting fingers at 1 metre with J14 left eye. By indirect ophthalmoscopy of her right eye, a round pigmented, cystic structure was observed in the vitreous cavity (fig 1). The left fundus showed myopic chorioretinal changes with a tilted optic disc. Retinoscopy showed marked myopic anisometropia of +1.00 –1.25×75° right eye and –10.50 –2.00×95° left eye. Ultrasonography disclosed a 3.05 mm cystic, round, hypoechogenic vitreal structure (fig 1). It was partially mobile with vitreal after-movements and was tethered to fine vitreal strands. Its walls showed internal reflectivity of 60%, whereas its contents had very low (<5%) reflectivity.

The younger sister was fitted with spectacles at the age of 7 months because of anisometropic myopia. Her visual acuity (picture cube) was 0.03 right eye with unsteady fixation and 0.2 left eye. In her left eye a cystic, pigmented lesion was attached to the posterior iris surface and extended into the anterior vitreous (fig 2). It was located in the superotemporal quadrant causing adjacent lenticular cortical opacities. Indirect ophthalmoscopy revealed bilateral mild retinal myopic changes. High frequency echography of the iris lesion disclosed a cyst with hypoechogenic content measuring 3.68 mm in diameter (fig 2). Cycloplegic refraction showed anisometropic myopia of –7.5–1.00 ×90° right eye and –0.5–3.50 ×80° left eye.

Figure 1 (A) Homogeneously pigmented vitreous cyst. (B) B-scan echography that demonstrates a round, echolucent vitreous cyst bound by fine vitreal strands.

Figure 2 (A) Cystic structure extending from the posterior layer of the iris in the superotemporal quadrant. (B) High frequency echography that shows a cystic structure attached to the posterior iris.

The poor vision in the fellow eye (without cyst) was attributed to the high anisometropia present in both sisters. They were prescribed glasses with anti-amblyopic therapy by patching. No invasive therapy of the cysts in the eye with better vision was recommended.

Comment

This reports the unusual association between vitreous and iris cysts found in two siblings. The familial clustering of pigmented intraocular cysts suggests a hereditary aetiology in these sisters. Sallo *et al*⁶ reported on the occurrence of primary marginal pigment epithelial iris cysts in four members of one family, thus postulating an autosomal dominant heredity pattern. Nork and Millecchia¹ reported an association between vitreous cyst in a patient and corneal dermoid in her son. A familial association between vitreous cyst and iris cyst was not previously reported. Our study, in accordance with previous reports, emphasises the need for examination of family members once an intraocular cyst has been diagnosed in a young child.

The second unusual association described in these siblings is the deeply amblyopic fellow eye with high anisometric myopic astigmatism. It is not clear whether axial myopia developed because of the amblyopia or whether the amblyopia is secondary to anisometropia. Amblyopia in anisometropia results from the suppression of cortical vision centres that receive inputs from the chronically defocused eye. Also, the eyes of a child

with anisometric amblyopia look normal to the family, leading to delay in detection and treatment.

The physical characteristics of the vitreous cyst we described, including its confinement to the region of Cloquet's canal, are similar to those reported by others.¹ This suggests that the cysts may be remnants of the persistent fetal vasculature, though this manifestation was not included in Goldberg's description of this disease.⁶ However, since no surgical excision was performed, we may postulate regarding their cellular origin. Nork and Millecchia¹ suggested after histological studies, that the cyst origin was pigment epithelial-type cells. In our study, indirect evidence that the cysts originated from pigment epithelium include their homogeneous brown pigmentation, medium reflectivity, and continuation of iris cyst with the posterior iris surface. The cellular origin of the vitreal cyst is less obvious. It can either be a primary congenital hyaloidal remnant or a cyst that detached from the iris during childhood. Only few have reported on vitreous cysts jarring loose from the ciliary body pigment epithelium.²

R Amer, I Anteby

Department of Ophthalmology, Hadassah University Hospital, Jerusalem, Israel

Correspondence to: Radgonda Amer, MD, Department of Ophthalmology, Hadassah University Hospital, POB 12000, Jerusalem 91120, Israel; radgonda@hotmail.com

doi: 10.1136/bjo.2003.041277

Accepted for publication 29 February 2004

Proprietary interest: none.

References

- 1 Nork TM, Millecchia LL. Treatment and histopathology of a congenital vitreous cyst. *Ophthalmology* 1998;**105**:825–30.
- 2 Lisch W, Rochels R. Pathogenesis of congenital vitreous cysts. *Klin Monatsbl Augenheilkd* 1989;**195**:375–8.
- 3 Awan KJ. Biomicroscopy and argon laser photostyloscopy of free-floating vitreous cysts. *Ophthalmology* 1985;**92**:1710–11.
- 4 Shields JA, Kline MW, Augsburger JJ. Primary iris cysts: a review of the literature and report of 62 cases. *Br J Ophthalmol* 1984;**68**:152.
- 5 Sallo FB, Haivani I. Recurring transitory blindness caused by primary marginal pigment epithelial iris cysts. *Am J Ophthalmol* 2002;**133**:407–9.
- 6 Goldberg MF. Persistent fetal vasculature (PFV): an integrated interpretation of signs and symptoms associated with persistent hyperplastic primary vitreous (PHPV). LIV Edward Jackson Memorial Lecture. *Am J Ophthalmol* 1997;**124**:587–626.

Central retinal artery occlusion and ophthalmoplegia following spinal surgery

Visual loss and ophthalmoplegia are very infrequent complications after spinal surgery.^{1–2} Visual loss may be caused by ischaemic optic neuropathy, central retinal artery or vein occlusion, or occipital stroke.^{2–3} Previous reports have attributed this complication to patient positioning, intraoperative blood loss, and controlled hypotension or shock.^{2–3} Associated risk factors include anaemia, prolonged surgical time, bradyarrhythmia, hypertension, diabetes, smoking, vascular disease, and increased blood viscosity.^{2–6} Ophthalmoplegia after spinal surgery is even more unusual than visual loss, and only few

reports exist in the literature.^{6–8} Moreover, magnetic resonance image (MRI) studies to differentiate between cavernous sinus thrombosis and direct compression of orbital contents have not been previously described. We therefore report two patients who developed this unusual combination of ophthalmoplegia and central retinal artery occlusion (CRAO) after spinal surgery.

Case 1

A 62 year old male ex-smoker underwent a L2–L3 posterior spinal decompression and segmental instrumentation for lumbar stenosis and scoliosis, in prone position with ocular protection (gauze swab and tape). The surgery lasted 2 hours and 45 minutes. Before the procedure blood pressure was 140/60 mm Hg and during operation it was maintained at 90/60 mm Hg. Just after surgery he complained of visual loss and left ocular and nasal pain. Examination revealed left palpebral oedema, local erythema, blindness, and total ophthalmoplegia of the left eye. Left pupil was dilated and fixed. The funduscopic examination showed retinal oedema, a central cherry-red spot at the macula, and attenuated arteries. The rest of his neurological examination was normal.

The haematocrit dropped from 43% to 34%. The brain MRI was normal and the orbit MRI revealed enlargement and hyperintensity of left ocular muscles in T2 weighted images sparing their tendons (fig 1). Ocular motility recovered in 4 weeks but visual loss persisted until the last follow up at 7 months.

Case 2

A 23 year old man with a history of tobacco abuse and asthma underwent a prolonged cervical arthrodesis in prone position caused by C7 vertebral collapse with spinal contusion. Immediately after surgery he complained of left visual loss and he was referred to our hospital. Details of duration of surgery, ocular protection, intraoperative blood pressure, and haematocrit were unavailable. Upon examination he showed blindness of the left eye with palpebral oedema, orbital pain, and total external ophthalmoplegia. The funduscopic examination revealed a pale retina with a macular cherry-red spot. The pupil was dilated and fixed. The MRI studies showed a normal brain, but swelling of the left extraocular muscles; MRI angiography and ophthalmic echo Doppler were normal. After 3 months

Figure 1 MRI of the orbit. T2 weighed image shows proptosis and oedema of extraocular muscles in the left eye, sparing their tendons.

Table 1 Reported patients with postoperative CRAO, ophthalmoplegia, or both

Author	Sex/age	Surgery	Position	Instrumentation	Operating time (minutes)	Blood pressure (mm Hg)	CRAO	Ophthalmoplegia
Slocum 1948 ⁹	F/55	Neurosurgical	Prone	No	180	80/60	Right	No
Givner 1950 ⁴	M/late 50s	Abdominal	Supine	No	265	Shock	Right	No
	F/33	Abdominal	Supine	No	85	Mild shock	Right	No
	F/37	Abdominal	Supine	No	90	Shock	Left	No
	F/40	Abdominal	Supine	No	Unknown	Unknown	Right	No
Gillan 1953 ⁵	M/26	Abdominal	Supine	No	At least 100	Systolic 70	Left	No
	F/48	Abdominal	Supine	No	At least 140	Systolic 55	Left	No
Hollenhorst 1954 ⁸	M/48	Spine	Prone	No	120	Normal	Left	No
	M/29	Neurosurgical	Sitting	No	120	Normal	Left	No
	M/65	Spine	Prone	No	120	Normal	Left	No
	M/50	Spine	Prone	No	180	90/60	Left	Yes
	M/39	Spine	Prone	No	180	Normal	Left	Yes
	F/36	Neurosurgical	Prone	No	240	Normal	Left	Yes
	M/50	Spine	Upright	No	240	Normal	Right	Yes
	M/29	Neurosurgical	Prone	No	360	92/55	Left	Yes
Bradish 1987 ¹⁰	F/12	Spine	Prone	Yes	180	Systolic 50	Left	No
West 1990 ⁷	F/50	Spine	Prone	Yes	270	Systolic 70–90	Left	Yes
Wolfe 1992 ⁶	F/28	Spine	Prone	Yes	135	90/60	Right	Yes
Grossman 1993 ¹³	F/12	Spine	Prone	Yes	360	Mean 70–80	Right	No
Halfon 2004	M/62	Spine	Prone	Yes	165	140/60–90/60	Left	Yes
	M/23	Spine	Prone	Yes	Unknown	Unknown	Left	Yes

CRAO, central retinal artery occlusion.

vision did not recover but ocular motility partially improved.

Comment

Our two patients developed complete ophthalmoplegia and CRAO after spinal surgery. Intraoperative ocular protection was used at least in the first patient. Common features included prone position and postoperative signs of orbital swelling. Imaging studies revealed signs of oedema in extraocular muscles sparing their tendons. To our knowledge, extraocular muscles abnormalities in MRI have not been previously reported. Ophthalmoplegia partially improved in one patient and fully recovered in the other within a few weeks, but visual loss persisted in both cases.

Stevens *et al.*² in a retrospective review of 3450 spinal surgeries, identified seven patients (incidence 0.2%) with visual loss caused by ischaemic optic neuropathy, occipital infarction, or central retinal vein occlusion, but neither presented with CRAO or extraocular muscle abnormalities.

Only few patients with CRAO following surgery have been previously described in the English literature (table 1). Moreover, Little¹ analysed 27 930 cases of controlled hypotension anaesthesia and found only three cases of CRAO. No details of surgical positioning or procedure were given. Since the original report of Slocum *et al.*,⁹ it has been emphasised that CRAO may be the result of extrinsic ocular pressure caused by head rest or anaesthetic mask malposition in the presence of hypotension, shock, and prolonged anaesthesia.^{4, 5, 8–10}

Ophthalmoplegia related to spinal surgery is also an exceptional complication. West *et al.*⁷ described a patient who developed unilateral total external ophthalmoplegia and unilateral visual loss following scoliosis surgery. An orbit computed tomography scan performed 1 week later showed left proptosis and swelling of the left medial rectus and ruled out cavernous sinus thrombosis. The authors suggested that perioperative ocular compression might be the cause. Wolfe *et al.*⁶ reported a young woman who suffered a

right CRAO after an instrumented spinal surgery. Postoperatively, the patient had blindness, moderate ptosis, restriction of the extraocular movements, and paraesthesias in the supraorbital region of the same eye. A CT scan of the brain and orbits showed only mild oedema of the right optic nerve. Ocular motility improved but the visual loss did not. Although prognosis is usually very poor, a recent patient described in Japan was treated successfully with urokinase and PGE1, stellate ganglion block, and hyperbaric oxygen therapy.¹¹

Hollenhorst *et al.*⁸ reported eight cases of unilateral visual loss after inadvertent orbital pressure during general anaesthesia for neurosurgical procedures. The most severe cases had proptosis, ptosis, and paralysis of extraocular musculature with no recovery. Moreover, they provoked visual loss and ophthalmoplegia by orbital compression for 60 minutes in seven rhesus monkeys, in the setting of hypovolaemia and hypotension. They proposed that partial or complete collapse of the arterial and venous channels of the orbit, occurred as a result of a tamponade action of the ocular contents. When the external pressure is released, the ischaemic vascular channels dilate and there is a transudation of fluid through the permeable walls into the tissue spaces. This results in orbital oedema, proptosis, paresis of ocular movement, and massive oedema of the retina. Based on findings from this animal model, it is likely that external pressure during the surgical procedure induces occlusion of the arterial and venous orbital vessels. The increased MRI signal in extraocular muscles is probably the expression of post-ischaemic oedema facilitated by the richer vascular supply and the more prominent extravascular space characteristic of these muscles.¹² Reversibility of ophthalmoplegia will probably depend on the degree of ischaemia suffered by both the extraocular muscles and the III, IV, and VI cranial nerves.

CRAO and ophthalmoplegia are unusual, but severe, complications after spinal surgery. Postoperative signs of orbital swelling, only in the affected eyes, were clear evidence of intraoperative compression in our patients.

The increase in intraorbital pressure associated with hypotension, shock, anaemia, prolonged operative time, and bradyarrhythmia are considered to be the main risk factors for developing CRAO and ophthalmoplegia. Adequate eye protection during surgery, and meticulous attention to keep the eyes free from pressure, can reduce the risk of these potentially avoidable serious complications.

Acknowledgement

We thank Dr John Stewart for his critical reading of the manuscript.

M J Halfon, P Bonardo, S Valensi,
M C Zaffaroni, M M Fernandez Pardal,
D Ribero Ayerza, R Ebner, P Anderson,
R C Reisin

Hospital Británico, Perdriel 74, Buenos Aires
C1280AEB, Argentina

Correspondence to: Dr Mario J Halfon, Hospital Británico, Perdriel 74, Buenos Aires C1280AEB, Argentina; mjhalfon@yahoo.com

doi: 10.1136/bjo.2003.039651

Accepted for publication 14 February 2004

References

- Little DM. Induced hypotension during anesthesia and surgery. *Anesthesiology* 1955;16:320–32.
- Stevens WR, Glazer PA, Kelley SD, *et al.* Ophthalmic complications after spinal surgery. *Spine* 1997;12:1319–24.
- Myers MA, Hamilton SR, Bogosian AJ, *et al.* Visual loss as a complication of spine surgery: a review of 37 cases. *Spine* 1997;12:1325–9.
- Givner I, Jaffe N. Occlusion of the central retinal artery following anesthesia. *Arch Ophthalmol* 1950;43:197–201.
- Gillan JG. Two cases of unilateral blindness following anesthesia with vascular hypotension. *Can Med Assoc J* 1953;69:294–6.
- Wolfe SW, Lospinus MF, Burke SW. Unilateral blindness as a complication of patient positioning for spinal surgery. A case report. *Spine* 1992;17:600–5.
- West J, Askin G, Clarke M, *et al.* Loss of vision in one eye following scoliosis surgery. *Br J Ophthalmol* 1990;74:243–4.
- Hollenhorst RW, Svien HJ, Benoit CF. Unilateral blindness occurring during anesthesia for

- neurosurgical operations. *Arch Ophthalmol* 1954;**52**:819–30.
- 9 **Slocum HC**, O'Neal KC, Allen CR. Neurovascular complications from malposition on the operating table. *Surg Gynecol Obstet* 1948;**86**:729–34.
 - 10 **Bradish CF**, Flowers M. Central retinal artery occlusion in association with osteogenesis imperfecta. *Spine* 1987;**12**:193–4.
 - 11 **Takeuchi M**, Morita K, Nakatsuka H, et al. A case of central retinal artery occlusion after anterior posterior fusion of the lumbar spine. *Masui* 2001;**50**:899–901.
 - 12 **Kaissar BA**, Kim JH, Bravo S, et al. Histologic basis for increased extraocular muscle enhancement in gadolinium-enhanced MR imaging. *Radiology* 1991;**179**:541–2.
 - 13 **Grossman W**, Ward WT. Central retinal artery occlusion after scoliosis surgery with a horseshoe headrest. Case report and literature review. *Spine* 1993;**18**:1226–8.

MAILBOX

Prophylaxis of ophthalmia neonatorum

I read with great interest the article by Isenberg *et al.*¹ The authors deserve to be commended for their pioneering interest in childhood blindness. There are certain points that I would like to clarify and supplement with regard to their study.

(1) The authors have mentioned that only babies born by a vaginal delivery were studied, since the eyes of babies delivered by caesarean section were previously proved to be nearly always sterile. This would result in a gross underestimation of the incidence of ophthalmia neonatorum in this study, for the following reason.

By convention, ophthalmia neonatorum is defined as conjunctivitis arising within 1 month after birth. Hence, some of these conjunctival infections could originate from sources other than the maternal vaginal and cervical flora. In fact, some cases of ophthalmia neonatorum, especially those caused by *Staphylococcus aureus*, could have originated at home, as previously reported by the authors themselves. In the same study, no significant difference in the frequency or type of infection was seen among the infants delivered vaginally or by caesarean section.²

Other authors, too have made similar observations. Krohn *et al.*³ have found some cases of ophthalmia neonatorum to have been acquired from the infants' nasopharyngeal passages or from their care givers after birth. Verma *et al.*⁴ in a prospective study from India, found no correlation between the microbiology of the conjunctival swabs of the infected eyes (*Staph aureus* was the commonest isolate) and the vaginal and cervical swabs of the mothers (*Escherichia coli* was the commonest isolate). They concluded that most of the cases of ophthalmia neonatorum were acquired postnatally. In the light of these previously reported studies, I feel that exclusion of cases that were delivered by caesarean section was not warranted and weakens the power of this study. The efficacy of the second drop of povidone-iodine was not tested on a significant proportion of the cases of ophthalmia neonatorum (those cases affecting the babies delivered by caesarean section).

(2) It would be relevant to note the percentage of ophthalmia neonatorum cases with neonatal dacryocystitis due to congenital nasolacrimal duct obstruction in this series. Such cases obviously would not have

benefited from a second drop of povidone-iodine.

(3) The Indian study by Verma *et al.*⁴ found a seasonal incidence of ophthalmia neonatorum with two peaks: in February and May–June, in most probability as a result of the monsoon and the hot tropical climate with the attendant eye seeking flies. It would be interesting to know whether any such seasonal variation was noted in this study, which was carried out in Kenya, a country with a hot tropical climate, like India. If so, the efficacy of the second drop could be investigated again during such peaks.

V Vedantham

Correspondence to: Vasumathy Vedantham, Retina-Vitreous Service, Aravind Eye Hospital and Postgraduate Institute of Ophthalmology, 1 Anna Nagar, Madurai - 625 020, Tamilnadu, India; drvasumathy@yahoo.com

doi: 10.1136/bjo.2004.042846

Accepted for publication 22 January 2004

References

- 1 **Isenberg SJ**, Apt L, Signore MD, et al. A double application approach to ophthalmia neonatorum prophylaxis. *Br J Ophthalmol* 2003;**87**:1449–52.
- 2 **Isenberg SJ**, Apt L, Wood M. A controlled trial of povidone-iodine as prophylaxis against ophthalmia neonatorum. *N Engl J Med* 1995;**332**:562–6.
- 3 **Krohn MA**, Hillier SL, Bell TA, et al. The bacterial etiology of conjunctivitis in early infancy. *Am J Epidemiol* 1993;**138**:326–32.
- 4 **Verma M**, Chhatwal J, Varghese PV. Neonatal conjunctivitis: a profile. *Indian Pediatr* 1994;**31**:1357–61.

Authors' reply

We greatly appreciate the inquiry of Dr Vedantham and are happy to reply to his questions.

While he is correct regarding the definition of ophthalmia neonatorum including all infections acquired by an infant during the first 30 days of life, for the purposes of our study,¹ we were primarily interested in those cases resulting from neonatal exposure in the birth canal. This source of ophthalmia neonatorum is the one that would be influenced mainly by a second drop of povidone-iodine placed later on the day of birth. Infections postnatally that Dr Vedantham listed as arising, while technically still within the definition of "ophthalmia neonatorum," would not be impacted by this second drop and therefore would not be directly affected by this study. Indeed, Dr Vedantham's interest in neonatal dacryocystitis would also fall within the same question since the reflux from the tear duct causing this infection generally does not arise until well after the first day of life.

The proportion of ophthalmia neonatorum cases acquired postnatally compared with those acquired during the birth process probably differs by country. The difference in predominant causative organisms (*Staphylococcus aureus* and *Enterococcus* in the former situation as in India² v chlamydia and gonococcus in the latter) probably reflects the origin of the infection. Historically, in Kenya a high proportion of the infections probably arose from the birth process as reflected in the type of infecting organism. Thus, our Kenyan study, was primarily directed towards those infections acquired during birth.

The fact that ophthalmia neonatorum in some countries tends to peak in certain seasons and not in others should cause increased vigilance for this disorder in those countries. Our study, however, encompassed more than one full year of births in Kikuyu, Kenya. Therefore, the study included both the peaks and troughs of the incidence of ophthalmia neonatorum.

We thank Dr Vedantham for his interest and hope that in many countries, including India, ophthalmia neonatorum prophylaxis will either continue unabated or be initiated preferentially with the use of povidone-iodine.

S J Isenberg, L Apt

UCLA School of Medicine, Harbor-UCLA Medical Center, 1000 W Carson Street, Box 6, Torrance, CA 90509, USA

Correspondence to: S J Isenberg, UCLA School of Medicine, Harbor-UCLA Medical Center, 1000 W Carson Street, Box 6, Torrance, CA 90509, USA; isenberg@ucla.edu

doi: 10.1136/bjo.2004.045096

Accepted for publication 19 February 2004

References

- 1 **Isenberg SJ**, Apt L, Del Signore M, et al. A double application approach to ophthalmia neonatorum prophylaxis. *Br J Ophthalmol* 2003;**87**:1449–52.
- 2 **Verma M**, Chhatwal J, Varghese PV. Neonatal conjunctivitis: a profile. *Indian Pediatr* 1994;**31**:1357–61.

Retinectomy for intractable glaucoma

We read with interest the article by Joussem *et al* on the long term results of retinectomy for the treatment of intractable glaucoma.¹ We congratulate the authors for studying this innovative method for the management of refractory glaucoma with a long follow up of 5 years.

The high incidence of complications in the study, however, has aroused our concern as only 15.9% of patients completed the study uneventfully. Further vitreoretinal surgeries were required in 47.7% because of retinal complications. Moreover, the incidence of hypotony, phthisis, and enucleation was 25%, 20%, and 16% respectively and these figures are higher compared with other treatments such as glaucoma implants and cyclodiode. We have previously studied the use of the Ahmed valve implant for complicated glaucoma and hypotony, phthisis, and enucleation occurred in 10.8%, 3.1%, and 1.5%, respectively.² A recent study on the management of refractory glaucoma by cyclodiode similarly found a lower rate of hypotony and phthisis of 9.5% and 5.3% respectively.³ The high complication rates in the study by Joussem *et al* may be because of the negative case selection with a high incidence of aphakic (30%) and infantile and juvenile glaucoma (7%). A further controlled study comparing retinectomy with other treatments may therefore be warranted.

In this evidence based era, emphasis should be placed on outcomes which are "patient oriented evidence that matters" (POEMs).⁴ It was stated by the authors that the main intentions of the surgery were to relieve pain and to preserve the eye without discomfort. Unfortunately, these POEMs were not included in the final outcome measures. Instead, success was determined

by "disease oriented evidence" (DOE) like intraocular pressure and retinal attachment, which are surrogate outcomes.⁵ These DOEs may not correlate well with the patients' symptoms and it would be valuable if the authors can include the level of pain and discomfort as other outcome measures for the study.

V W Y Wong, T Y Y Lai, D S C Lam
Hong Kong Eye Hospital, 147K Argyle Street,
Kowloon, Hong Kong

Correspondence to: Dr Timothy Y Y Lai, Hong Kong Eye Hospital, 147K Argyle Street, Kowloon, Hong Kong; tyylai@netvigator.com

doi: 10.1136/bjo.2004.042820

Accepted for publication 22 January 2004

References

- 1 Jousen AM, Walter P, Jonescu-Cuyper CP, et al. Retinectomy for treatment of intractable glaucoma: long term results. *Br J Ophthalmol* 2003;**87**:1094-103.
- 2 Lai JS, Poon AS, Chua JK, et al. Efficacy and safety of the Ahmed glaucoma valve implant in Chinese eyes with complicated glaucoma. *Br J Ophthalmol* 2000;**84**:718-21.
- 3 Murphy CC, Burnett CA, Spry PG, et al. A two centre study of the dose-response relation for transscleral diode laser cyclophotocoagulation in refractory glaucoma. *Br J Ophthalmol* 2003;**87**:1252-7.
- 4 Shaughnessy AF, Slawson DC, Bennett JH. Becoming an information master: a guidebook to the medical information jungle. *J Fam Pract* 1994;**39**:489-99.
- 5 Temple R. Are surrogate markers adequate to assess cardiovascular disease drugs? *JAMA* 1999;**282**:790-5.

Blue light and the circadian clock

Drs Mainster and Sparrow have provided an excellent perspective on the relative merits and difficulties of extending intraocular lens (IOL) absorption into the blue portion of the spectrum.¹

However, they have not considered an unintentional consequence of blockage of the blue portion of the spectrum—reducing the activity of intrinsically photosensitive retinal ganglion cells.²⁻³ These cells subserve several non-visual ocular photoreceptive tasks, most prominently the entrainment of the circadian clock to external light-dark cycles.⁴ Pupillary light responses in mice are also at least partially controlled by this system, which appears to use a novel opsin (melanopsin)⁵⁻⁶ and possibly also a flavoprotein (cryptochrome)⁷⁻⁸ as photopigments.

Experiments in mice have suggested that the action spectrum for these photopigments peak in the blue, at approximately 480 nm, but with substantial sensitivity to blue light to 430 nm.⁹ This system appears to be functional in humans as documented by the action spectrum for light suppression of the pineal hormone, melatonin.¹⁰⁻¹¹

The clinical importance of these photoreceptors is presently unknown, although it appears that loss of retinal ganglion cells predisposes children and young adults to disorders of sleep timing that outer retinal disease does not.¹² While, as the authors note, there may be substantial benefit in blocking blue light phototoxicity, particularly for patients with pre-existing outer retinal degeneration, these lenses may have unintended consequences with respect to the

timing of sleep and wakefulness or levels of certain neurohormones.

R N Van Gelder

Washington University Medical School, Campus Box 8096, 660 S Euclid Avenue, St Louis, MO 63110, USA; vangelder@vision.wustl.edu

doi: 10.1136/bjo.2004.042861

Accepted for publication 22 January 2004

References

- 1 Mainster MA, Sparrow JR. How much blue light should an IOL transmit? *Br J Ophthalmol* 2003;**87**:1523-9.
- 2 Berson DM. Strange vision: ganglion cells as circadian photoreceptors. *Trends Neurosci* 2003;**26**:314-20.
- 3 Berson DM, Dunn FA, Takao M. Phototransduction by retinal ganglion cells that set the circadian clock. *Science* 2002;**295**:1070-3.
- 4 Freedman MS, Lucas RJ, Soni B, et al. Regulation of mammalian circadian behavior by non-rod, non-cone, ocular photoreceptors. *Science* 1999;**284**:502-4.
- 5 Panda S, Provencio I, Tu DC, et al. Melanopsin is required for non-image-forming photic responses in blind mice. *Science* 2003;**301**:525-7.
- 6 Hattar S, Lucas RJ, Mrosovsky N, et al. Melanopsin and rod-cone photoreceptive systems account for all major accessory visual functions in mice. *Nature* 2003;**424**:75-81.
- 7 Van Gelder RN, Wee R, Lee JA, et al. Reduced pupillary light responses in mice lacking cryptochromes. *Science* 2003;**299**:222.
- 8 Selby CP, Thompson C, Schmitz TM, et al. Functional redundancy of cryptochromes and classical photoreceptors for nonvisual ocular photoreception in mice. *Proc Natl Acad Sci USA* 2000;**97**:14697-702.
- 9 Lucas RJ, Douglas RH, Foster RG. Characterization of an ocular photopigment capable of driving pupillary constriction in mice. *Nat Neurosci* 2001;**4**:621-6.
- 10 Brainard GC, Hanifin JP, Greeson JM, et al. Action spectrum for melatonin regulation in humans: evidence for a novel circadian photoreceptor. *J Neurosci* 2001;**21**:6405-12.
- 11 Thapan K, Arendt J, Skene DJ. An action spectrum for melatonin suppression: evidence for a novel non-rod, non-cone photoreceptor system in humans. *J Physiol (Lond)* 2001;**535**:261-7.
- 12 Wee R, Van Gelder RN. Sleep disturbances in young subjects with visual dysfunction. *Ophthalmology* (in press).

Author's reply

I appreciate Van Gelder's thoughtful comments regarding the potential consequences of a ultraviolet + blue light absorbing intraocular lens (IOL) on circadian rhythmicity. I agree that the clinical importance of retinal ganglion photoreceptors is currently unknown and that decreasing the amount of blue light reaching them might affect their function. Conversely, if photosensitive ganglia respond to circadian changes in their blue light exposure rather than just the magnitude of that exposure, a ultraviolet + blue light absorbing IOL may not impair ganglion function.

Van Gelder re-emphasises our finding that IOL chromophore selection balances the potential loss of useful visual function against a reduction in the risk of acute ultraviolet-blue phototoxicity. Our paper did not state, however, that ultraviolet + blue absorbing IOLs were desirable for people with outer retinal degeneration. Indeed, blue light is more important in scotopic than photopic vision. Individuals with age related macular degeneration have greater night-time visual problems than their peers without it, and these scotopic problems may be exacerbated

if a significant amount of blue light is blocked by an IOL.

M A Mainster

University of Kansas Medical School, 3901 Rainbow Boulevard, MS3009, Kansas City, KS 66160-7379, USA; mmmainste@kumc.edu

doi: 10.1136/bjo.2004.045120

Accepted for publication 19 February 2004

Modification of classification of ocular chemical injuries

A recent paper by Kobayashi and co-workers¹ on temporary amniotic membrane patching for acute chemical burns highlights the difficulty in the consistent classification of this type of injury.

Roper-Hall's classification of acute chemical injuries to the eye is based on the original classification of Ballen²⁻³ and there is little difference between them. However, in neither classification is the grade based on the most severe sign. This immediately leads to the problem of trying to classify an eye having, for example, corneal signs of one grade and conjunctival signs of another. There is also difficulty in grading the conjunctival and limbal injuries.

Dua *et al*⁴ recognised the problem of assessing limbal stem cell damage and proposed a quasi-analogue scale in order to incorporate intergrade variations. They suggested using limbal fluorescein staining as a marker of limbal stem cell damage. However, their grading scheme is complex and departs significantly from that of Roper-Hall. Although fluorescein staining is a useful sign, it does not necessarily imply limbal stem cell damage or failure and has not been shown to be a better index of limbal damage than perilimbal ischaemia. Importantly, it is also becoming apparent, that both the fornices and mucocutaneous junction of the conjunctiva are important for conjunctival regeneration.⁵⁻⁷ In fact, Roper Hall stressed the importance of involvement of contiguous areas of the conjunctiva, which may lead to symblepharon formation.⁴

Although there are limitations with Roper-Hall's classification, it is simple and remains popular. Rather than replace Roper-Hall's and Ballen's classification,²⁻³ we suggest a modification, which addresses some of the issues raised and makes the classification more robust.

One of the questions that needs to be answered is whether to base the grade of injury upon the most severe sign or on a combination of ocular surface signs. A combination of signs using three parameters (cornea, limbus, and conjunctiva), each with three levels requires 27 possible combinations to avoid crossover. To avoid this complexity and without evidence to indicate a difference in prognosis, it would seem reasonable therefore to base the grade of injury on the most severe sign.

Although limbal ischaemia does not necessarily imply limbal stem cell failure, it remains to be shown that it is less indicative than fluorescein limbal staining of limbal damage. We propose therefore to continue to use limbal ischaemia in the grading of injury.

With regard to conjunctival involvement, in order to be able to include the total area of involvement, we suggest dividing the conjunctival surface into bulbar and tarsal areas, as is natural. The bulbar and tarsal conjunctiva comprise approximately two thirds and

Figure 1 Estimation of conjunctival injury. For example, $1/6\text{th} + 1/6\text{th} = 1/3\text{rd}$.

one third of the total conjunctival surface respectively. Using conjunctival fluorescein staining as an indicator of the extent of conjunctival damage, the area of involvement can be based on the fraction of the third involved, limiting any division into not less than sixths—that is, the tarsal surfaces together comprise a third of conjunctiva (see fig 1). This includes the issue that a vertical distribution of conjunctival injury is as important as a horizontal distribution.

Corneal involvement in terms of prognosis remains an area of difficulty. Although it may be assumed that limbal and conjunctival damage implies a worse prognosis than isolated corneal damage, this has yet to be shown. In addition, a severe chemical injury involving the cornea but not the limbus, or vice versa, would be expected to be an uncommon event. We therefore propose to retain the degree of corneal damage (as proposed by Roper-Hall² and Ballen³) in grading of the injury (see table 1).

Thus, grade I is identified by any isolated corneal epithelial injury. Grade II includes limbal or conjunctival involvement, but involves less than one third of the area involved. Grade III includes either a hazy cornea, defined as obscurity of the iris or pupil details (as per Roper-Hall's and Ballen's original descriptions), and/or greater than one third of limbal or conjunctival damage. With the advent of recent surgical techniques such as amniotic membrane transplants and limbal allografts, the prognosis of more severe ocular chemical injuries previously classified as Roper-Hall grade IV have improved and no longer carry a uniformly poor prognosis.² Therefore, we reason that these cases can be included in grade III of our proposed classification.

In conclusion, in the absence of good evidence for re-classifying ocular surface injuries, it would seem reasonable to keep

to the Roper-Hall/Ballen classification and to move it forward by addressing the weaknesses of that system. We hope that the proposed grading system improves the consistency with which chemical injuries are reported in the literature, serves as a basis for controlled comparative evaluation of modern treatment, and stimulates further work in this area.

S Harun, S Srinivasan, K Hollingworth, M Batterbury, S Kaye

8z Link, Royal Liverpool University Hospital, Prescot Street, Liverpool L7 8XP, UK

Correspondence to: Mr Shabbir Harun, 8z Link, Royal Liverpool University Hospital, Prescot Street, Liverpool L7 8XP, UK; harun@doctors.org.uk

doi: 10.1136/bjo.2004.046797

Accepted 15 March 2004

References

- 1 Kobayashi A, Shirao Y, Yoshita T, *et al*. Temporary amniotic membrane patching for acute chemical burns. *Eye* 2003;17:149–58.
- 2 Roper-Hall MJ. Thermal and chemical burns. *Trans Ophthalmol Soc UK* 1965;85:631–53.
- 3 Ballen PH, Hemstead NY. Treatment of chemical burns of the eye. *Eye, Ear, Nose and Throat Monthly* 1964;43:57–61.
- 4 Dua HS, King AJ, Joseph A. A new classification of ocular surface burns. *Br J Ophthalmol* 2001;85:1379–83.
- 5 Wirtschafter JD, Ketcham JM, Weinstock RJ, *et al*. Mucocutaneous junction as the major source of replacement palpebral conjunctival epithelial cells. *Invest Ophthalmol Vis Sci* 1999;40:3138–46.
- 6 Wei ZG, Wu RL, Lavkar RM, *et al*. In vitro growth and differentiation of rabbit bulbar, fornix and palpebral conjunctival epithelia. Implications on conjunctival epithelial transdifferentiation and stem cells. *Invest Ophthalmol Vis Sci* 1993;34:1814–28.
- 7 Brodovsky SC, McCarty CA, Snibson G, *et al*. Management of alkali burns: an 11 year retrospective review. *Ophthalmology* 2000;107:1829–35.

Author's response

The response by Harun *et al* is to be commended in so far as it highlights the problems with the current Roper-Hall classification system and the difficulties it poses in evaluating outcome and efficacy of treatments in ocular surface burns. As a proposed modification, however, it is a retrograde step.

The three major issues with the Roper-Hall¹ classification were that it lumped all injuries with 50% or more of limbal involvement into one category, did not take into account conjunctival involvement in the actual classification, and placed undue emphasis on the degree of corneal haze.

The proposed modification by Harun *et al* goes a step backwards by grouping all injuries with more than 33% limbal involvement into

one category. The grading of a patient with all 12 clock hours of limbus involvement would then be the same as one with just over 3 clock hours of limbus involvement! The prognosis given to these two patients cannot be the same, given that the Roper-Hall and the Dua, King, and Joseph² classifications are prognostic classifications. Furthermore, a patient presenting with less than one third limbus involvement does not necessarily come with less than one third conjunctival involvement, which could be much more. The proposed modification does not allow for such variances, which are frequent. The Dua, King, and Joseph classification has the flexibility to allow for such variables and also to progressively document change both improvement and deterioration, over the acute phase of the injury. The authors rightly point out that the degree of ischaemia does not always correspond to degree of limbal involvement. Yet limbal involvement without ischaemia, in the form of loss of stem cells, can have an equally important impact on prognosis. That is precisely why the Dua, King, and Joseph classification considers limbal involvement (to encompass ischaemia as well) rather than limbal ischaemia alone.

The point about conjunctival involvement is well made in the proposed modification. This does not differ significantly from the Dua, King, and Joseph classification. The latter was the first to take this aspect of burns into account in determining severity and prognosis. The authors mention the importance of tarsal conjunctival involvement. This is a valid though often an impractical consideration. Associated swelling, induration, thickening, shrinkage and the like, of the lids make tarsal conjunctival evaluation impractical if not impossible in some cases, in the immediate post-injury period. It was for this practical consideration that the Dua, King, and Joseph classification included only the extent of bulbar conjunctival involvement in determining the grade. It is interesting to note that the authors disregard limbal fluorescein staining as an indicator of limbal damage (as proposed in the Dua, King, and Joseph classification) but propose fluorescein staining as an indicator of conjunctival damage in evaluating extent of conjunctival damage. This implies that fluorescein staining is appropriate to evaluate both conjunctival epithelial damage and conjunctival ischaemia but not limbal epithelial damage and limbal ischaemia. There is no rationale for this.

Corneal haze can be an indicator of the offending chemical rather than the severity of the insult. It is not uncommon to find a clear and transparent cornea, which is totally denuded of its epithelium, immediately after a chemical injury. This can stay so for a few days before becoming rapidly hazy or opaque, or remain clear and become re-epithelialised. Corneal endothelial damage leading to stromal oedema and haze can occur later in the course of an acute chemical injury. Conversely, a hazy cornea with a resultant scar could do well following a corneal graft procedure if the limbal involvement is minimal. The proposed modification retains corneal haze as a grading parameter and includes a hazy cornea in grade 3 only. There are many chemical injuries, which involve 3–6 clock hours of the limbus (30–50%) with a clear cornea. These do not fall well in any grade in the proposed new classification and highlight the inherent

Table 1 Modified classification of ocular chemical injuries

Grade	Cornea	Limbal ischaemia	Conjunctival involvement	Prognosis
I	Clear: epithelial damage only	None	None	Good
II	Clear: epithelial damage only	<1/3	<1/3	Good
III	Hazy cornea	OR >1/3	OR >1/3	Guarded

problem in the Roper-Hall classification and its proposed modification.

Most important of all, the proposed classification is purely theoretical and has not been validated. The Dua, King, and Joseph classification is based on several years of clinical experience of managing burns including more than 67 patients. It is simple and easy to use (clock hours of limbus involvement and percentage of conjunctival involvement), flexible, and allows for all combinations of different extents of involvement of the two structures. It is validated as a prognostic indicator and allows for accurate comparison of cases. The proposed new classification/modification fails on all these counts.

H S Dua

Division of Ophthalmology and Visual Sciences, B Floor, Eye ENT Centre, University Hospital, Queens Medical Centre, Derby Road, Nottingham NG7 2UH, UK; harminder.dua@nottingham.ac.uk

doi: 10.1136/bjo.2004.046797

Accepted 23 March 2004

Further communications regarding this subject can be found at eLetters on the BJO website (www.bjophthalmol.com)

References

- 1 Roper-Hall MJ. Thermal and chemical burns. *Trans Ophthalmol Soc UK* 1965;**85**:631-53.
- 2 Dua HS, King AJ, Joseph A. New classification of ocular surface burns. *Br J Ophthalmol* 2001;**85**:1379-83.

BOOK REVIEW

Complications in Ophthalmic Plastic Surgery

Ed by B G Brazzo. Pp 264; £130.50 (€169.95). New York: Springer Verlag, 2003. ISBN 0-387-00283-9.

There is no doubt that complications in surgery are an inevitable fact.

Nevertheless, the unforeseen surgical outcomes always play a fundamental part in the self improving process. Certainly, the experience provides the safest way, for both the patient and the surgeon, to prevent harm and smooth the final result.

This book represents one of the most serious, and not very common, works focusing on the complications in ophthalmic plastic surgery. It is a considerably well organised book, which apparently requires some basic knowledge of oculoplastics and facial aesthetic surgery. The format is based on three distinguished parts: cosmetic surgery, ptosis, and lower eyelid malposition. A

very competent number of contributors cover the topics of their specific interest. In the first part the authors are dealing with the blepharoplasties, the laser resurfacing, and the forehead lift. The ptosis chapter is referred to the most common ptosis techniques but brow suspension is remarkably absent. The third part, although it is entitled "Lower eyelid malposition," includes and some unrelated, though welcomed topics, like DCR, enucleation, and orbital fractures.

The necessity of the communication between the surgeon and the patient is vigorously emphasised and didactically analysed in every single chapter. Deep understanding of the patient's expectations as well as detailed information about the pragmatic results is recommended throughout the chapters of the book. There is quite a sufficient reference to preoperative evaluation of the patient regarding measurements, anaesthesia, and surgical preparation.

Although the covered operations are extensively described, a countable number of other surgical techniques, and their possible complications, are not mentioned. The latter is probably related to the editor's orientation to aesthetic oculoplastic surgery.

The anatomical and pathophysiological mechanisms of the most common complications are thoroughly explained. At the same time, the authors give many enlightening tips, based on their broad experience, for preventing the problems, and meticulously describe the management of the intraoperative and postoperative complications. The number of the illustrations do not adequately correspond to the addressed complications and the quality of the pictures varies, depending on the author's collection. Additionally, the shortage of references in some of the most interesting chapters (ptosis, enucleation) should certainly not be overlooked, for the magnitude of such a book.

Every attempt to give precious advice about the frustrating and unavoidable surgical complications is always warmly welcomed. Brian Brazzo's book is predominately a useful guide to the understanding, prevention, and management of the commonest problems in oculoplastic surgery. Despite the expected problems of every first edition this generally represents a meticulous work on specific issues and thus is recommended for the ophthalmic surgeon and especially for surgeons who are chiefly interested in oculoplastics and cosmetic surgery.

N Chalvatzis

Bristol Eye Hospital, Lower Maudlin Street,
Bristol BS1 2LX, UK;
nikocha@hotmail.com

CORRECTION

In the letter titled Sequential treatment of central retinal vein occlusion with intravitreal

tissue plasminogen activator and intravitreal triamcinolone (*Br J Ophthalmol* 2004;**88**:1100-1101) the authors were listed incorrectly. The correct listing is as follows: J M Lahey, J J Kearney, M C Cheung. The journal apologises for this error.

It has come to our attention that, owing to a production error, the letter by N Islam, K Mireskandari, and G E Rose has been published twice, in the June issue (*Br J Ophthalmol* 2004;**88**:833-834) and the August issue (*Br J Ophthalmol* 2004;**88**:1092-1093). The initial publication in the June issue should be taken as the article of record. We apologise to the authors and readers for any confusion this accidental duplicate publication may have caused.

NOTICES

Childhood cataract

The latest issue of Community Eye Health (No 50) deals with the manitude, management, economics and impact of childhood cataract. For further information please contact: Journal of Community Eye Health, International Resource Centre, International Centre for Eye Health, Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK (tel: +44 (0)20 7612 7964; email: Anita.Shah@lshtm.ac.uk; online edition: www.jceh.co.uk). Annual subscription (4 issues) UK £28/US\$45. Free to developing country applicants.

Ophthalmic Anesthesia Society

The 18th annual meeting of the Ophthalmic Anesthesia Society will be held on 1-3 October 2004 in Chicago, USA. For further details: Ophthalmic Anesthesia Society (OAS), 793-A Foothill Blvd, PMB #119, San Luis Obispo, CA 93405, USA (tel: 001 805 534 0300; fax: 001 805 534 9030; email: info@eyeanesthesia.org; website: www.eyeanesthesia.org).

4th International Congress on Autoimmunity

The 4th International Congress on Autoimmunity will take place 3-7 November 2004 in Budapest, Hungary. Further details: Kenes International Global Congress Organisers and Association Management Services, 17 Rue du Cendrier, PO Box 1726, CH-1211 Geneva 1, Switzerland (tel: +41 22 908 0488; fax: +41 22 732 2850; email: autoim04@kenes.com; website: www.kenes.com/autoim2004).