Highlights from this issue

doi:10.1136/bjophthalmol-2014-306207

Keith Barton, James Chodosh, Jost Jonas, *Editors in Chief*

In this month's issue, Nash *et al*¹ continue their series of Statistics notes on behalf of the Ophthalmic Statistics Group. In this article they discuss methods of analysing continuous variables of clinical trial data baseline and post-treatment.

In a systematic review and meta-analysis of trials of topical antibiotics for bacterial keratitis using Cochrane methodology, McDonald *et al*² report no evidence of difference in comparative effectiveness between fluoroquinolones and aminoglycoside-cephalosporin combinations, although there they observed reported differences in the safety profile.

Oh *et al*³ describe and validate the utility of the Microsoft Kinect-based head tracker for measuring head posture using the cervical range of motion as a reference. They observed that measurements of head posture using the Kinect head-tracker were very close to those of the cervical range of motion with relatively favourable test-retest reliability.

Serna-Ojeda and Pedroza-Seres⁴ report the clinical course of 49 patients in Mexico with pars planitis who presented at median age 8 years and underwent immunosuppressive therapy with clinical and visual improvement and reduction in the steroid dose.

Ishikawa *et al*⁵ report the usefulness of measuring choroidal thickness by enhanced depth imaging optical coherence tomography in evaluating uveitic activity in patients with Behçet's disease.

Nowinska *et al*⁶ observed in 24 Polish patients with phenotypically homogenous macular corneal dystrophy on confocal microscopy and optical coherence tomography, that the genetic mutations were quite heterogenous on analysis of single nucleotide polymorphisms.

Tan *et al*⁷ propose a classification of polypoidal choroidal vasculopathy based on indocyanine green angiography and fluorescein angiography characteristics, using 107 treatment naïve patients. The subtypes that derive from this classification were observed to have prognostic significance.

In a case control study of single nucleotide polymorphisms to investigate the association of Complement Factor I and CD46 genes with acute anterior uveitis (AAU), Wang *et al*⁸ examined 300 AAU patients and 300 healthy controls and report the Rs7356506 in the CF1 gene to be protective against AAU. They conclude that CF1 is likely to be involved in the pathogenesis of AAU.

REFERENCES

- 1 Nash R, Bunce C, Freemantle N, *et al*. Ophthalmic Statistics Note 4: analysing data from randomised controlled trials with baseline and follow-up measurements. *Br J Ophthalmol* 2014;98:1467–9.
- 2 McDonald EM, Ram FSF, Patel DV, *et al*. Topical antibiotics for the management of bacterial keratitis: an evidence-based review of high quality randomised controlled trials. *Br J Ophthalmol* 2014;98:1470–7.
- 3 Oh B-L, Kim J, Kim J, *et al*. Validity and reliability of head posture measurement using Microsoft Kinect. *Br J Ophthalmol* 2014;98:1560–4.
- 4 Serna-Ojeda JC, Pedroza-Seres M. Treatment with immunosuppressive therapy in patients with pars planitis: experience of a reference centre in Mexico. *Br J Ophthalmol* 2014;98:1503–7.
- 5 Ishikawa S, Taguchi M, Muraoka T, *et al*. Changes in subfoveal choroidal thickness associated with uveitis activity in patients with Behçet's disease. *Br J Ophthalmol* 2014;98:1508–13.
- 6 Nowinska AK, Wylegala E, Teper S, *et al*. Phenotype and genotype analysis in patients with macular corneal dystrophy. *Br J Ophthalmol* 2014;98:1514–21.
- 7 Tan CSH, Ngo WK, Lim LW, *et al*. A novel classification of the vascular patterns of polypoidal choroidal vasculopathy and its relation to clinical outcomes. *Br J Ophthalmol* 2014;98:1528–33.
- 8 Wang Y, Huang X-F, Yang M-M, *et al*. CFI-rs7356506 is a genetic protective factor for acute anterior uveitis in Chinese patients. *Br J Ophthalmol* 2014;98:1592–1596.

CrossMark