

e-Table 11: Change in microscotomas in center 3 and center 4

| | | Center=03 N=17 | Center=04 N=17 | Total N=34 | p-value |
|--|--------------|-------------------|-------------------|-------------------|---------|
| Number of microscotomas at D0 | | | | | |
| | N | 16 | 17 | 33 | |
| | Min-Max | [0.00;29.00] | [0.00;25.00] | [0.00;29.00] | 0.4936 |
| | Mean+/- SD | 12.31+/-12.92 | 7.24+/-8.28 | 9.70+/-10.92 | |
| | Median [IQR] | 7.00[0.00;27.50] | 3.00[1.00;12.00] | 3.00[0.00;20.00] | |
| Number of microscotomas at M06 | | | | | |
| | N | 10 | 15 | 25 | |
| | Min-Max | [0.00;22.00] | [0.00;21.00] | [0.00;22.00] | |
| | Mean+/- SD | 5.10+/-8.31 | 5.13+/-6.16 | 5.12+/-6.93 | |
| | Median [IQR] | 1.00[0.00;4.00] | 2.00[1.00;9.00] | 2.00[0.00;8.00] | |
| Change in number of microscotomas between D0 and M06 | | | | | |
| | N | 10 | 15 | 25 | |
| | Min-Max | [-29.00;4.00] | [-25.00;9.00] | [-29.00;9.00] | 0.6282 |
| | Mean+/- SD | -7.50+/-11.42 | -2.87+/-9.16 | -4.72+/-10.16 | |
| | Median [IQR] | -4.50[-7.00;0.00] | -1.00[-6.00;2.00] | -1.00[-7.00;0.00] | |