

Supplementary Table 3. Changes in mean BCVA and CRT from baseline over time in pseudophakic, treatment-naïve patients (n=39)

| Visit | Mean BCVA (SD), letters | Change in mean BCVA (SD) from baseline, letters | p value | Mean CRT (SD), µm | Change in mean CRT (SD) from baseline, µm | p value |
|---------------|----------------------------|---|------------|-------------------------|--|------------|
| Baseline n | 58.6 (17.7) 39 | – | – | 416.9 (100.8) 39 | – | – |
| Week 6 n | 63.7 (18.8) 29 | +3.6 (8.9) | 0.041 | 321.2 (60.2) 32 | –85.4 (62.5) | <0.001 |
| Week 16 n | 60.3 (19.3) 34 | +3.2 (8.8) | 0.044 | 366.1 (89.1) 34 | –54.2 (78.8) | <0.001 |
| Week 24 n | 58.2 (18.5) 23 | +1.6 (11.8) | 0.521 | 360.5 (98.6) 26 | –71.5 (99.3) | 0.001 |
| Week 52 n | 65.3 (16.7) 29 | +4.1 (11.3) | 0.063 | 336.7 (94.8) 29 | –91.5 (115.0) | <0.001 |

BCVA, best-corrected visual acuity; CRT, central retinal thickness; SD, standard deviation.