

e - Table 6: Change in peripapillary retinal nerve fiber layer thickness during the follow-up between active peeling and no-peeling group

pRNFL: peripapillary retinal nerve fiber layer thickness (μm); *: statistically significant

All patients	Difference	Confidence interval (95%)	p-value
Temporal superior pRNFL			
M1 : ILM peeling versus No ILM peeling group	-0.78	[-9.13 ; 7.57]	0.8530
M6 : ILM peeling versus No ILM peeling group	-7.15	[-14.50 ; 0.20]	0.0565
M12: ILM peeling versus No ILM peeling group	-7.06	[-14.79 ; 0.67]	0.0729
Temporal pRNFL			
M1 : ILM peeling versus No ILM peeling group	2.17	[-7.60 ; 11.94]	0.6596
M6 : ILM peeling versus No ILM peeling group	-11.79*	[-20.47 ; -3.10]	0.0083
M12: ILM peeling versus No ILM peeling group	-10.37*	[-18.54 ; -2.20]	0.0135
Temporal inferior pRNFL			
M1 : ILM peeling versus No ILM peeling group	2.43	[-5.01 ; 9.87]	0.5182
M6 : ILM peeling versus No ILM peeling group	0.48	[-8.84 ; 9.80]	0.9190
M12: ILM peeling versus No ILM peeling group	1.34	[-7.50 ; 10.18]	0.7642
Mean temporal + temporal superior + temporal inferior pRNFL			
M1 : ILM peeling versus No ILM peeling group	-0.56	[-5.98 ; 4.86]	0.8382
M6 : ILM peeling versus No ILM peeling group	-7.72*	[-13.96 ; -1.74]	0.0119
M12: ILM peeling versus No ILM peeling group	-6.60*	[-12.59 ; -0.61]	0.0311