

e-Table 9: Baseline Characteristics of randomized Patients by centers

BCVA: best-corrected visual acuity; SD: standard deviation

	Centre1 N=31	Centre2 N=28	Centre3 N=17	Centre4 N=17	Centre5 N=7	Total N=100
Female N(%)	18 (58.06%)	17 (60.71%)	10 (58.82%)	9 (52.94%)	5 (71.43%)	59 (59.00%)
Age Mean (SD)	72.84+/-7.20	70.89+/-8.77	71.12+/-5.49	67.88+/-8.13	72.14+/-6.12	71.11+/-7.57
Right Included Eye N (%)	15 (48.39%)	13 (46.43%)	12 (70.59%)	6 (35.29%)	2 (28.57%)	48 (48.00%)
Patient's history						
Included eye's glaucoma	2 (6.45%)	1 (3.57%)	0 (0.00%)	2 (11.76%)	0 (0.00%)	5 (5.00%)
Diabetes	0 (0.00%)	5 (17.86%)	0 (0.00%)	3 (17.65%)	1 (14.29%)	9 (9.00%)
BCVA Mean (SD)	72.19+/-8.39	70.18+/-8.10	67.00+/-10.25	70.00+/-8.18	61.43+/-10.00	69.62+/-9.01
Eye Status						
Phakic	22 (70.97%)	18 (64.29%)	16 (94.12%)	14 (82.35%)	1 (14.29%)	71 (71.00%)
Posterior vitreous detachment	24 (80.00%)	16 (59.26%)	0 (0.00%)	6 (35.29%)	5 (71.43%)	51 (52.04%)
Intraretinal edema	10 (32.26%)	7 (25.00%)	0 (0.00%)	6 (35.29%)	2 (28.57%)	25 (25.00%)
Pseudo macular hole	4 (12.90%)	2 (7.14%)	6 (35.29%)	0 (0.00%)	1 (14.29%)	13 (13.00%)

Cystoid macular edema	3 (9.68%)	6 (21.43%)	0 (0.00%)	1 (5.88%)	0 (0.00%)	10 (10.00%)
Subretinal fluid	0 (0.00%)	2 (7.14%)	2 (11.76%)	0 (0.00%)	0 (0.00%)	4 (4.00%)
Patient Discomfort						
(moderate/severe)						
Blurry Visions	29 (93.55%)	23 (82.14%)	9 (56.25%)	14 (82.35%)	7 (100.0%)	82 (82.83%)
Metamorphopsia	13 (41.94%)	12 (42.86%)	10 (62.50%)	12 (70.59%)	4 (57.14%)	51 (51.52%)
Relative scotoma	1 (3.23%)	14 (50.00%)	0 (0.00%)	0 (0.00%)	1 (14.29%)	16 (16.16%)
Diplopia	2 (6.45%)	5 (17.86%)	1 (6.25%)	3 (17.65%)	1 (14.29%)	12 (12.12%)
Micropsia	1 (3.23%)	2 (7.14%)	0 (0.00%)	4 (23.53%)	0 (0.00%)	7 (7.07%)